

DEX-0257



BB

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C07H 21/04, C07K 1/00, C12N 1/20, C12P 21/06		A1	(11) International Publication Number: <b>WO 00/11014</b>
			(43) International Publication Date: 2 March 2000 (02.03.00)
(21) International Application Number: PCT/US99/19330			
(22) International Filing Date: 24 August 1999 (24.08.99)			
(30) Priority Data: 60/097,917 25 August 1998 (25.08.98) US 60/098,634 31 August 1998 (31.08.98) US			
(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): MOORE, Paul, A. [US/US]; 19005 Leatherbark Drive, Germantown, MD 20874 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). SHI, Yang-gu [CN/US]; 437 West Side Drive, Gaithersburg, MD 20878 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). FLORENCE, Kimberly, A. [US/US]; 12805 Atlantic Avenue, Rockville, MD 20851 (US). SOPPET, Daniel, R. [US/US]; 15050 Stillfield Place, Centreville, MD 22020 (US). LAFLEUR, David, W. [US/US]; 3142 Quesada Street, N.W., Washington, DC 20015 (US). EN-			
		(74) Agents: HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).	
		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: 49 HUMAN SECRETED PROTEINS			
(57) Abstract			
<p>The present invention relates to novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## 49 Human Secreted Proteins

### *Field of the Invention*

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and their production.

### *Background of the Invention*

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include the commercially valuable human insulin, interferon, Factor VIII, human growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of

the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent medical disorders by using secreted proteins or the genes that encode them.

5

### *Summary of the Invention*

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant and synthetic methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders and conditions related to the polypeptides and polynucleotides, and therapeutic methods for treating such disorders and conditions. The invention further relates to screening methods for identifying binding partners of the polypeptides.

15

### *Detailed Description*

#### Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide/sequences of the present invention.

30



In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released  
5 into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000  
10 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding  
15 sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid  
20 sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants  
25 of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig  
30 analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the

ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

- 5 A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42° C in a solution comprising 50% formamide, 5x SSC (750  
10 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

- Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions.
- 15 Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M  
20 NaH<sub>2</sub>PO<sub>4</sub>; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 ug/ml salmon sperm blocking DNA; followed by washes at 50°C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

- Note that variations in the above conditions may be accomplished through the  
25 inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above,  
30 due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a

complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using digo dT as a primer).

5       The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of  
10 single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA  
15 backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

      The polypeptide of the present invention can be composed of amino acids  
20 joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more  
25 detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of  
30 modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or

may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

**Polynucleotides and Polypeptides of the Invention****FEATURES OF PROTEIN ENCODED BY GENE NO: 1**

The translation product of this gene shares sequence homology with the TAP2 of *Rattus norvegicus* (See, e.g., Genbank Accession No. gi|407479) AAD15830; all  
 5 references available through this accession are hereby incorporated by reference herein.) which is thought to be important in antigen presentation in T-cells. As such, the protein product of this gene is useful for a variety of diagnostic tests for various immune system dysfunctions, and for intervention of the antigen presentation process (e.g., to enhance the immune response to vaccines and diminish the immune response  
 10 associated with autoimmune disease).

The major histocompatibility complex (Mhc) regions of mice, rats, and humans all contain a pair of related genes, TAP1 and TAP2, which encode members of a large superfamily of proteins of similar structure and function. A functional TAP1/TAP2 heterodimer is probably required for efficient presentation of antigens to  
 15 CD8(+) T cells. This heterodimer resides in the membrane of the endoplasmic reticulum, and transports peptides from the cytoplasm into the endoplasmic reticulum lumen for binding to Mhc class I molecules.

The translation product of this gene also shares sequence homology with the transport-associated proteins and ATP-binding proteins (see, e.g., Genbank accession  
 20 numbers CAB05918 (z83328.1) and AAB95060 (AF040659.1); all references available through these accessions are hereby incorporated by reference herein.)

Preferred polypeptides of the invention comprise the following amino acid sequence:

EPHRGPHLPPDLGHHHGQRPGQLQNINVFLRNTVKVTGVVVFMSLSWQLSLV  
 25 TFMGFPIIMMVSNIYGKYYKRLSKEVQNALARASNTAEETISAMKTVRSFAN  
 EEEAEVYLRKLQQVYKLNKRAAAYMYVWGSGLTLLVVQVSILYYGGH  
 LVISGQMTSGNLIAFIYEFVLGDCMENVSFSLSPGKVTALVGPSGSGKSSCVN  
 ILENFYPLEGGRVLLDGKPISAYDHYKYLHRVISLSQEPVLFARSITDNISYGLP  
 TVPFEMVVEAAQKANAHGFIMELQDGYSTETGEKGAQLSGGQKQQRVAWPG  
 30 LWCGTPQSSSWMKPPALWMPRASI (SEQ ID NO: 123);  
 MSSATWTAASWRTSATSTSLTRCWISGQPACTAAACCWGATIGVAKNSALG  
 PRRLRASWLVITLVCLFVGIYAMVKLLLFSEVRRPIRDPFWFALFVWTYISLG

ASFLWLLSTVRPGTQALEPGAATEAEGFPGSGRPPPEQASGATLQKLLSYT  
KPDVAFLVAASFFLIVAALGETFLPYTGRAIDGIVIQKSMQFSTAVVIVCLL  
AIGSSFAAGIRGGIFTLIFARLNIRLRNCLFRSLVSQETSFFDENRTGDLISRLTS  
DTTMVSDLVSRTSMSSCGTQSRRAWWSACSASHGSSPWSPSWASPS (SEQ

5 ID NO: 124) or

HLLRPAHCAFRDGGGGRTEGQCPRLLHHGTPGRLQHRDRGEGRPAVRWPEAA  
GGMARALVRNPPVLILDEATSALDAESEYLIQQAIHGNLQKHTVLIHRLST  
VEHAHLIVVLDKGRVVQQGTHQQLAQGLYAKLVQRQMLGLQPAADFTA  
GHNEPVANGSHKA (SEQ ID NO: 125). Polynucleotides encoding these

10 polypeptides are also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome 12. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 12.

15 This gene is expressed primarily in brain and testes, and to a lesser extent in amniotic cells, merkel cells and fetal tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural, reproductive, or immune disorders, particularly

20 immunodeficiency, infection, lymphomas, auto-immunities, cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in

25 certain tissues or cell types (e.g., immune, hematopoietic, neural, reproductive, developmental, differentiating, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or

30 bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 67 as residues: His-65 to Cys-73, His-144 to Gly-152. Polynucleotides encoding said polypeptides are also provided.

The major histocompatibility complex (Mhc) regions of mice, rats, and humans all contain a pair of related genes, TAP1 and TAP2, which encode members of a large superfamily of proteins of similar structure and function. A functional TAP1/TAP2 heterodimer is probably required for efficient presentation of antigens to CD8(+) T cells. Furthermore, the tissue distribution in merkel cells and homology to TAP indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of conditions in which antigen presentation is at issue, such as general microbial infection, auto-immunity, inflammation or cancer. Alternatively, expression within brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3275 of SEQ ID NO:11, b is an integer of 15 to 3289, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 2

The translation product of this gene shares sequence homology with a human retrovirus related env polyprotein pseudogene. This similarity indicates that the human protein described herein is also sent to the cell surface.

Preferred polypeptides of the invention comprise the following amino acid sequence:

RLTKTISFSLQNQTAFINSLAKTPYQALTGAALAGSYPIWENENTLSWYLPSPPTLLSPPVLFCVIQLIFXLPANWSGTCTLVFQAPTINILPPNQITILISVEASISSSPIR  
NKWALHLITLLTGLGITAALGTGIAGITTSITSYQTLFTTSLNTVEDMHTSITSL  
QR

QLDFLVGVILQNWRVLDLLTTEKGGTCIYLQECCFCVNESGIVHIAVRRLHDDRAAEL (SEQ ID NO: 126), YPIWENENTLSWYLPSPPTLLSPPVLFCV (SEQ ID NO: 127), or RVDLLTTEKGGTCIYLQECCFCVNE (SEQ ID NO: 128).

Polynucleotides encoding these polypeptides are also provided. Further,

The gene encoding the disclosed cDNA is believed to reside on the X chromosome. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for the X chromosome.

This gene is expressed primarily in human prostate cancer (stage C fraction), subtracted kidney cortex, adult brain and breast, and to a lesser extent in a variety of normal and transformed tissue types.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, anti-viral therapies, including cancer and other proliferative diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides



are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the prostate, kidney, brain and breast, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, neural, endocrine, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, seminal fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 68 as residues: Ser-28 to Ser-37, Ser-50 to Ser-58. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in various transformed tissues, combined with its similarity to retroviral env proteins indicates that the protein product is useful for the detection, diagnosis, and treatment of a variety of cancers and other proliferative disorders. This gene may show utility in gene therapy applications in viral prophylaxis. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2328 of SEQ ID NO:12, b is an integer of 15 to 2342, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 3

The translation product of this gene shares sequence homology with Cytochrome P450 monooxygenase which is thought to be important in NADPH-dependent oxidation of a number of cellular substrates (See Genbank Accession Nos. 5 gi|1185452, gb|AAC50370.1, and gb|AAB87635.1, in addition to Geneseq Accession No. R72378; all information and references available through these accessions are hereby incorporated herein by reference).

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 11 - 27 of the amino acid sequence referenced in 10 Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 28 to 501 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

Included in this invention as a preferred domain is the cytochrome P450 15 cysteine heme-iron ligand signature domain, which was identified using the ProSite analysis tool (Swiss Institute of Bioinformatics). Cytochrome P450's [1,2,3] are a group of enzymes involved in the oxidative metabolism of a high number of natural compounds (such as steroids, fatty acids, prostaglandins, leukotrienes, etc) as well as drugs, carcinogens and mutagens. Based on sequence similarities, P450's have been 20 classified into about forty different families [4,5]. P450's are proteins of 400 to 530 amino acids; the only exception is *Bacillus* BM-3 (CYP102) which is a protein of 1048 residues that contains a N-terminal P450 domain followed by a reductase domain. P450's are heme proteins. A conserved cysteine residue in the C-terminal part of P450's is involved in binding the heme iron in the fifth coordination site. From 25 a region around this residue, we developed a ten residue signature specific to P450's. The consensus pattern is as follows: [FW]-[SGNH]-x-[GD]-x-[RHPT]-x-C-[LIVMFAP]-[GAD] [C is the heme iron ligand].

Preferred polypeptides of the invention comprise the following amino acid sequence: FSLGRRHCLG (SEQ ID NO: 129). Polynucleotides encoding these 30 polypeptides are also provided.

Further preferred are polypeptides comprising the cytochrome P450 cysteine heme-iron ligand signature domain of the sequence referenced in Table for this gene,

and at least 5, 10, 15, 20, 25, 30, 50, or 75 additional contiguous amino acid residues of this referenced sequence. The additional contiguous amino acid residues is N-terminal or C-terminal to the cytochrome P450 cysteine heme-iron ligand signature domain. Alternatively, the additional contiguous amino acid residues is both N-terminal and C-terminal to the cytochrome P450 cysteine heme-iron ligand signature domain, wherein the total N- and C-terminal contiguous amino acid residues equal the specified number. The above preferred polypeptide domain is characteristic of a signature specific to cytochrome P450 cysteine heme-iron proteins. Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with cytochrome P450 cysteine heme-iron proteins. Such activities are known in the art, some of which are described elsewhere herein. The following references were reference above and are hereby incorporated by reference herein: [ 1] Nebert D.W., Gonzalez F.J. Annu. Rev. Biochem. 56:945-993(1987). [ 2] Coon M.J., Ding X., Pernecky S.J., Vaz A.D.N. FASEB J. 6:669-673(1992). [ 3] Guengerich F.P. J. Biol. Chem. 266:10019-10022(1991).[ 4] Nelson D.R., Kamataki T., Waxman D.J., Guengerich F.P., Estrabrook R.W., Feyereisen R., Gonzalez F.J., Coon M.J., Gunsalus I.C., Gotoh O., Okuda K., Nebert D.W. DNA Cell Biol. 12:1-51(1993).[ 5] Degtyarenko K.N., Archakov A.I. FEBS Lett. 332:1-8(1993).

When tested against Jurket T-cells and U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activation site) promoter element. Thus, it is likely that this gene activates T-cells and myeloid cells through the Jaks-STAT signal transduction pathway. GAS is a promoter element found upstream in many genes which are involved in the Jaks-STAT pathway. The Jaks-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

The gene encoding the disclosed cDNA is believed to reside on chromosome 11. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 11.

This gene is expressed primarily in fetal liver spleen, and to a lesser extent in lung, LNCAP prostate cell line, control synovial fibroblasts, human testes tumor, and Hodgkin's lymphoma.

Therefore, polynucleotides and polypeptides of the invention are useful as  
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hepatic or developmental diseases and/or disorders, particularly cancer, and other proliferative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential  
10 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the liver, spleen, lung, prostate, testes, and lymphatic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., liver, spleen, developmental, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, bile,  
15 seminal fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic  
20 epitopes shown in SEQ ID NO: 69 as residues: Leu-30 to Gly-38, Arg-67 to Val-72, Ser-127 to Trp-133, Gly-148 to Phe-154, Thr-171 to Phe-177, Thr-201 to Asp-206, Ser-265 to Pro-273, Glu-283 to Lys-297, Pro-346 to Lys-357, Phe-409 to Glu-418, Glu-423 to Ser-428, Leu-443 to Cys-448. Polynucleotides encoding said polypeptides are also provided.

25 The tissue distribution in fetal liver and homology to cytochrome P450 monooxygenase indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g., hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). In  
30 addition the expression in fetus would suggest a useful role for the protein product in developmental abnormalities, fetal deficiencies, pre-natal disorders and various would-healing models and/or tissue trauma.

The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions and as nutritional supplements. It may also have a very wide range of biological activities. Representative uses are described in the "Chemotaxis" and "Binding Activity" sections below, in Examples 11, 12, 13, 14, 15, 16, 18, 19, and 20, and elsewhere herein. Briefly, the protein may possess the following activities: cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g., for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g., for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g., for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g., for treating infections, tumors); hemostatic or thrombolytic activity (e.g., for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g., for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1652 of SEQ ID NO:13, b is an

integer of 15 to 1666, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 4

5 The translation product of this gene shares sequence homology with C1qR(P), the human C1q/MBL/SPA receptor that mediates enhanced phagocytosis in vitro (see, e.g., Genbank Accession number AAB53110.1 (U94333.1); all references available through this accession are hereby incorporated by reference herein. Also see, Immunity 1997 Feb;6(2):119-129).

10 Preferred polypeptides of the invention comprise one or more of the following amino acid sequences:

EHPTADRAGCSASGACYSLHHATMKRQAAEEACILRGGALSTVRAGAE LRA  
VLALLRAGPGPGXGSKDLLFWVALERRRSHCXLENEPLRGFSWLSSDPGGLE  
SDTLQWVEEPQRSCTARRWV (SEQ ID NO: 130), and/or

15 SRPPVGSSPQLEGDAMPPXRQRYLCKYQFEVLC PAPRPGAASNLSYRAPFQL  
HSAALDFSPPGTEVSALCRGQLPISVT CIADEIGARWDKLSGDVLCPCPGRYL  
RAGKCAELPNCLDDLGGFACECATGFELGKDGRSCVTS GEGQPTLGGTGVPT  
RRPPATATSPVPQRTWP IRVDEKLGETPLVPEQDNSVTSIPEIPRWGSQSTMST  
LQMSLQAESKATITPSGSVISKFNSTTSSATPQAFDSSSA VVFIFVSTAVVVLVI  
20 LTMTVLGLVKLCFHESPSSQPRKESMGPPGWRVILKPAALGSSSAHCTNNGV  
KVGDCDLRDRAEGALLAESPLGSSDA (SEQ ID NO: 131). Polynucleotides  
encoding such polypeptides are also provided.

This gene is believed to reside on chromosome 14. Therefore, polynucleotides and polypeptides related to this gene are useful in linkage analysis as markers for  
25 chromosome 14.

This gene is expressed primarily in chondrosarcoma, smooth muscle tissue, bone marrow, chondrosarcoma, fetal tissue (e.g., heart) and to a lesser extent in ovarian cancer, adult pulmonary tissues, and brain.

Therefore, polynucleotides and polypeptides of the invention are useful as  
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, chondrosarcoma, immune disorders, ovarian cancer, respiratory and

gastrointestinal disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at

5 significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the

10 disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 70 as residues: Pro-18 to Gly-30. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in bone marrow and homology to C1qR(P) indicates

15 that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation;

20 survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene

25 product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues,

30 such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus

erythematosis, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma.

Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer (e.g., ovarian, chondrosarcoma), and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation.

The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions.



Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Alternatively, the protein is useful in the detection, treatment, and/or prevention of vascular conditions, which include, but are not limited to, microvascular disease, vascular leak syndrome, aneurysm, stroke, atherosclerosis, arteriosclerosis, or embolism. For example, this gene product may represent a soluble factor produced by smooth muscle that regulates the innervation of organs or regulates the survival of neighboring neurons. Likewise, it is involved in controlling the digestive process, and such actions as peristalsis. Similarly, it is involved in controlling the vasculature in areas where smooth muscle surrounds the endothelium of blood vessels. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:14 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2013 of SEQ ID NO:14, b is an integer of 15 to 2027, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

5

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 5

The translation product of this gene shares sequence homology with human renal dipeptidase, which is a glycosyl-phosphatidylinositol- anchored ectoenzyme thought to be important in the metabolism of dihydro peptide bonds (See Genbank  
10 Accession No.bbs|148378; dbj|BAA02433.1; gb|AAB59410.1; dbj|BAA02431.1; and Geneseq Accession Nos. W29665 and R30823; all information and references available through these accessions are hereby incorporated herein by reference). Included in this invention as a preferred domain is the renal dipeptidase active site domain, which was identified using the ProSite analysis tool (Swiss Institute of  
15 Bioinformatics). Renal dipeptidase (rDP) (EC 3.4.13.19), also known as microsomal dipeptidase, is a zinc-dependent metalloenzyme which hydrolyzes a wide range of dipeptides. It is involved in renal metabolism of glutathione and its conjugates. It is a homodimeric disulfide-linked glycoprotein attached to the renal brush border microvilli membrane by a GPI-anchor. A glutamate residue has recently been shown  
20 [1] to be important for the catalytic activity of rDP. RDP seems to be evolutionary related to hypothetical proteins in the PQQ biosynthesis operons of *Acinetobacter calcoaceticus* and *Klebsiella pneumoniae*. The concensus pattern is as follows: [LIVM]-E-G-[GA]-x(2)-[LIVMF]-x(6)-L-x(3)-Y-x(2)-G-[LIVM]-R [E is the active site residue].

25 Preferred polypeptides of the invention comprise the following amino acid sequence: VEGGHSLDNSLSILRTFYMLGVR (SEQ ID NO: 138). Polynucleotides encoding these polypeptides are also provided.

Further preferred are polypeptides comprising the renal dipeptidase active site domain of the sequence referenced in Table for this gene, and at least 5, 10, 15, 20,  
30 25, 30, 50, or 75 additional contiguous amino acid residues of this referenced sequence. The additional contiguous amino acid residues is N-terminal or C- terminal to the renal dipeptidase active site domain. Alternatively, the additional contiguous

amino acid residues is both N-terminal and C-terminal to the renal dipeptidase active site domain, wherein the total N- and C-terminal contiguous amino acid residues equal the specified number. The above preferred polypeptide domain is characteristic of a signature specific to renal dipeptidase proteins. Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with renal dipeptidase proteins. Such activities are known in the art, some of which are described elsewhere herein. The following references were referenced above and are hereby incorporated herein by reference: [ 1 ] Adachi H., Katayama T., Nakazato H., Tsujimoto M. Biochim. Biophys. Acta 1163:42-48(1993) and [ 2 ] Rawlings N.D., Barrett A.J. Meth. Enzymol. 248:183-228(1995).

The polypeptide of this gene has been determined to have two transmembrane domains at about amino acid position 17 - 33 and 470 - 486 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type IIIB membrane proteins.

When tested against Jurket T-cell cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activation site) and NF-kB (Nuclear Factor kB) pathway. Thus, it is likely that this gene activates T-cells through the Jaks-STAT signal transduction pathway and is involved in the activation of apoptosis. GAS is a promoter element found upstream in many genes which are involved in the Jaks-STAT pathway. The Jaks-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. Similarly, NF-kB is a transcription factor activated by a wide variety of agents, leading to cell activation, differentiation, or apoptosis. Reporter constructs utilizing the NF-kB promoter element are used to screen supernatants for such activity.

Preferred polypeptides of the invention comprise the following amino acid sequence: VEGGHS , RYLTLTH SEQ ID NO: 133, CNTPWA SEQ ID NO: 134, APVIFSHS SEQ ID NO: 135, RNVPDD SEQ ID NO: 136, or GLEDVS SEQ ID NO: 137(SEQ ID NO:). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in infant brain, and to a lesser extent, in primary dendritic cells, L428 cells, melanocytes, keratinocytes, eosinophils, ovarian tumor, thymus stromal cells, treated bone marrow, and Hodgkins lymphoma, and (to a lesser extent) in a variety of other normal and transformed cell types.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, renal, urogenital, or neural disorders, particularly neurodegenerative and/or developmental disorders of the brain, including cancer and other proliferative  
10 disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and neurological systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types  
15 (e.g., immune, neural, renal, urogenital, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 71 as residues: Thr-36 to Arg-41, Pro-55 to Pro-60, Pro-67 to Leu-72, Asn-111 to Ser-118, Cys-138 to Asp-144, Asn-290 to Pro-296, Gly-350 to Phe-358, Gly-379 to Glu-384, Gln-399 to Cys-426, Ser-428 to Ser-438. Polynucleotides encoding said polypeptides are also provided.

25 The tissue distribution combined with its homology to the human renal dipeptidase indicates that this gene or gene product could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney  
30 stones, in addition to Wilms Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Representative uses are described here and elsewhere herein. Alternatively, the tissue distribution in

brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system.

Moreover, expression within embryonic tissue and other cellular sources marked by proliferating cells combined with the detected GAS and NF-kB biological activity indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2320 of SEQ ID NO:15, b is an integer of 15 to 2334, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 6

Preferred polypeptides of the invention comprise the following amino acid

sequence:

TWLRLGSSQIWLGTAPRGPRIHPEQAGLAGAPVKSTSSEESQPGGQCQSSGGA  
 QTLPSLRAAPVAALGSLSSYPDSCPRATTPELCPGAPTLHLADSISGPVSPPGSS  
 LGPDAWTLCAKHHQAKGMTLGT PKVLRLQPVSPCWGPKSWRVPGPFQGR  
 5 RRGESRQQGRGKRRSARSAQSPTGPESAAWPC (SEQ ID NO: 139);  
 TVATACVWAACTGCWARPPVPTWAGCAARCAAEDARAGVGDLPATGGAA  
 TGRRALTPAPRGPCILSPQPWALGLPGAPLPAALPGRARGRPGLPALPALSTL  
 PGCPALDPAGAGTLCPPPGAAEPAGP (SEQ ID NO: 140) or  
 RSGQPGECSMLRKFSLQRLLSPLDQAQTRWGLALACVAGDKGPPRPWNISAA  
 10 PAHPHVTTPGMETSGGPARDGGLILEREAAFNKPAPGE (SEQ ID NO: 141).

Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in immune cells (e.g., eosinophils, T-cells, and macrophage), leukemic and lymphoid cells, rectum, colon, and tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as  
 15 reagents for differential identification of the tissue(s) or cell type(s) present in a  
 biological sample and for diagnosis of diseases and conditions which include, but are  
 not limited to, inflammatory disorders, particularly immunodeficiency, tumor  
 necrosis, infection, lymphomas, auto-immunities, breast cancer, disorders of the colon  
 and rectum, metastasis, inflammation, anemias (leukemia) and other hematopoietic  
 20 disorders. Similarly, polypeptides and antibodies directed to these polypeptides are  
 useful in providing immunological probes for differential identification of the  
 tissue(s) or cell type(s). For a number of disorders of the above tissues or cells,  
 particularly of the immune system, expression of this gene at significantly higher or  
 lower levels is routinely detected in certain tissues or cell types (e.g., immune,  
 25 hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph,  
 serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample  
 taken from an individual having such a disorder, relative to the standard gene  
 expression level, i.e., the expression level in healthy tissue or bodily fluid from an  
 individual not having the disorder.

30 The tissue distribution indicates that polynucleotides and polypeptides  
 corresponding to this gene are useful for the diagnosis and treatment of a variety of  
 immune system disorders. Expression of this gene product in a variety of immune

cells indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity, immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The tissue distribution in kidney indicates the protein product of this gene could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilm's Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:16 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2594 of SEQ ID NO:16, b is an integer of 15 to 2608, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 7

The translation product of this gene shares sequence homology with the human IgE receptor which is thought to be important in immune regulation, particularly in immune cell aggregation (See Genbank Accession No gi|337418; all references available through this accession are hereby incorporated herein by reference). Moreover, the protein is believed to share structural features to the TM4SF superfamily of proteins.

The polypeptide of this gene has been determined to have two transmembrane domains at about amino acid position 73 - 89 and 106 - 122 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type IIIa membrane proteins. Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with human IgE receptor proteins. Such activities are known in the art, some of which are described elsewhere herein.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

RCQRNKDIMMSSKPTSHAEVNETIPNPYPSSFMAPGFQQ  
 PLGSINLENQAQGAQRAQPYGITSPGIFASSQPGQGNIQMINPSVGTAVMNFK  
 EEAKALGVIQIMVGLMHIGFGIVLCLISFSFREVLGFASTAXIGGYPFWGGLSFI  
 ISGSLSVSASKELSRCLVKGSLGMNIGRSILAFIVILLVDMCINGVXGQDYW  
 XVLSGKGISATLMIFSXLEFFVACATAHFANQANTTTNMSVLVIPNMYESNPX  
 TPASSAPPRCNNYSANAPKRKRGISLISWRKTTCKNFLRCLLLSTMISSL



(SEQ ID NO: 142). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in colon.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders and disorders of the digestive tract. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and digestive system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, gastrointestinal, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, bile, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 73 as residues: Met-2 to Ser-8, Glu-14 to Ser-23, Leu-39 to Gly-53. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution and homology to the human IgE receptor indicates that that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Expression of this gene product in colon indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of proliferative mechanisms in the digestive tract. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene has homology to a gene of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such

as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1277 of SEQ ID NO:17, b is an integer of 15 to 1291, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 8

This gene is expressed primarily in T-cells and lymph node.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly immunodeficiencies or inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph,

serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5           The tissue distribution in T-cells and lymph node indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the
- 10       expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).
- 15           Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as
- 20       T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that
- 25       influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue
- 30       markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as,

antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3115 of SEQ ID NO:18, b is an integer of 15 to 3129, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 9

This gene is expressed primarily in immune and haemopoietic cells, particularly messangial cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or haemopoietic diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and haemopoietic systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, haemopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in immune cells and hemopoietic cells indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses

are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma.

Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3615 of SEQ ID NO:19, b is an integer of 15 to 3629, where both a and b correspond to the positions of nucleotide  
 5 residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 10

The translation product of this gene shares sequence homology with various dehydrogenase and oxidoreductase polypeptides and appears to belong in the alcohol  
 10 dehydrogenase /xybitol dehydrogenase family (see, e.g., Genbank accession number AAD36790.1 (AE001811.1) and CAA68181 (X99908.1); all references available through this accession are hereby incorporated by reference herein.).

Preferred polypeptides of the invention comprise the following amino acid sequence:

15 MGRLDGKVIILTAAQIGQAAALAFAREGAKVIATDINESKLQELEKYPGIQ  
 TRVLDVTKKKQIDQFANEVERLDVLFNVAGFVHHGTVLDCEEKDWDFSMNL  
 NVRNVMYLMIKAFLPKMLAQKSGNIINMSSVASSVKGVNRCVYSTTKAAV  
 IGLTKSVAADFIQQGIRCNCVCPGTVDTPSLQERIQARGNPPEEARNDFLKRQK  
 TGRFATAEEIAMLCVYLASDESAYVTGNPVIIDGGWSL (SEQ ID NO: 143).

20 Also provided are fragments thereof having dehydrogenase activity and polypeptides comprising at least 30 residues of the foregoing amino acid sequence. Polynucleotides encoding such polypeptides are also provided.

This gene is expressed primarily in fetal tissue (e.g., liver, spleen, lung), gall  
 25 bladder, heart, bone marrow and to a lesser extent in smooth muscle, and parathyroid tumor.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, cardiovascular and developmental disorders. Similarly,  
 30 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and

fetal systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 76 as residues: Pro-78 to Gln-85, Arg-87 to Arg-94, Asp-96 to Gly-104. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in bone marrow indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,

and in the differentiation and/or proliferation of various cell types. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically



excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1130 of SEQ ID NO:20, b is an integer of 15 to 1144, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 11

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 115- 131 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 132 to 152 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

GTIGLYWVGSII MSVVVFVPGNIVGKYGTRICPAFFLSIPYTCLPVWAGFRIYN  
 QPSENYNYP SKVIQEAQAKDLLRRPFDLMLVVCLLLATGFCLFRGLIALDCPS  
 ELCRLYTQFQEPY LKDPAA YPKIQMLAYMFYSVPYFVTALYGLVVPGC SWM  
 PDITLIHAGGLAQAQFSHIGASLHARTAYVYRVPEEAKILFLALNIA YGVLPQL  
 LAYRCIYKPEFFIKTKAE EKVE (SEQ ID NO: 144). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in osteoclastoma, and to a lesser extent, in other human tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, skeletal diseases and/or disorders, particularly osteoclastoma and osteoporosis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the

tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skeletal system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., skeletal, immune, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 77 as residues: Thr-32 to Lys-40, Lys-146 to Glu-152. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in osteoclastoma indicates a role in the detection and treatment of disorders and conditions affecting the skeletal system, in particular osteoporosis as well as disorders afflicting connective tissues (e.g., arthritis, trauma, tendonitis, chondromalacia and inflammation), such as in the diagnosis or treatment of various autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as chondrodysplasias (ie. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Representative uses are described here and elsewhere herein. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 1429 of SEQ ID NO:21, b is an integer of 15 to 1443, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 12

The polypeptide of this gene has been determined to have transmembrane domains at about amino acid position 1-23 and 149-167 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type IIb membrane  
10 proteins.

Preferred polypeptides of the invention comprise the following amino acid sequence:

MSNHDPRGCTRRRAQKPLAIQPRLFHASAPDEGTQGTLKGTQKGGCILVQCQ  
SEGGAAGAWTGPPSPARDRRVRPPGTAKAQLERRRHVPRLHGLGVGGCEVR  
15 TGIVARISGSTPWAGGKPLGLHGAMGEAGAGDTGCCAKGPSAAPLPAEGRG  
QGAGPGGLVGRGERRDQQTLLGMAEDTGXSPSRPSAPAPRAPVPARQPLPRA  
RLGAATAISKSRSSRVAPALAAASASSHQR (SEQ ID NO: 145). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in neutrophils, haemopoietic cells, tymus  
20 tumor, osteosarcoma, synovial sarcoma, B-cell lymphoma, dendritic cells, pineal gland, brain, prostate and to a lesser extent in other tissues, including cancers.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are  
25 not limited to, immune and haemopoietic disorders, particularly neutropenia or neutrophilia, and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and haemopoietic systems, expression of  
30 this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, haemopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or

another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic,  
5 epitopes shown in SEQ ID NO: 78 as residues: Ser-23 to Ala-32, Gly-40 to Glu-47. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune cells (e.g., neutrophils) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses  
10 are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine  
15 production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such  
20 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus  
25 erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,  
30 and in the differentiation and/or proliferation of various cell types. Expression of this gene product in tonsils also indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell

lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses). The tissue distribution in pineal gland and brain indicates polynucleotides and

5 polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection,

10 treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder,

15 learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation,

20 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show

25 utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution in thymus tumor, B-cell lymphoma, osteosarcoma, and synovial sarcoma indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of these and related diseases.

Many polynucleotide sequences, such as EST sequences, are publicly

30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1039 of SEQ ID NO:22, b is an integer of 15 to 1053, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 13

The translation product of this gene shares sequence homology with lymphoblastic leukaemia antigen, which is thought to be important in cancers including leukemia.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 46 - 62 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 63 to 69 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

This gene is expressed primarily in lung, and infant adrenal gland.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the pulmonary and endocrine system, including cancers and developmental diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the pulmonary and endocrine systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., pulmonary, endocrine, developmental, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression

level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in adrenal gland indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various endocrine disorders and cancers. Representative uses are described in the "Biological Activity", "Hyperproliferative Disorders", and "Binding Activity" sections below, in Example 11, 17, 18, 19, 20 and 27, and elsewhere herein. Briefly, the protein can be used for the detection, treatment, and/or prevention of Addison's Disease, Cushing's Syndrome, and disorders and/or cancers of the pancreas (e.g., diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism), thyroid (e.g., hyper-, hypothyroidism), parathyroid (e.g., hyper-, hypoparathyroidism), hypothalamus, and testes.

Similarly, expression within fetal tissues and other cellular sources marked by proliferating cells, combined with the homology to the human lymphoblastic leukaemia antigen, indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:23 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 727 of SEQ ID NO:23, b is an integer of 15 to 741, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 14

This gene is expressed primarily in human tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic diseases and/or disorders, particularly leukemia and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 80 as residues: Gly-33 to Arg-40, Ser-106 to Met-112, Ala-154 to Gly-163. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in tonsils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Expression of this gene product indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen



presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory conditions such as inflammatory bowel disease, sepsis, acne, and psoriasis and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 932 of SEQ ID NO:24, b is an integer of 15 to 946, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 15**

The polypeptide of this gene has been determined to have two transmembrane domains at about amino acid position 49 - 65 and 141 - 157 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type IIIa membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

5 STXTXTIGXAGTPAGTGPEFPGRPTRPGEXPVDFSKQYSA\$WMCLSLAALA  
CSAGDTWASEVGPVLSKSSPRLITTWEKVPVGTNGGVTVVGLVSSLLGGTFV  
GIAYFLTQLIFYNDLDISAPQWPPIAFGGLAGLLGSIVDSYLGATMQYTGLDES  
TGMVVNSPTNXARHIAGKPILDNNAVNLFS\$VLIALLLPTAAWGFWPRG  
(SEQ ID NO: 146). Polynucleotides encoding these polypeptides are also provided.

10 This gene is expressed primarily in colon, brain, and to a lesser extent, in epiglottis.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are  
15 not limited to, diseases and/or disorders of the central nervous system and gastrointestinal or digestive tract. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the digestive and central nervous system, expression of  
20 this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., gastrointesinal, neural, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or  
25 bodily fluid from an individual not having the disorder.

The tissue distribution in brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11,  
30 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia,

obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. The protein product of this gene may also be useful for the detection, treatment, or prevention of a variety of gastrointestinal and digestive tract disorders, particularly proliferative disorders, such as ulcers and cancers. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 817 of SEQ ID NO:25, b is an integer of 15 to 831, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

## 25 FEATURES OF PROTEIN ENCODED BY GENE NO: 16

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 21-39 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 40-41 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

Preferred polypeptides comprise the following amino acid sequence:

## MSQRAG

RRPGGWNPSLSVVEVCRGCRGTGPLPWGASLFPCSASPLFPLPLNRRGDVHG  
TLGGRMLNRVECRDGVAAAWLCLHDAAAIRGAVGRCPMWTQPTHWVLLL  
CWALHFYCR (SEQ ID NO: 147) Also preferred are the polynucleotides encoding

5 these polypeptides.

This gene is expressed primarily in tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly tonsillitis and tonsillar neoplasms. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, saliva, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Expression of this gene product in tonsils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel

disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1280 of SEQ ID NO:26, b is an integer of 15 to 1294, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

## 25 FEATURES OF PROTEIN ENCODED BY GENE NO: 17

This gene is expressed primarily in fetal tissue (e.g., bone, liver, spleen), smooth muscle, chondrosarcoma, osteoblasts, osteosarcoma, and placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of developing and growing organs and tissues, bone disease, osteosarcoma, and other cancers. Similarly, polypeptides and antibodies directed to

these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the fetal and developing systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., fetal tissue, bone, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, bone marrow, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

10           The tissue distribution in osteosarcoma, osteoblasts, and chondrosarcoma indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of bone disease and diseases of the skeletal system. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular  
15           division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern  
20           formation.

          Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of  
25           potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types  
30           of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in

modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1642 of SEQ ID NO:27, b is an integer of 15 to 1656, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 18

This gene is expressed primarily in liver, fetal liver, and to a lesser extent in bone marrow stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hepatic dysfunction, immune disorders, and disease of the hemopoietic system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the liver and immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune; liver, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma,

urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 84 as residues: Glu-44 to Asp-50. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in liver and fetal liver indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment  
10 of liver disorders and cancers (e.g., hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). In addition the expression in fetus would suggest a useful role for the protein product in developmental abnormalities, fetal deficiencies, pre-natal disorders and various wound-healing models and/or tissue trauma. The  
15 tissue distribution in bone marrow indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product  
20 indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- 25 Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as  
30 T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus



erythematosis, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1336 of SEQ ID NO:28, b is an integer of 15 to 1350, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 19

The translation product of this gene shares sequence homology with vacuolar proton-ATPase subunit M9.2 (see, e.g., Genbank accession numbers CAA75571 (Y15286.1); all references available through this accession are hereby incorporated by reference herein.). Preferred polypeptide encoded by this gene comprise the following amino acid sequence:

MTAHSFALPVIIFTTFWGLVGIAGPWFVPKGPNRGVIITMLVATAVCCYLFWL  
IAILAQLNPLFGPQLKNETIWYVRFLWE (SEQ ID NO: 148) and  
AQRAARLGTRAPAAPAARPCILPGHPAPGHDGALIRPPGHHLHHVLGPRRHR  
GPWFVPKGPNRGVIITMLVATAVCCYLFWL IAILAQLNPLFGPQLKNETIWYV

RFLWE (SEQ ID NO: 149). Polynucleotides encoding such polypeptides are also provided.

This gene is expressed primarily in infant brain, pancreas islet cell tumor, ovary tumors, immune cells (e.g., T-cells), normal cerebellum, endometrial tumor  
5 tissues and to a lesser extent in other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodevelopmental disorders, endocrine system disorders, disorders  
10 of the immune system, and ovarian cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, ovaries, immune system, expression of this gene at significantly higher or lower levels is routinely  
15 detected in certain tissues or cell types (e.g., immune, neural, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 The tissue distribution and homology to vacuolar proton-ATPase subunit M9.2 indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neurodevelopmental disorders. The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative  
25 disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis,  
30 encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive

disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

5           Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. The tissue distribution in immune cells (e.g., T-cells) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses  
10 are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine  
15 production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

          Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such  
20 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus  
25 erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,  
30 and in the differentiation and/or proliferation of various cell types. The tissue distribution in endocrine tissues such as the pancreas indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or

prevention of various endocrine disorders and cancers. Representative uses are described in the "Biological Activity", "Hyperproliferative Disorders", and "Binding Activity" sections below, in Example 11, 17, 18, 19, 20 and 27, and elsewhere herein. Briefly, the protein can be used for the detection, treatment, and/or prevention of

5 Addison's Disease, Cushing's Syndrome, and disorders and/or cancers of the pancreas (e.g., diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism); thyroid (e.g., hyper-, hypothyroidism), parathyroid (e.g., hyper-, hypoparathyroidism), hypothalamus, and testes. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to

10 isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly

15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

20 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1752 of SEQ ID NO:29, b is an integer of 15 to 1766, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

## 25 FEATURES OF PROTEIN ENCODED BY GENE NO: 20

This gene is expressed primarily in placenta, induced endothelial cells, immune cells (e.g, T-cells, B-cells, leukocytes), brain, fetal tissue, epididymus, lung, lung cancer, thyroid tumor and to a lesser extent in many other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as

30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, fetal developmental disorders, immune disorders, cancer of the lungs,

thyroid, and cancer, in general. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at

5 significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the

10 disorder.

The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative

15 Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction,

20 aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

25 Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. The tissue distribution in immune cells (e.g., T-cells, B-cells, and leukocytes) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders.

30 Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the

proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- 5           Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as
- 10   T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that
- 15   influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates
- 20   this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell
- 25   differentiation and/or apoptosis in pattern formation.

- Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of
- 30   potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the

polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of  
5 degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation.

The tissue distribution in endothelial cells indicates that polynucleotides and  
10 polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders involving the vasculature. Elevated expression of this gene product by endothelial cells indicates that it may play vital roles in the regulation of endothelial cell function; secretion; proliferation; or angiogenesis. Alternately, this may represent a gene product expressed by the endothelium and transported to distant sites of action  
15 on a variety of target organs. Expression of this gene product by hematopoietic cells also indicates involvement in the proliferation; survival; activation; or differentiation of all blood cell lineages. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a  
20 nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
25 related to SEQ ID NO:30 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general  
30 formula of a-b, where a is any integer between 1 to 2776 of SEQ ID NO:30, b is an integer of 15 to 2790, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 21**

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 1-29 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

Preferred polypeptides comprise the following amino acid sequence:

MTLEEHRDRPRLGMCVCACVYACMLMHVCVHACLCVCVCVCVEPWSS  
R

QSKDTGGWHMEEQVTPPSLAQLKSGQVRGEMGEGRGEKGEEALTGGAEAL  
SLLGRRSPSTPLFLDREDKQAKDARNLSSTVAPDF (SEQ ID NO: 150). Also preferred are the polynucleotides encoding these polypeptides.

This gene is expressed primarily in tonsils and activated monocytes and to a lesser extent in activated neutrophils and anergic T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 87 as residues: Thr-27 to Arg-33, Gly-37 to Ser-42, Pro-52 to Arg-72. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune cells (e.g., T-cells, neutrophils) indicates polynucleotides and polypeptides corresponding to this gene are useful for the



diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Expression of this gene product in tonsils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

related to SEQ ID NO:31 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or  
5 more polynucleotides comprising a nucleotide sequence described by the general formula of  $a-b$ , where  $a$  is any integer between 1 to 1403 of SEQ ID NO:31,  $b$  is an integer of 15 to 1417, where both  $a$  and  $b$  correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where  $b$  is greater than or equal to  $a + 14$ .

## 10 FEATURES OF PROTEIN ENCODED BY GENE NO: 22

This gene is expressed primarily in CD34 positive cells (Cord Blood) and resting T-cells and to a lesser extent in anergic T-cells and neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a  
15 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at  
20 significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the  
25 disorder.

The tissue distribution in immune cells (e.g., T-cells, neutrophils) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in  
30 Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including

blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene  
5 product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues,  
10 such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits  
15 hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their  
20 interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
25 related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general  
30 formula of a-b, where a is any integer between 1 to 1892 of SEQ ID NO:32, b is an integer of 15 to 1906, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 23**

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 33 - 49 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 50 to 62 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

HEKILTPIWPSSTDLEKPHLEMLFLNVILFSLTVFTLISTAHTLDRAVRSDWLLL  
VLIYACLEELIPELIFNLYCQGNATLFF (SEQ ID NO: 151). Polynucleotides

encoding these polypeptides are also provided.

This gene is expressed primarily in activated T cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic diseases and/or disorders, particularly inflammatory conditions or immunodeficiencies such as AIDS. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in T-cells indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of

immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or, 5 activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene 10 product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, 15 such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits 20 hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their 25 interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are 30 related to SEQ ID NO:33 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 529 of SEQ ID NO:33, b is an integer of 15 to 543, where both a and b correspond to the positions of nucleotide  
5 residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 24

This gene is expressed primarily in activated T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as  
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic diseases and/or disorders, particularly inflammatory or immunodeficiency disorders, such as AIDS. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological  
15 probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal  
20 fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a  
25 variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Expression of this gene product in indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells.  
30 This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1438 of SEQ ID NO:34, b is an integer of 15 to 1452, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 25**

This gene is expressed primarily in multiple sclerosis tissue, immune cells (e.g., T-cells and dendritic cells), brain, uterus, ovary, stomach, placenta, and fetal tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, multiple sclerosis, disorders of the fetus and female reproductive system, immune disorders, particularly immunodeficiency, tumor necrosis, infection,

lymphomas, auto-immunities, cancer, metastasis, wound healing, inflammation, anemias (leukemia) and other hematopoietic disorders, in addition to developmental or proliferative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential  
5 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, developmental, reproductive system, differentiating, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid,  
10 serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic  
15 epitopes shown in SEQ ID NO: 91 as residues: Met-1 to Lys-6. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune cells (e.g., T-cells, dendritic cells) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses  
20 are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine  
25 production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such  
30 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues,



such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The tissue distribution in female reproductive organs indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating female infertility and cancers including but not limited to ovaries and uterus. The protein product is likely involved in preparation of the endometrium of implantation and could be administered either topically or orally. Alternatively, this gene could be transfected in gene-replacement treatments into the cells of the endometrium and the protein products could be produced. Similarly, these treatments could be performed during artificial insemination for the purpose of increasing the likelihood of implantation and development of a healthy embryo. In both cases this gene or its gene product could be administered at later stages of pregnancy to promote healthy development of the endometrium. The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated

expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:35 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2894 of SEQ ID NO:35, b is an integer of 15 to 2908, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 26

The gene encoding the disclosed cDNA is believed to reside on chromosome 20. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 20.

This gene is expressed primarily in fetal and developing tissues, tumors of male and female reproductive tissue (e.g., ovary and testes), and immune cells (e.g., T-cells).

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental and growth disorders, disorders of the immune system, disorders and cancers of ovaries and testes. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the fetal, reproductive, or developing systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., developing, differentiating, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene

expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 92 as residues: Val-57 to Ala-63. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune cells (e.g., T-cells) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The tissue distribution in ovaries and testes indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of diseases of the reproductive organs, including but not limited to ovarian and testicular cancer. The

expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative  
5 uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent  
10 of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the  
15 polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in  
20 modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their  
25 interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
30 related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 939 of SEQ ID NO:36, b is an integer of 15 to 953, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 27

This gene is expressed primarily in T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and haemopoietic disorders, particularly immunodeficiencies, such as AIDS, or inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and haemopoietic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, haemopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in T-cells indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3850 of SEQ ID NO:37, b is an integer of 15 to 3864, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 28

In another embodiment, polypeptides comprising the amino acid sequence of

the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

PANKAGAAIEAGIGISLMVLSPWACLFVVFPPYIQSSLRSDKHLQLSNILPTPS

- 5 HHHLPASICIQLRAGN (SEQ ID NO: 152). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in brain (early stage human brain and cerebellum) and immune cells (activated neutrophils, activated T-cells, neutrophils and dendritic cells) and, to a lesser extent in other tissues.

- 10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural or immune diseases and/or disorders, particularly inflammatory disorders, in addition to cancer and other proliferative disorders. Similarly,
- 15 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain or the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, neural,
- 20 and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, amniotic fluid, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 25 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the
- 30 detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses ,



autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system.

- 5 Alternatively, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Expression of this gene product in a variety of hematopoietic derived cells (T-cells, neutrophils, etc.) indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of
- 10 potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- Since the gene is expressed in cells of lymphoid origin, the natural gene
- 15 product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages,
- 20 and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or
- 25 immunotherapy targets for the above listed tissues.

- Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:38 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
- 30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 1397 of SEQ ID NO:38; b is an integer of 15 to 1411, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:38, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 29

When tested against Jurket T-cell cell lines, supernatants removed from cells containing this gene activated the NF-kB (Nuclear Factor kB) promoter element. Thus, it is likely that this gene activates T-cells. NF-kB is a transcription factor activated by a wide variety of agents, leading to cell activation, differentiation, or  
10 apoptosis. Reporter constructs utilizing the NF-kB promoter element are used to screen supernatants for such activity.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 5 - 21 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 22 to  
15 40 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by  
20 the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

AGSPAGTGPEFPGRPTRPISTHVFEYECICKIPRFMCEYVLLLYIVLLCNRSYA  
VFTQCVLRSSPIDSSRNAVLL (SEQ ID NO: 153). Polynucleotides encoding these polypeptides are also provided.

25 This gene is expressed primarily in colon, synovium, chondrosarcoma and to a lesser extent in dendritic cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are  
30 not limited to, immune, gastrointestinal, or skeletal diseases and/or disorders, particularly cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential

identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and digestive systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., lymph, skeletal, gastrointestinal, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or  
5 another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution of this gene product in synovium would suggest a role  
10 in the detection and treatment of disorders and conditions affecting the skeletal system, in particular osteoporosis as well as disorders afflicting connective tissues (e.g., arthritis, trauma, tendonitis, chondromalacia and inflammation), such as in the diagnosis or treatment of various autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and  
15 specific joint abnormalities as well as chondrodysplasias (ie. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid).

The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents  
20 that modulate their interactions and as nutritional supplements. It may also have a very wide range of biological activities, particularly NFk-B activation. Representative uses are described in the "Chemotaxis" and "Binding Activity" sections below, in Examples 11, 12, 13, 14, 15, 16, 18, 19, and 20, and elsewhere herein. Briefly, the protein may possess the following activities: cytokine, cell  
25 proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g., for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g., for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or  
30 nerves (e.g., for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g., for treating infections, tumors); hemostatic or thrombolytic activity (e.g., for treating

haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g., for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures.

- 5 Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 10 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:39 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
- 15 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1168 of SEQ ID NO:39, b is an integer of 15 to 1182, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

20

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 30**

The gene encoding the disclosed cDNA is believed to reside on chromosome 11. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 11.

- 25 This gene is expressed primarily in immune cells (e.g., activated T-cells), testes, fetal, spleen, and to a lesser extent in colon tumor, teratocarcinoma cells, brain and number of other tissues and cell types.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
- 30 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders related to T-cell dysfunction, such as immunodeficiencies or inflammatory conditions, in addition to neural, or

reproductive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at  
5 significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, neural, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g., seminal fluid, lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression  
10 level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 96 as residues: Pro-25 to Arg-38. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in activated T-cells and spleen indicates that  
15 polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune related disorders and diseases, including hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases; immunodeficiency diseases, such as acquired immunodeficiency syndrome,  
20 autoimmunity, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma; infections, and other inflammatory diseases and complications. Additionally, the tissue distribution in T-cells and spleen indicates polynucleotides and polypeptides corresponding to this gene are useful for the  
25 diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including  
30 blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of

degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues.

The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. In addition, the secreted protein can be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions and as nutritional supplements. It may also have a very wide range of biological activities although no evidence for any is provided in the specification. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g., for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of haematopoiesis (e.g., for treating anaemia or as adjunct to chemotherapy); stimulation of growth of bone, cartilage, tendons, ligaments and/or nerves (e.g., for treating wounds, periodontal disease, neurological diseases stroke, fibrosis); inhibition or stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g., for treating infections, tumours); haemostatic or thrombolytic activity (e.g., for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g., for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative disease; for regulation of metabolism, behaviour, and many others. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2443 of SEQ ID NO:40, b is an integer of 15 to 2457, where both a and b correspond to the positions of nucleotide  
5 residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 31

The translation product of this gene shares sequence homology with epidermal growth factor which is thought to be important in the growth and proliferation of  
10 epidermal cells, fibroblasts and number of other cell types and tissues.

The gene encoding the disclosed cDNA is believed to reside on chromosome 16. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 16.

This gene is expressed primarily in neutrophils.

15 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly autoimmunities or connective tissue diseases. Similarly, polypeptides and antibodies directed to these polypeptides are  
20 useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and connective tissue systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, epithelial, endothelial, and cancerous and wounded  
25 tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and  
30 polypeptides corresponding to this gene are useful for for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14,



16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or  
5 other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such  
10 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus  
15 erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,  
20 and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or  
25 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:41 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 1833 of SEQ ID NO:41, b is an integer of 15 to 1847, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 32

This gene is expressed primarily in kidneys, tonsils and dendritic cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, renal disorders and neoplasms, tonsillitis and immune disorders, particularly infections. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the renal and immune systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., renal, urogenital, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in kidney indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of renal conditions such as acute renal failure, kidney fibrosis and kidney tubule regeneration and neoplasms. Conditioned media, generated from the transient expression of this gene in CHO cells has been shown to activate a IFNg-reponsive element (GAS) in a B cell line (Reh). The same conditioned media had no effect on T-cell (Jurkat) and pro-monocyte (U937) derived cell lines, suggesting that the protein product of this gene may exhibit IFNg-like activity in a (B)cell-specific manner. This experimental data in conjunction with expression on dendritic cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies (e.g., AIDS), immuno-suppressive

conditions (transplantation) and hematopoietic disorders. In addition this gene product is applicable in conditions of general microbial infection, inflammation or cancer.

Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents  
5 that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
10 related to SEQ ID NO:42 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general  
15 formula of a-b, where a is any integer between 1 to 2583 of SEQ ID NO:42, b is an integer of 15 to 2597, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 33

20 The translation product of this gene was shown to have homology to the classII MHC transactivator CIITA of Mus musculus (See, e.g., Genbank Accession No gi|1870520 and AAB48859.1; all references available through this accession are hereby incorporated by reference herein.), which is thought to regulate MHC class II  
25 gene expression in B lymphocytes via direct interaction with the MHC class II-specific transcription factors. Furthermore, the CIITA protein is thought to play an indirect role in reducing tumorigenicity and inducing long-term tumor immunity.

Preferred polypeptides of the invention comprise the following amino acid sequence:

30 MP SGMSAAVPISGLLDLSHNSISQESALYLLETLPSCPRVREASVNLGSEQSF  
RIHFSREDQAGKTLRLSECSFRPEHVSRLATGLSKSLQLTELTLTQCCLGQKQL  
AILLSLVGRPAGLFSRLRVQEPWADRARVLSLLEVCAQASGSVTEISISSETQQQL  
CVQLEFPRQEENPEAVALRLAHCDLGAHHSLLXGQLMETCARLXQLSLSQV

NLCEDDDASSLLLQSLLLSLSELKTFRLTSSCVSTEGLAHLASGLGHCHHLEEL  
 DLSNNQFDEEGTKALMRALEGKWMLKRLDLSHLLLNSSTLALLTHRLSQMT  
 CLQSLRLNRNSIGDVGCCHLSEALRAATSLEELDLSHNQIGDAGVQHLATILP  
 GLPELRKIDLSGNSISSAGGVQLAESLVLCCRLEELMLGCNALGDP TALGLAQ  
 5 ELPQHRLRVLHLPFSLHGPGGALSLARPWMDPPIWKRSAWRKTTWLEGSCVS  
 VWSSRCSDR (SEQ ID NO: 154); or  
 HQLSRGSAVGRVSRSLQAPGGVDAWLQCPGGSHSPGAGSGAAPAPEGPTPTI  
 QPSGPRWGPEPGQALDGSPHLEEISLAENNLAGGVLRFCMELPLLRLQIDL VSC  
 KIDNQTA KLLTSSFTSCPALEVILLSWNLLGDEAAAE LAQVLPQMGR LKRVD  
 10 LE  
 KNQITALGAWLLAEGLAQGSSIQVIRLWNNPIPCDMAQHLSQEPRLDFAFFD  
 NQPQAPWGT (SEQ ID NO: 155). Polynucleotides encoding these polypeptides are  
 also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome  
 15 16. Accordingly, polynucleotides related to this invention are useful as a marker in  
 linkage analysis for chromosome 16.

The polypeptide of this gene has been determined to have transmembrane  
 domains at about amino acid positions 1-47 and 24-47 of the amino acid sequence  
 referenced in Table 1 for this gene. Based upon these characteristics, it is believed  
 20 that the protein product of this gene shares structural features to type IIIa membrane  
 proteins.

This gene is expressed primarily in immune cells (e.g., eosinophils, T-cells,  
 dendritic) and other cell types of hematopoietic origin and to a lesser extent in ovary  
 tumor and heart.

25 Therefore, polynucleotides and polypeptides of the invention are useful as  
 reagents for differential identification of the tissue(s) or cell type(s) present in a  
 biological sample and for diagnosis of diseases and conditions which include, but are  
 not limited to, immune disorders, particularly immunodeficiency, tumor necrosis,  
 infection, lymphomas, auto-immunities, cancer, metastasis, inflammation, anemias  
 30 (leukemia) and other hematopoietic disorders. Similarly, polypeptides and antibodies  
 directed to these polypeptides are useful in providing immunological probes for  
 differential identification of the tissue(s) or cell type(s). For a number of disorders of

the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another  
5 tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in immune cells (T-cells, dendritic, and eosinophils) indicates polynucleotides and polypeptides corresponding to this gene are useful for  
10 the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including  
15 blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for  
20 immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders,  
25 such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful  
30 in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue

markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:43 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is  
10 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3102 of SEQ ID NO:43, b is an integer of 15 to 3116, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.

15

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 34

Polynucleotides of the invention do not comprise the nucleic acid sequence shown as Genbank accession no.gb|Z85986|HS108K11, which is hereby incorporated herein by reference.

- 20 The gene encoding the disclosed cDNA is believed to reside on chromosome 6. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 6.

This gene is expressed primarily in smooth muscle and cells of hematopoietic origin.

- 25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, vascular disorders, particularly heart disease, vasculitis, atherosclerosis, in addition to immune disorders, such as immunodeficiency, auto-  
30 immunities, cancer, metastasis, anemias (leukemia) and other hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the

tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the [insert system where a related disease state is likely, e.g., immune], expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, endothelial, muscle, vascular, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue expression in hematopoietic tissues indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,

and in the differentiation and/or proliferation of various cell types. In addition, the expression in smooth muscle might indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of cardiovascular and disorders such as atherosclerosis, restenosis, stoke, angina, thrombosis

5 hypertension, inflammation and vascular wound healing. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or

10 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:44 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

15 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3446 of SEQ ID NO:44, b is an integer of 15 to 3460, where both a and b correspond to the positions of nucleotide

20 residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 35

The translation product of this gene was shown to have homology to the human cytochrome c-like polypeptide from lung adenocarcinoma A549 (See

25 Genbank Accession No.bbs|175350) which is thought to be involved in metabolic processes, specific to tumors or transformed cells.

Preferred polypeptides of the invention comprise the following amino acid sequence:

EKLFCFEMLLICKFSPNSVPPETCAILNQGLMDLGLCRMCLGNNMFAGSMLG

30 KSHRHSPFSINQRHNALRKAAGTPAQKSLGIVQVS PN SEQ ID NO: 156,  
GCAGCALVTICLQAVCLVKAIALHSRLTRDTMHCGRPQGGLPRKAWVLSRF  
PPTETA (SEQ ID NO: 157);



PETQCTAEGRRDPCPEKPGYCPGFPQLRQPEIWPRGKGKTLHPPARHM (SEQ ID NO: 158); SEIGENRP (SEQ ID NO: 159); HDTDSFAH (SEQ ID NO: 160); or ALRKAAG (SEQ ID NO: 161). Polynucleotides encoding these polypeptides are also provided. Polynucleotides of the invention do not comprise the nucleic acid sequence shown as Genbank accession no.gb|AC004706|AC004706, which is hereby  
5 incorporated herein by reference.

This gene is expressed primarily in fetal tissue, lung, melanocyte, retina, brain, T-cell lymphoma, and to a lesser extent, in other tissues and cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as  
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hepatic, pulmonary, developmental, or growth disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell  
15 type(s). For a number of disorders of the above tissues or cells, particularly of the immune and pulmonary systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, pulmonary, hematopoietic, developing, differentiating, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid, serum, plasma, urine, synovial  
20 fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of  
25 neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome,  
30 meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive

compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

- 5           Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. The tissue distribution in T-cell lymphoma indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the
- 10 "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or
- 15 other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such
- 20 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus
- 25 erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,
- 30 and in the differentiation and/or proliferation of various cell types. Moreover, the expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may

show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions  
5 involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of  
10 potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types  
15 of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new  
20 insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or  
25 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 2608 of SEQ ID NO:45, b is an integer of 15 to 2622, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 36

The translation product of this gene was shown to have homology to the human protein kinase C substrate 80K-H (See Genbank Accession No. P14314), which is important in the regulation of various signal transduction pathways. Polynucleotides of the invention do not comprise the nucleic acid sequence shown as  
10 Genbank accession no. gb|G21007|G21007, which is hereby incorporated herein by reference.

This gene is expressed primarily in breast and prostate cancer, retina, ovary, parathyroid tumor, fetal tissue and to a lesser extent in ovary.

Therefore, polynucleotides and polypeptides of the invention are useful as  
15 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive, endocrine, or ocular disorders, particular breast or other cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the  
20 tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive and endocrine systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, endocrine, differentiating, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, breast milk, aqueous humor, vitreous humor, serum,  
25 plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic  
30 epitopes shown in SEQ ID NO: 102 as residues: Gln-14 to Val-20, Arg-28 to Trp-35. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in breast cancer cell lines, combined with the homology to a PKC substrate indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis, treatment, and/or prevention of a variety of tumors, particularly of the breast or other neoplasms. Alternatively, the tissue distribution in ovary and parathyroid tumors indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various endocrine disorders and cancers, particularly Addison's Disease, Cushing's Syndrome, and disorders and/or cancers of the pancreas (e.g., diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism), thyroid (e.g., hyper-, hypothyroidism), parathyroid (e.g., hyper-, hypoparathyroidism), hypothalamus, and testes. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in

modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution in immune cells (e.g., germinal B-cells) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the

protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution in parathyroid tumor, prostate cancer and breast cancer tissue indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment these and related disorders.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:46 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1970 of SEQ ID NO:46, b is an integer of 15 to 1984, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.

## 20 FEATURES OF PROTEIN ENCODED BY GENE NO: 37

Preferred polypeptides of the invention comprise the following amino acid sequence:

MRGPVCGFSLVEMLLALALGLMLILGVTQIALSSRTTYASQSAASLLQDDAR  
 FALGKLIQEIRQAGMFGCLSAASISNAPAGFDRPIGWSTTGSSRSLTLVTADVG  
 25 EGGSKPDWTVLSDCTGSAHAYVGSPPAANARANPLPTCAKLT (SEQ ID NO:  
 162); or  
 MGYYSRQRQAGMVLLISLVFLLLLALLGVSSMQGAISQEKITGSLRQRNQSF  
 QQAESGLRLGESLVQASGFALRPCHSTAACAPPAESVSVVGPGTNPVSTVTWI  
 GMKDGVIYGIQNLGPGTGLVNSRQRPRPSIA (SEQ ID NO: 163).

30 Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in cord blood, ovary, tumors of the parathyroid, testes, and pancreas, and to a lesser extent in fetal tissue, retina, brain, colon, endometrial stromal, HL-60 cells, and many other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as  
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hematopoietic, or developmental disorders, cancer of the ovaries and endocrine system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential  
10 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic and immune systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, reproductive, developmental, endocrine, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid,  
15 serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic  
20 epitopes shown in SEQ ID NO: 103 as residues: Lys-29 to Ser-38, Ser-55 to Trp-61, Gln-63 to Ser-69. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in endocrine tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various endocrine disorders and cancers, particularly Addison's Disease,  
25 Cushing's Syndrome, and disorders and/or cancers of the pancreas (e.g., diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism), thyroid (e.g., hyper-, hypothyroidism), parathyroid (e.g., hyper-, hypoparathyroidism), hypothalamus, and testes. The tissue distribution in immune cells (e.g., neutrophils and T-cells) indicates polynucleotides and polypeptides corresponding to this gene are  
30 useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein.



Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may

also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation.

10       The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, 15 the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive 20 compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, 25 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show 30 utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

related to SEQ ID NO:47 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or  
5 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1973 of SEQ ID NO:47, b is an integer of 15 to 1987, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.

#### 10 FEATURES OF PROTEIN ENCODED BY GENE NO: 38

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 1-24 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

15 This gene is expressed primarily in activated monocytes, dendritic cells, and cancerous ovary.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are  
20 not limited to, ovarian cancer, immune or hematopoietic disorders, particularly immunodeficiencies or inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene  
25 at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily  
30 fluid from an individual not having the disorder.

The tissue distribution in monocytes and dendritic cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the

treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. The tissue distribution in dendritic cells indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The tissue distribution in ovarian cancer tissue indicates that polynucleotides and polypeptides

corresponding to this gene are useful for the treatment and diagnosis of ovarian, as well as, other cancers:

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2099 of SEQ ID NO:48, b is an integer of 15 to 2113, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 39

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 36-57 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type II membrane proteins.

Preferred polypeptides of the invention comprise the following amino acid sequence:

ENESTKEPSLLQYLCVQSPAGLNGFNVLLSGSQTPPTVGPSSGQLPSFSVPCM  
VLPSPLGPFVLYSPAMGPVSSTLGALPNTGPVNFSLPGLGSIAQLLVGPTA  
VVNPKSSTLPSADPQLQSQPSLNLSPVMSRSHSVVQQPESPVYVGHVSVVKL  
HQSPVPVTPKSIQRTHRETFKTPGSLGDPVLKRRERNNHETPARPRGD (SEQ  
ID NO: 164). Polynucleotides encoding such polypeptides are also provided.

This gene is expressed primarily in 8 week-old embryo, fetal lung, testes, colon, the tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or

activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

5           Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as  
10   T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that  
15   influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. endothelium, and T-cell lymphoma.

20           Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive and immune defects, cancer, T-cell lymphoma, and developmental disorders. Similarly, polypeptides and antibodies directed to these  
25   polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the pulmonary and immune systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma,  
30   urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e.,

the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in T-cell lymphoma indicates that polynucleotides and polypeptides corresponding to this gene are useful for study and treatment of developmental and pulmonary defects and neoplasms of blood, reproductive and other organs. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The tissue distribution in testes tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of male reproductive and endocrine disorders. It may also prove to be valuable in the

diagnosis and treatment of testicular cancer, as well as cancers of other tissues where expression has been observed. The expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment,  
5 and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

10       Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have  
15 applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue  
20 differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the  
25 protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30       Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:49 and may have been publicly available prior to conception of



the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3451 of SEQ ID NO:49, b is an integer of 15 to 3465, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 40

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 145-171 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 168-282 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

This gene is expressed primarily in T helper cells, breast cancer, kidney, fetal tissue and to a lesser extent in thymus and cells from some other hemopoietic and endocrine sources.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and inflammatory conditions, cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic and lymphoid systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 106 as residues: Pro-135 to Ile-145, Trp-173 to Gly-188, Pro-199 to Gln-219, Ser-225 to Ala-237, Pro-240 to Gly-253, Ser-262 to Gly-275. Polynucleotides encoding said polypeptides are also provided.

- 5       The tissue distribution in breast cancer tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neoplasms of breast and other organs. The tissue distribution in immune cells (e.g., T-helper cells) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders.
- 10       Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation
- 15       of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such
- 20       as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus
- 25       erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,
- 30       and in the differentiation and/or proliferation of various cell types. The tissue distribution in kidney indicates the protein product of this gene could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal

tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilm's Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Moreover, the expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1223 of SEQ ID NO:50, b is an integer of 15 to 1237, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 41

The gene encoding the disclosed cDNA is believed to reside on chromosome 16. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 16.

This gene is expressed primarily in adipose tissue, brain, Hodgkin's lymphoma, and to a lesser extent in fetal tissue colon tumor, synovium, salivary gland, immune cells (e.g., neutrophils), and other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disorders, immune, or metabolic disorders, particularly diseases or disorders of adipose tissue or Hodgkin's lymphoma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the adipose tissues and immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, adipose, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e.,

the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 107 as residues: Arg-54 to Leu-60, Ala-73 to Gly-78.

5 Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in adipose tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of obesity, or other metabolic disorders, such as Tay-Sach's Disease, phenylketonuria, galactosemia, hyperlipidemias, porphyrias, and Hurler's syndrome, in addition to  
10 various immune disorders and neoplasia. The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example  
15 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms,  
20 hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

25 Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. The tissue distribution in immune cells (e.g., neutrophils) indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses  
30 are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation;

survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

- 5           Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as
- 10   T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that
- 15   influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. In addition, the expression of this gene product in synovium would suggest a role in the detection and
- 20   treatment of disorders and conditions afflicting the skeletal system, in particular osteoporosis, bone cancer, connective tissue disorders (e.g., arthritis, trauma, tendonitis, chondromalacia and inflammation). The protein is also useful in the diagnosis or treatment of various autoimmune disorders (i.e., rheumatoid arthritis, lupus, scleroderma, and dermatomyositis), dwarfism, spinal deformation, joint
- 25   abnormalities, and chondrodysplasias (i.e. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid, etc.). Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional
- 30   supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:51 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1383 of SEQ ID NO:51, b is an integer of 15 to 1397, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 42

Polynucleotides of the invention do not comprise the nucleic acid sequence shown as Genbank accession no.gb|G15452|G15452, which is hereby incorporated herein by reference.

Preferred polypeptides of the invention comprise the following amino acid sequence:

RHERHEYRRALDHEEEALSSGSVQEA EAM LDEPQEQAEGSLTVYVISEHSSLL  
PQDMMSYIGPKRTAVVRGIMHREAFNIIGRRIVQVAQAMSLTEDVLAAALAD  
HLPEDKWSAEKRRPLKSSLGYEITFSLNPDPKSHDVYWDIEGAVRRYVQPFL  
NALGAAGNFSVDSQILYYAMLGVNPRFDSASSSYLDMHSLPHVINPVESRL  
GSSAASLYPVLNFLLYVPELAHSPLYIQDKDGAPVATNAFHSPRWGGIMVYN  
VDSKTYNASVLPVRVEVDMVRVMEVFLAQLRLLFGIAQPQLPPKCLLSGPTS  
EGLMTWELDRLLWARSVENLATATTTLTSLAQLLGKISNIVIKDDVASEVYK  
AVAAVQKSAEELASGHLASAFVASQEAVTSSSELAFFDPSLLHLLYFPDDQKF  
AIYIPLFLPMAVPILLSLVKIFLETRKSWRKPEKTD (SEQ ID NO: 165).

Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in immune and haemopoietic cells, tumors of the ovaries, endometrium, and parathyroid.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, immune or haemopoietic disorders, particularly cancers, ovarian cancer, cancers of the endocrine system and endometrium, and disorders of the retina. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the haemopoietic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, haemopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 108 as residues: Phe-11 to Gly-16, Pro-33 to Ser-42. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in T-cells and bone marrow indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders,



such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Moreover, the expression within fetal tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as,

antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:52 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2257 of SEQ ID NO:52, b is an integer of 15 to 2271, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 43

The gene encoding the disclosed cDNA is believed to reside on chromosome 19. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 19.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 25-44 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

This gene is expressed primarily in infant brain, fetal heart, uterine cancer, colon, metastatic melanoma, spleen, liver, thymus and other cancers.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of developing tissues, haemopoietic or immune system, cardiovascular or musculoskeletal, or neural tissues, uterine cancer and metastatic melanoma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system and developing systems, expression of this gene at

significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, muscle, immune, hematopoietic, hepatic, developing, differentiating, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, bile, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample  
5 taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in infant brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of  
10 neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia,  
15 obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system.  
20 Alternatively, expression within embryonic tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis in pattern formation. Thus  
25 this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy.

The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions and as nutritional supplements. It may also have a  
30 very wide range of biological activities. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g., for treating human

immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g., for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g., for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g., for treating infections, tumors); hemostatic or thrombolytic activity (e.g., for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g., for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. The tissue distribution in uterine cancer and metastatic melanoma indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of these cancers.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2755 of SEQ ID NO:53, b is an integer of 15 to 2769, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

## 25 FEATURES OF PROTEIN ENCODED BY GENE NO: 44

This gene shares sequence homology with Serglycin (see GenSeq accession number Q44278; all references available through this accession are hereby incorporated by reference herein.) Serglycin is involved in the regulation of haematopoietic cell function and development.

30 This gene is expressed primarily in brain frontal cortex.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological and cognitive conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system (CNS), expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 110 as residues: Tyr-28 to Cys-40. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for study and treatment of central nervous system disorders, esp. schizophrenia, neurodegenerative and memory disorders. The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to  
5 identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
10 related to SEQ ID NO:54 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general  
15 formula of a-b, where a is any integer between 1 to 1375 of SEQ ID NO:54, b is an integer of 15 to 1389, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 45**

20 This gene is expressed primarily in human stomach.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, gastrointestinal disorders, particularly gastritis, stomach ulcers, and  
25 stomach cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the gastrointestinal system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g.,  
30 gastrointesinal, endothelial, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, bile, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard

gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in stomach cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating or diagnosing disease involving the stomach such as inflammation, ulceration or cancers. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:55 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 734 of SEQ ID NO:55, b is an integer of 15 to 748, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 46

The gene encoding the disclosed cDNA is believed to reside on chromosome 11. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 11.

This gene is expressed primarily in brain and fetal liver, and to a lesser extent, in other cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental, neurological, behavioral, hepatic or immune diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides

are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune or nervous systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g.,  
5 hepatic, immune, hematopoietic, neural, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, bile, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

10 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 112 as residues: Ala-24 to Lys-31. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution of this gene predominantly in fetal liver indicates that the gene could be important for the treatment or detection of immune or  
15 hematopoietic disorders including arthritis, asthma, immunodeficiency diseases and leukemia. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of neurodegenerative disease states and behavioural  
20 disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional  
25 supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of  
30 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or



more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 4188 of SEQ ID NO:56, b is an integer of 15 to 4202, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

5

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 47

Preferred polypeptides of the invention comprise the following amino acid sequence:

KLLLTKEQKLELARLQVDTSGSKEFGTSGIPAKCRFPKIFVNTDDTYEELHLI

10 VYKVTTVFLPAL (SEQ ID NO: 166). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in brain and lymph node of breast cancer.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, breast cancer and neural disorders, particular neurodegenerative, neurological, or psychological disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neurological and psychological disorders, including but not limited to: trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, toxic neuropathies induced by neurotoxins, inflammatory diseases such as meningitis and encephalitis, demyelinating diseases, neurodegenerative diseases

30

such as Parkinson's Disease, Huntington's Disease, Alzheimer's Disease, peripheral neuropathies, multiple sclerosis, neoplasia of neuroectodermal origin, etc.

In addition, the secreted protein can be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to  
5 identify agents that modulate their interactions and as nutritional supplements. It may also have a very wide range of biological activities although no evidence for any is provided in the specification. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g., for treating human  
10 immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of haematopoiesis (e.g., for treating anaemia or as adjunct to chemotherapy); stimulation of growth of bone, cartilage, tendons, ligaments and/or nerves (e.g., for treating wounds, periodontal disease, neurological diseases stroke, fibrosis); inhibition or stimulation of follicle stimulating hormone (for control of  
15 fertility); chemotactic and chemokinetic activities (e.g., for treating infections, tumours); haemostatic or thrombolytic activity (e.g., for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g., for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative disease; for regulation of metabolism, behaviour, and many others. Also contemplated is the  
20 use of the corresponding nucleic acid in gene therapy procedures.

The tissue distribution in cancerous breast tissue indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of breast cancer, and cancer in general. Furthermore, the protein may also be used to determine biological activity, to raise  
25 antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly  
30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:57 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 840 of SEQ ID NO:57, b is an integer of 15 to 854, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 48

A preferred polypeptide fragment of the invention comprises the following amino acid sequence:

MEPQLGPEAAALRPGWLALLLWVSALSCSFSLPASSLSSLVPQVRTSYNFGRT  
FLGLDKCNACIGTSICKKFFKERNKI (SEQ ID NO: 167). Polynucleotides encoding these polypeptides are also provided.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

QLPLWSPASVQPRVDSQRARGSPPEPKMEPQLGPEAAALRPGWLALLLWVS  
ALSCSFSLPASSLSSLVPQVRTSYNFGRTFLGLDKCNACIGTSICKKFFKEEIRS  
DNWLASHLGLPPDSLLSYPANYSDDSKIWRPVEIFRLVSKYQNEISDRKICAS  
ASAPKTCSIERVLRKTERFQKWLQAKRLTPDLVQDC HQGQRELKFLCMLR  
(SEQ ID NO: 168). Polynucleotides encoding these polypeptides are also provided.

A preferred polypeptide variant of the invention comprises the following amino acid sequence:

MEPQLGPEAAALRPGWLALLLWVSALSCSFSLPASSLSSLVPQVRTSYNFGRT  
FLGLDKCNACIGTSICKKFFKEEIRSDNWLASHLGTASRFPLXSYPCLLQMX  
KIWXPCGXLLTGQ QXSNEISKQEIXCLLHPPPKNLHIDV (SEQ ID NO: 169).  
Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in endothelial cells, and to a lesser extent, in the adult pulmonary system.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, vascular and pulmonary diseases and/or disorders, particularly atherosclerosis, and microvascular disease. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the pulmonary and vascular systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., endothelial, pulmonary, cardiovascular, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, pulmonary surfactant, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 114 as residues: Arg-45 to Gly-51, Glu-75 to Asn-81, Ala-99 to Ile-107, Lys-119 to Asp-126, Leu-145 to Gln-152. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution of this gene in the pulmonary system, and in particular endothelial cells, indicates that it could play a role in the treatment/detection of lung lymphoma or sarcoma formation, pulmonary edema and embolism, bronchitis and cystic fibrosis. Expression in endothelial cells suggest a role in the treatment and/or detection of vascular disorders including vasculitis, cardiovascular disorders such as myocardial infarction, myocarditis, ischemia and stroke. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:58 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1441 of SEQ ID NO:58, b is an integer of 15 to 1455, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 49

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

GPRARVQGFSGADIVKFMALGSMYLVLTIVAKVLRGAEPCCGPLKNRVLRP  
CPLPVHCPLPIPSAEGIPWVAYLPIRWFISCCPGHCIQIPMCTS (SEQ ID NO:

170). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in kidney, and to a lesser extent, in a wide variety of human tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, renal or urogenital diseases and/or disorders, particularly kidney cancer or nephritis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the urological or renal system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., renal, urogenital, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in kidney indicates that this gene or gene product could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilms Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Representative uses are described here and elsewhere herein. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:59 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 579 of SEQ ID NO:59, b is an integer of 15 to 593, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:59, and where b is greater than or equal to a + 14.

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HKGCN17	203105 08/13/98	pSport1	11	3289	1	3289	218	218	67	1	23	24	155
2	HETAD68	203081 07/30/98	Uni-ZAP XR	12	2342	1	2342	84	84	68	1	32	33	69
3	HPIAT78	203081 07/30/98	Uni-ZAP XR	13	1666	1	1666	30	30	69	1	29	30	501
4	HMWGY65	203105 08/13/98	Uni-ZAP XR	14	2027	1	1976	42	42	70	1	21	22	188
5	HDTAB58	203081 07/30/98	pCMVSPORT 2.0	15	2334	1874	2334	133	133	71	1	33	34	486
5	HDTAB58	203081 07/30/98	pCMVSPORT 2.0	60	496	159	496	223	223	116	1	37	38	48
6	HEOMQ62	203105 08/13/98	pSport1	16	2608	1	2608	130	130	72	1	24	25	87
7	HWLJQ88	203081 07/30/98	pSport1	17	1291	1	1291	114	114	73	1	30	31	299
7	HWLJQ88	203081 07/30/98	pSport1	61	1292	1	1292	403	403	117	1	25	26	203
8	HMICP03	203105 08/13/98	Uni-ZAP XR	18	3129	1	3129	86	86	74	1	34	35	47

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
9	HABAB01	203105 08/13/98	pCMVSPORT 3.0	19	3629	1	3629	147	147	75	1	15	16	43
10	HE2AT09	203105 08/13/98	Uni-ZAP XR	20	1144	437	1144	435	435	76	1	17	18	140
11	HSDJA15	203081 07/30/98	Uni-ZAP XR	21	1443	1	1443	247	247	77	1	20	21	152
12	HAMGW29	203105 08/13/98	pCMVSPORT 3.0	22	1053	12	1053	25	25	78	1	27	28	179
13	HAPSR85	203081 07/30/98	Uni-ZAP XR	23	741	1	741	428	428	79	1	21	22	69
14	HTOHD42	203081 07/30/98	Uni-ZAP XR	24	946	1	946	155	155	80	1	24	25	190
15	HWLJH65	203081 07/30/98	pSport1	25	831	1	831	129	129	81	1	18	19	165
16	HTOJA73	203105 08/13/98	Uni-ZAP XR	26	1294	1	1294	100	100	82	1	21	22	41
17	HPMGI45	203105 08/13/98	Uni-ZAP XR	27	1656	1	1656	119	119	83	1	25	26	48
18	HFVIC62	203105 08/13/98	pBluescript	28	1350	1	1350	114	114	84	1	31	32	56
19	HHENW77	203105 08/13/98	pCMVSPORT 3.0	29	1766	1	1714	205	205	85	1	19	20	42



Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
20	HMSIV91	203105 08/13/98	Uni-ZAP XR	30	2790	1	2790	95	95	86	1	19	20	40
21	HMSKC04	203105 08/13/98	Uni-ZAP XR	31	1417	1	1417	133	133	87	1	22	23	73
22	HSZG33	203105 08/13/98	Uni-ZAP XR	32	1906	1	1906	122	122	88	1	23	24	46
23	HTEBC92	203081 07/30/98	Uni-ZAP XR	33	543	1	543	63	63	89	1	22	23	62
23	HTEBC92	209215 08/21/97	Uni-ZAP XR	62	398	241	398		156	118	1	13	14	18
24	HTXEL29	203081 07/30/98	Uni-ZAP XR	34	1452	1	1452	322	322	90	1	25	26	69
24	HTXEL29	209090 06/05/97	Uni-ZAP XR	63	1202	1	1202		294	119	1			12
25	HDPWA44	203105 08/13/98	pCMVSPORT 3.0	35	2908	1	2908	40	40	91	1	30	31	56
26	HMACS20	203105 08/13/98	Uni-ZAP XR	36	953	1	953	227	227	92	1	43	44	67
27	HAJAY88	203105 08/13/98	pCMVSPORT 3.0	37	3864	1	3864	201	201	93	1	25	26	44
28	HBOEG69	203081 07/30/98	pSport1	38	1411	1	1411	302	302	94	1	19	20	54

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of -Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
29	HWLEQ37	203081 07/30/98	pSport1	39	1182	1	1182	110	110	95	1	19	20	40
30	HE9CS37	203105 08/13/98	Uni-ZAP XR	40	2457	65	2454	183	183	96	1	22	23	40
31	HNGEI34	203105 08/13/98	Uni-ZAP XR	41	1847	1	1847	70	70	97	1	18	19	49
32	HTOAT76	203105 08/13/98	Uni-ZAP XR	42	2597	1	2597	100	100	98	1	41	42	57
33	HDPVH60	203105 08/13/98	pCMVSPORT 3.0	43	3116	1	3100	8	8	99	1	45	46	51
34	HL YCR65	203105 08/13/98	pSport1	44	3460	1	3460	28	28	100	1	19	20	44
35	HARAY91	203105 08/13/98	pBluescript SK-	45	2622	1	2622	214	214	101	1	46	47	57
36	HCHNT03	203105 08/13/98	pSport1	46	1984	133	1984	228	228	102	1	17	18	40
37	HCUBW95	203105 08/13/98	ZAP Express	47	1987	1	1987	131	131	103	1	20	21	142
38	HDPLV95	203105 08/13/98	pCMVSPORT 3.0	48	2113	1	2113	12	12	104	1	18	19	43
39	HEMGB12	203105 08/13/98	Uni-ZAP XR	49	3465	1	3438	78	78	105	1	48	49	62

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
40	HHENP27	203105 08/13/98	pCMVSPORT 3.0	50	1237	1	1237	12	12	106	1	22	23	282
41	HSPBF70	203105 08/13/98	pSport1	51	1397	288	1397	429	429	107	1	19	20	97
42	HTXKB57	203105 08/13/98	Uni-ZAP XR	52	2271	1	2271	290	290	108	1	16	17	61
43	HUKAA55	203105 08/13/98	Lambda ZAP II	53	2769	129	2769	263	263	109	1	32	33	46
44	HFXGT58	203105 08/13/98	Lambda ZAP II	54	1389	12	1389	238	238	110	1	28	29	46
45	HROAS46	203081 07/30/98	Uni-ZAP XR	55	748	1	748	123	123	111	1	19	20	41
46	HUSFF19	203081 07/30/98	pBluescript	56	4202	863	2447	1080	1080	112	1	20	21	41
47	HBWBX21	203105 08/13/98	ZAP Express	57	854	1	854	201	201	113	1	17	18	43
48	HUVDI43	203081 07/30/98	Uni-ZAP XR	58	1455	103	1455	128	128	114	1	31	32	182
48	HUVDI43	203081 07/30/98	Uni-ZAP XR	64	1517	1	1517	133	133	120	1	31	32	79
48	HUVDI43	203027 06/26/97	Uni-ZAP XR	65	526	69	526	89	89	121	1	31	32	146

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
49	HTLCU49	203081 07/30/98	Uni-ZAP XR	59	593	1	593	170	170	115	1	20	21	80
49	HTLCU49	203027 06/26/97	Uni-ZAP XR	66	664	1	664	249	249	122	1	20	21	80

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing) and the translated SEQ ID NO:Y (where Y may be any of the polypeptide sequences disclosed in the sequence listing) are sufficiently

accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also  
5 hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used, for example, to generate antibodies which bind specifically to proteins containing the polypeptides and the secreted proteins encoded by the cDNA clones identified in Table 1.

10 Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the  
15 actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the  
20 generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods.  
25 The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

30 The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed

herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or a deposited clone, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the secreted protein.

The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or a cDNA contained in ATCC deposit Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide encoded by the cDNA contained in ATCC deposit Z. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of the polypeptide sequence of SEQ ID NO:Y' and/or a polypeptide sequence encoded by the cDNA contained in ATCC deposit Z are also encompassed by the invention.

#### Signal Sequences

The present invention also encompasses mature forms of the polypeptide having the polypeptide sequence of SEQ ID NO:Y and/or the polypeptide sequence encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species of the protein. Further, it has long been known that cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, Virus Res. 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, Nucleic Acids Res. 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of



these methods is in the range of 75-80%. (von Heinje, supra.) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., Protein Engineering 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

### **Polynucleotide and Polypeptide Variants**

The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X, the complementary strand thereto, and/or the cDNA sequence contained in the deposited clone.

The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y and/or encoded by the deposited clone.

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence contained in a deposited cDNA clone or the complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited clone, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein). Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

The present invention is also directed to polypeptides which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% identical to, for example, the polypeptide sequence shown in SEQ ID NO:Y, the polypeptide sequence encoded by the cDNA contained in a deposited clone, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described herein).

By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In

other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using

the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, an amino acid sequences shown in Table 1 (SEQ ID NO:Y) or to the amino acid sequence encoded by cDNA contained in a deposited clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid-residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., J. Biol. Chem. 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions,

inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies  
5 for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions  
10 where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function.  
15 For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains  
25 are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr,  
30 and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues,



where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for  
5 example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of  
10 charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev.  
15 Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid  
20 substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1  
25 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

30

#### **Polynucleotide and Polypeptide Fragments**

The present invention is also directed to polynucleotide fragments of the polynucleotides of the invention.

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence which: is a portion of that contained in a deposited clone, or encoding the polypeptide encoded by the cDNA in a deposited clone; is a portion of that shown in SEQ ID NO:X or the complementary strand thereto, or is a portion of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:Y. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in a deposited clone or the nucleotide sequence shown in SEQ ID NO:X. In this context "about" includes the particularly recited value, a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X, or the complementary strand thereto, or the cDNA contained in a deposited clone. In this context "about" includes the particularly recited ranges, and ranges larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to these nucleic acid molecules under

stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, and ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are

specifically contemplated by the present invention. Moreover, polynucleotides encoding these domains are also contemplated.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

#### 10 Epitopes & Antibodies

The present invention is also directed to polypeptide fragments comprising, or alternatively consisting of, an epitope of the polypeptide sequence shown in SEQ ID NO:Y, or the polypeptide sequence encoded by the cDNA contained in a deposited clone. Polynucleotides encoding these epitopes (such as, for example, the sequence disclosed in SEQ ID NO:X) are also encompassed by the invention, is the nucleotide sequences of the complementary strand of the polynucleotides encoding these epitopes. And polynucleotides which hybridize to the complementary strand under stringent hybridization conditions or lower stringency conditions.

In the present invention, "epitopes" refer to polypeptide fragments having antigenic or immunogenic activity in an animal, especially in a human. A preferred embodiment of the present invention relates to a polypeptide fragment comprising an epitope, as well as the polynucleotide encoding this fragment. A region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." In contrast, an "immunogenic epitope" is defined as a part of a protein that elicits an antibody response. (See, for instance, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998-4002 (1983).)

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

30 In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 15, at least 20, at least 25, and most preferably between about 15 to about 30

amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. (See, 5 for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983).)

Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; 10 and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985).) A preferred immunogenic epitope includes the secreted protein. The immunogenic epitopes may be presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids 15 have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting.)

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, *in vivo* immunization, *in vitro* immunization, and phage display methods. See, e.g., 20 Sutcliffe et al., supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If *in vivo* immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling of the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to 25 a carrier using a linker such as -maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier-coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier 30 protein and Freund's adjuvant. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide

adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

5           As one of skill in the art will appreciate, and discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to heterologous polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, any combination thereof  
10 including both entire domains and portions thereof) resulting in chimeric polypeptides. These fusion proteins facilitate purification, and show an increased half-life *in vivo*. This has been shown, *e.g.*, for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, *e.g.*, EPA  
15 0,394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion can also be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, *e.g.*, Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of  
20 interest as an epitope tag to aid in detection and purification of the expressed polypeptide.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to  
25 modulate the activities of polypeptides corresponding to SEQ ID NO:Y thereby effectively generating agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P.A., et al., *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S., *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, L.O., et al., *J. Mol. Biol.* 287:265-76  
30 (1999); and Lorenzo, M. M. and Blasco, R., *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and

corresponding polypeptides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired molecule corresponding to SEQ ID NO:X polynucleotides of the invention by homologous, or site-specific, recombination. In another embodiment, polynucleotides corresponding to SEQ ID NO:X and corresponding polypeptides may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of coding polynucleotide corresponding to SEQ ID NO:X, or the polypeptide encoded thereby may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

### Antibodies

The present invention further relates to antibodies and T-cell antigen receptors (TCR) which specifically bind the polypeptides of the present invention. The antibodies of the present invention include IgG (including IgG1, IgG2, IgG3, and IgG4), IgA (including IgA1 and IgA2), IgD, IgE, or IgM, and IgY. As used herein, the term "antibody" (Ab) is meant to include whole antibodies, including single-chain whole antibodies, and antigen-binding fragments thereof. Most preferably the antibodies are human antigen binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')<sub>2</sub>, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V<sub>L</sub> or V<sub>H</sub> domain. The antibodies may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, rabbit, goat, guinea pig, camel, horse, or chicken.

Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entire or partial of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are any combinations of variable region(s) and hinge region, CH1, CH2, and CH3 domains. The present invention further includes monoclonal, polyclonal, chimeric, humanized, and human monoclonal and human polyclonal antibodies which specifically bind the polypeptides of the present invention. The present invention

further includes antibodies which are anti-idiotypic to the antibodies of the present invention.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for  
5 different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for heterologous compositions, such as a heterologous polypeptide or solid support material. *See, e.g.*, WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); US Patents 5,573,920, 4,474,893, 5,601,819, 4,714,681, 4,925,648; Kostelny et al., J.  
10 Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which are recognized or specifically bound by the antibody. The epitope(s) or polypeptide portion(s) may be specified as described herein, *e.g.*, by N-terminal and C-terminal  
15 positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

20 Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of the polypeptides of the present invention are included. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and  
25 less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. Further included in the present invention are antibodies which only bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein).

30 Antibodies of the present invention may also be described or specified in terms of their binding affinity. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-6}M$ ,  $10^{-6}M$ ,  $5 \times 10^{-7}M$ ,  $10^{-7}M$ ,  $5 \times 10^{-8}M$ ,  $10^{-8}M$ ,  $5 \times 10^{-9}M$ .



$9M$ ,  $10^9M$ ,  $5 \times 10^{-10}M$ ,  $10^{-10}M$ ,  $5 \times 10^{-11}M$ ,  $10^{-11}M$ ,  $5 \times 10^{-12}M$ ,  $10^{-12}M$ ,  $5 \times 10^{-13}M$ ,  $10^{-13}M$ ,  $5 \times 10^{-14}M$ ,  $10^{-14}M$ ,  $5 \times 10^{-15}M$ , and  $10^{-15}M$ .

Antibodies of the present invention have uses that include, but are not limited to, methods known in the art to purify, detect, and target the polypeptides of the present invention including both *in vitro* and *in vivo* diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., ANTIBODIES: A LABORATORY MANUAL, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference in the entirety).

The antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, or toxins. See, e.g., WO 92/08495; WO 91/14438; WO 89/12624; US Patent 5,314,995; and EP 0 396 387.

The antibodies of the present invention may be prepared by any suitable method known in the art. For example, a polypeptide of the present invention or an antigenic fragment thereof can be administered to an animal in order to induce the production of sera containing polyclonal antibodies. The term "monoclonal antibody" is not a limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technology.

Hybridoma techniques include those known in the art and taught in Harlow et al., ANTIBODIES: A LABORATORY MANUAL, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: MONOCLONAL ANTIBODIES AND

T-CELL HYBRIDOMAS 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). Fab and F(ab')<sub>2</sub> fragments may be produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments).

- 5 Alternatively, antibodies of the present invention can be produced through the application of recombinant DNA and phage display technology or through synthetic chemistry using methods known in the art. For example, the antibodies of the present invention can be prepared using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of a phage particle which carries polynucleotide sequences encoding them. Phage with a desired binding property are selected from a repertoire or combinatorial antibody library (e.g. human or murine) by selecting directly with antigen, typically antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT/GB91/01134; WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and US Patents 5,698,426, 5,223,409, 5,403,484, 5,580,717, 5,427,908, 5,750,753, 5,821,047, 5,571,698, 5,427,908, 5,516,637, 5,780,225, 5,658,727 and 5,733,743 (said references incorporated by reference in their entireties).

- As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host including mammalian cells, insect cells, plant cells, yeast, and bacteria. For example, techniques to recombinantly produce Fab, Fab' and F(ab')<sub>2</sub> fragments can also be employed using methods known in the art such as those disclosed in WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and

Sawai et al., AJRI 34:26-34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu, L. et al., PNAS 90:7995-7999 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; and US Patent 5,807,715. Antibodies can be humanized using a variety of techniques including CDR-grafting (EP 0 239 400; WO 91/09967; US Patent 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan E.A., Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (US Patent 5,565,332). Human antibodies can be made by a variety of methods known in the art including phage display methods described above. See also, US Patents 4,444,887, 4,716,111, 5,545,806, and 5,814,318; and WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741 (said references incorporated by reference in their entireties).

Further included in the present invention are antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide of the present invention. The antibodies may be specific for antigens other than polypeptides of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either *in vitro* or *in vivo*, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in *in vitro* immunoassays and purification methods using methods known in the art. See e.g., Harbor et al. *supra* and WO 93/21232; EP 0 439 095; Naramura et al., Immunol. Lett. 39:91-99 (1994); US Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992);

Fell et al., J. Immunol. 146:2446-2452 (1991) (said references incorporated by reference in their entireties).

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides of the present invention may be fused or conjugated to the above antibody portions to increase the *in vivo* half life of the polypeptides or for use in immunoassays using methods known in the art. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See e.g., US Patents 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, 5,112,946; EP 0 307 434, EP 0 367 166; WO 96/04388, WO 91/06570; Ashkenazi et al., PNAS 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., PNAS 89:11337-11341 (1992) (said references incorporated by reference in their entireties).

The invention further relates to antibodies which act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Included are both receptor-specific antibodies and ligand-specific antibodies. Included are receptor-specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. Also include are receptor-specific antibodies which both prevent ligand binding and receptor activation. Likewise, included are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor

activation, but do not prevent the ligand from binding the receptor. Further included are antibodies which activate the receptor. These antibodies may act as agonists for either all or less than all of the biological activities affected by ligand-mediated receptor activation. The antibodies may be specified as agonists or antagonists for biological activities comprising specific activities disclosed herein. The above antibody agonists can be made using methods known in the art. See e.g., WO 96/40281; US Patent 5,811,097; Deng et al., Blood 92(6):1981-1988 (1998); Chen, et al., Cancer Res. 58(16):3668-3678 (1998); Harrop et al., J. Immunol. 161(4):1786-1794 (1998); Zhu et al., Cancer Res. 58(15):3209-3214 (1998); Yoon, et al., J. Immunol. 160(7):3170-3179 (1998); Prat et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard et al., Cytokine 9(4):233-241 (1997); Carlson et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (said references incorporated by reference in their entireties).

As discussed above, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

### 30 Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a

second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for

example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

### **Vectors, Host Cells, and Protein Production**

The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the

vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac  
5 promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a  
10 translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin  
15 resistance genes for culturing in E. coli and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as E. coli, Streptomyces and Salmonella typhimurium cells; fungal cells, such as yeast cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture  
20 mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech,  
25 Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated  
30 transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that the



polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control

regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al.,  
5 Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, Proteins: Structures and  
10 Molecular Principles, W.H. Freeman & Co., N.Y., and Hunkapiller et al., Nature, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a polypeptide sequence of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid  
analogues can be introduced as a substitution or addition into the polypeptide sequence.  
15 Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid,  $\alpha$ -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid,  $\gamma$ -Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine,  
norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-  
20 butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine,  $\beta$ -alanine, fluoro-amino acids, designer amino acids such as  $\beta$ -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptides which are differentially modified  
25 during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin,  
30 papain, V8 protease,  $\text{NaBH}_4$ ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and  
5 addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the  
10 polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent NO: 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose,  
15 dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between  
20 about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or  
25 lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those  
30 skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., Exp. Hematol. 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may

be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, pharmaceutical compositions) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone (including fragments, variants, splice variants, and fusion proteins, corresponding to these polypeptides as described herein). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (*e.g.*, containing polypeptides having identical or different amino acid sequences) or a homotrimer (*e.g.*, containing polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (*i.e.*, polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid

residues contained in the polypeptide sequence (e.g., that recited in the sequence listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant

fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

5 Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by  
10 reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, associations  
15 proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers  
20 of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues  
25 located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers  
30 containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the

polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

#### Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be



selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an  
5 amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day, using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping  
10 strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread.  
15 This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to  
20 mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

25 Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins  
30 University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural  
5 alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this  
10 polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic  
15 marker.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Both methods rely on binding of the polynucleotide to DNA or RNA. For these techniques, preferred polynucleotides are usually 20 to 40 bases in length and complementary to  
20 either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991) ) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally  
25 results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

Polynucleotides of the present invention are also useful in gene therapy. One  
30 goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate

manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of  
5 restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification  
10 difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for  
15 amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

20 Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic  
25 biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for  
30 forensic purposes.

There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with

tissue, of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

#### Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell. Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99\text{m}}\text{Tc}$ ), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers

for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

5 A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example,  $^{131}\text{I}$ ,  $^{112}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ ), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety  
10 needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99\text{m}}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics  
15 of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells  
20 or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder.

Moreover, polypeptides of the present invention can be used to treat disease.  
25 For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B), to inhibit the activity of a polypeptide (e.g., an oncogene), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the  
30 activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

#### Gene Therapy Methods

Another aspect of the present invention is to gene therapy methods for treating disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the invention that operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Beldegrun et al., J. Natl. Cancer Inst., 85:207-216 (1993); Ferrantini et al., Cancer Research, 53:107-1112 (1993); Ferrantini et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura et al., Cancer Research 50: 5102-5106 (1990); Santodonato, et al., Human Gene Therapy 7:1-10 (1996); Santodonato, et al., Gene Therapy 4:1246-1255

(1997); and Zhang, et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

5 As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

10 In one embodiment, the polynucleotide of the invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the  
15 invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

The polynucleotide vector constructs of the invention used in the gene  
20 therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be  
25 readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of polynucleotide sequence of the invention. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the  
30 respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters,

such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotides of the invention.

Unlike other gene therapy techniques, one major advantage of introducing  
5 naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the invention can be delivered to the interstitial  
10 space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic  
15 fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by  
20 injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

25 For the naked acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue  
30 site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.



The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA  
5 constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These  
10 delivery methods are known in the art.

The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

In certain embodiments, the polynucleotide constructs of the invention are  
15 complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate  
20 intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA , 86:6077-6081 (1989), which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem., 265:10189-10192 (1990), which is herein incorporated by reference), in functional  
25 form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA ,  
30 84:7413-7416 (1987), which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication NO: WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., Felgner et al., Proc. Natl. Acad. Sci. USA, 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., Methods of Immunology, 101:512-527 (1983),

which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include  $\text{Ca}^{2+}$ -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta*, 394:483 (1975); Wilson et al., *Cell*, 17:77 (1979)); ether injection (Deamer et al., *Biochim. Biophys. Acta*, 443:629 (1976); Ostro et al., *Biochem. Biophys. Res. Commun.*, 76:836 (1977); Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348 (1979)); detergent dialysis (Enoch et al., *Proc. Natl. Acad. Sci. USA*, 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., *J. Biol. Chem.*, 255:10431 (1980); Szoka et al., *Proc. Natl. Acad. Sci. USA*, 75:145 (1978); Schaefer-Ridder et al., *Science*, 215:166 (1982)), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent NO: 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding polypeptides of the invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy, 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and  $\text{CaPO}_4$  precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding polypeptides of the invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express polypeptides of the invention.

In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotides of the invention contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses polypeptides of the invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al., Am. Rev. Respir. Dis., 109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld et al., Science, 252:431-434 (1991));

Rosenfeld et al., Cell, 68:143-155 (1992)). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al. Proc. Natl. Acad. Sci. USA, 76:6606 (1979)).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel., 3:499-503 (1993); Rosenfeld et al., Cell, 68:143-155 (1992); Engelhardt et al., Human Genet. Ther., 4:759-769 (1993); Yang et al., Nature Genet., 7:362-369 (1994); Wilson et al., Nature, 365:691-692 (1993); and U.S. Patent NO: 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express E1a and E1b, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, for example, the HARP promoter of the present invention, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, Curr. Topics in Microbiol. Immunol., 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct containing polynucleotides of the invention is inserted into the AAV vector using standard cloning methods, such as those found  
5 in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses.  
10 Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct of the invention. These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express the desired gene product.

15 Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding the polypeptide sequence of interest) via homologous recombination (see, e.g., U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO  
20 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the  
25 art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous  
30 polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

The promoter and the targeting sequences can be amplified using PCR.

Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

The polynucleotides encoding polypeptides of the present invention may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

Preferably, the polynucleotide encoding a polypeptide of the invention contains a secretory signal sequence that facilitates secretion of the protein.

Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be

homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more  
5 molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications  
10 during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers. (Kaneda et al., Science, 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a  
15 recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide  
20 construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

Therapeutic compositions useful in systemic administration, include  
25 recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection,  
30 aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad.



Sci. USA , 189:11277-11281 (1992), which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian. Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly

#### Biological Activities

The polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists could be used to treat the associated disease.

#### Immune Activity

The polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called

hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins),  
5 or infectious. Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating or detecting deficiencies or disorders of  
10 hematopoietic cells. A polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to:  
15 blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia,  
20 or hemoglobinuria.

Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotides or polypeptides, or agonists or  
25 antagonists of the present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used  
30 to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of a polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat  
5 inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)); ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over  
10 production of cytokines (e.g., TNF or IL-1.)

#### Hyperproliferative Disorders

A polynucleotides or polypeptides, or agonists or antagonists of the invention can be used to treat or detect hyperproliferative disorders, including neoplasms. A  
15 polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an  
20 immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.  
25

Examples of hyperproliferative disorders that can be treated or detected by a polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast,  
30 digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and

peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by a polynucleotides or polypeptides, or agonists or antagonists of the present invention.

5 Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

10

### Cardiovascular Disorders

Polynucleotides or polypeptides, or agonists or antagonists of the invention may be used to treat cardiovascular disorders, including peripheral artery disease, such as limb ischemia.

15 Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilogy of Fallot, ventricular heart septal defects.

25 Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium,

30

postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, 5 bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular 10 rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, 15 mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, 20 pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary 25 vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, angiodysplasia, angiomas, bacillary angiomas, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive 30 diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia,

peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, atacia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

Polynucleotides or polypeptides, or agonists or antagonists of the invention, are especially effective for the treatment of critical limb ischemia and coronary disease. As shown in the Examples, administration of polynucleotides and polypeptides of the invention to an experimentally induced ischemia rabbit hindlimb  
5 may restore blood pressure ratio, blood flow, angiographic score, and capillary density.

Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle  
10 accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppository solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides of the invention may be administered as part of a pharmaceutical composition, described in more detail below. Methods of delivering polynucleotides  
15 of the invention are described in more detail herein.

#### Anti-Angiogenesis Activity

The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate.  
20 Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing  
25 solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman  
30 *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J.*



*Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun,  
5 *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides or polypeptides, or agonists or antagonists of the invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or  
10 agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)).

Ocular disorders associated with neovascularization which can be treated with  
15 the polynucleotides or polypeptides or agonists or antagonists of the invention include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalm.* 85:704-710  
20 (1978) and Gartner *et al.*, *Surv. Ophthalm.* 22:291-312 (1978).

Additionally, disorders which can be treated with the polynucleotides and polypeptides of the present invention (including agonist and/or antagonists) include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic  
25 plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

Moreover, disorders and/or states, which can be treated with be treated with polynucleotides or polypeptides or agonists or antagonists of the present invention,  
30 but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis,

psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion  
5 fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization  
10 required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (Rochele minalia quintosa), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

15       **Diseases at the Cellular Level**

Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by the polynucleotides or polypeptides and/or antagonists or agonists of the invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors,  
20 including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders  
25 (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred  
30 embodiments, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by the polynucleotides or polypeptides, or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic

injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestosis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

5

### Wound Healing and Epithelial Cell Proliferation

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, for therapeutic purposes, for example, to stimulate  
10 epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. Polynucleotides or polypeptides, as well as agonists or antagonists of the invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye  
15 tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and  
20 antimetabolites. Polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote dermal reestablishment subsequent to dermal loss

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to increase the adherence of skin grafts to a wound bed and  
25 to stimulate re-epithelialization from the wound bed. The following are a non-exhaustive list of grafts that polynucleotides or polypeptides, agonists or antagonists of the invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepdermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft,  
30 epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omentop graft, patch graft, pedicle graft, penetrating graft, split skin

graft, thick split graft. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, can be used to promote skin strength and to improve the appearance of aged skin.

It is believed that the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may have a cytoprotective effect on the small intestine mucosa. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly.

Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat diseases associated with the under expression of the polynucleotides of the invention.

Moreover, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to prevent and heal damage to the lungs due to various pathological states. A growth factor such as the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of alveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using the polynucleotides or polypeptides, and/or agonists or antagonists of the invention. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to stimulate the proliferation and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

In addition, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

#### 10 Infectious Disease

A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polypeptide or polynucleotide and/or agonist or antagonist of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections

(e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. A polypeptide or polynucleotide, and/or agonist or antagonist of the present invention  
5 can be used to treat or detect any of these symptoms or diseases.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (e.g.,  
10 Corynebacterium, Mycobacterium, Norcardia), Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (Klebsiella, Salmonella, Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales,  
15 Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Pasteurellacea Infections (e.g., Actinobacillus, Haemophilus, Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis,  
20 uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis,  
25 Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat or detect any of these symptoms or diseases.

30 Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families: Amebiasis,



Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas. These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat or detect any of these symptoms or diseases.

Preferably, treatment using a polypeptide or polynucleotide and/or agonist or antagonist of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

### **Regeneration**

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage.

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds  
5 includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a polynucleotide or polypeptide and/or agonist or antagonist of the present invention to proliferate and differentiate nerve cells. Diseases that could be treated using this  
10 method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g.,  
15 Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide and/or agonist or antagonist of the present invention.

### Chemotaxis

20 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then  
25 fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted  
30 to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured

location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may inhibit chemotactic activity. These molecules  
5 could also be used to treat disorders. Thus, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention could be used as an inhibitor of chemotaxis.

### Binding Activity

10 A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or  
15 small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural  
20 receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell  
25 membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

30 The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving

competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product  
5 mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

Preferably, an ELISA assay can measure polypeptide level or activity in a  
10 sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

Additionally, the receptor to which a polypeptide of the invention binds can be identified by numerous methods known to those of skill in the art, for example, ligand  
15 panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided  
20 into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

Following fixation and incubation, the slides are subjected to auto-  
25 radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled polypeptides  
30 can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the

polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

5           Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of polypeptides of the invention thereby effectively generating agonists and antagonists of polypeptides of the invention. See generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and  
10   5,837,458, and Patten, P. A., et al., Curr. Opin Biotechnol. 8:724-33 (1997); Harayama, S. Trends Biotechnol. 16(2):76-82 (1998); Hansson, L. O., et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. Biotechniques 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding  
15   polypeptides of the invention may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired polynucleotide sequence of the invention molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides of the invention may be altered by being subjected to random mutagenesis by error-prone  
20   PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptides of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are  
25   family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B,  
30   decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the polypeptides of the invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide. The biological activity of the fragments may include an improved desired activity, or a  
5 decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and 3[H]  
10 thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the uptake of 3[H] thymidine in each case. The amount of fibroblast cell proliferation is  
15 measured by liquid scintillation chromatography which measures the incorporation of 3[H] thymidine. Both agonist and antagonist compounds may be identified by this procedure.

In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled  
20 polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is  
25 measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring  
30 about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from

suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to the polypeptides of the invention comprising the steps of: (a) incubating a candidate binding compound with the polypeptide; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with the polypeptide, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Also, one could identify molecules bind a polypeptide of the invention experimentally by using the beta-pleated sheet regions contained in the polypeptide sequence of the protein. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, the amino acid sequence of each beta pleated sheet regions in a disclosed polypeptide sequence. Additional embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, any combination or all of contained in the polypeptide sequences of the invention. Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the amino acid sequence of each of the beta pleated sheet regions in one of the polypeptide sequences of the invention. Additional embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions in one of the polypeptide sequences of the invention.

### **Drug Screening**

Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in

such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed  
5 cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the present  
10 invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound  
15 form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present invention, and is described in great detail in European Patent Application 84/03564,  
20 published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art.  
25 Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present  
30 invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of



any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

#### Antisense And Ribozyme (Antagonists)

5 In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained a deposited clone. In one embodiment, antisense sequence is generated internally by the organism, in another embodiment, the antisense sequence is separately administered  
10 (see, for example, O'Connor, Neurochem., 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense  
15 Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research, 6:3073 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

20 For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense  
25 RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the  
30 invention. Such a vector would contain a sequence encoding the antisense nucleic acid of the invention. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA.

Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding a polypeptide of the invention, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature*, 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell*, 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature*, 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of interest. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids of the invention, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA sequence of the invention it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, *e.g.*, the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., *Nature*, 372:333-335 (1994). Thus, oligonucleotides complementary to either the 5' - or 3' - non-translated, non-coding regions of a polynucleotide sequence of the invention could be used in an antisense approach to inhibit translation of endogenous mRNA.

Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5' -, 3' - or coding region of mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556 (1989); Lemaitre et al., Proc. Natl. Acad. Sci., 84:648-652 (1987); PCT Publication NO: WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication NO: WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Król et al., BioTechniques, 6:958-976 (1988)) or intercalating agents. (See, e.g., Zon, Pharm. Res., 5:539-549 (1988)). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxymethylaminomethyl-2-thiouracil, beta-D-mannosylqueosine,

5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v),  
5 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

10 In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

15 In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., Nucl. Acids Res., 15:6625-6641 (1987)). The oligonucleotide is a 2-O-methylribonucleotide (Inoue et al., Nucl. Acids Res.,  
20 15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett. 215:327-330 (1987)).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples,  
25 phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (Nucl. Acids Res., 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., Proc. Natl. Acad. Sci. U.S.A., 85:7448-7451 (1988)), etc.

While antisense nucleotides complementary to the coding region sequence of  
30 the invention could be used, those complementary to the transcribed untranslated region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science, 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs  
5 corresponding to the polynucleotides of the invention, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5' -UG-3' . The construction and production of hammerhead ribozymes is well  
10 known in the art and is described more fully in Haseloff and Gerlach, Nature, 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within each nucleotide sequence disclosed in the sequence listing. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA corresponding to the polynucleotides of the invention; i.e., to  
15 increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the polynucleotides of the invention *in vivo*.  
20 DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of  
25 the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic  
30 cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirous in cases such as restenosis after balloon angioplasty.

The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat the diseases described herein.

#### Other Activities

The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

The polypeptide of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. The polypeptide of the invention may have the ability to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

The polypeptide of the invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte

growth. Along the same lines, the polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

The polypeptide of the invention may also be employed to maintain organs  
5 before transplantation or for supporting cell culture of primary tissues.

The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

The polypeptide or polynucleotides and/or agonist or antagonists of the  
10 present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue,  
15 pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, polypeptides or polynucleotides and/or agonist or antagonists of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

Polypeptide or polynucleotides and/or agonist or antagonists of the present  
20 invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

25 Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

### 30 Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

5 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

10 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

15 Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

20 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

25 Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

30 A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.



A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under  
5 stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which  
10 comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide  
15 sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of  
20 at least 50 contiguous nucleotides is included in the nucleotide sequence of the complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

25 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule  
30 comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least

one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of

positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence  
5 at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

10 Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino  
15 acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted  
20 protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the  
25 amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in  
30 the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred, is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA

clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of

the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

### Examples

#### Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited</u>
	<u>Plasmid</u>	
	Lambda Zap	pBluescript (pBS)
30	Uni-Zap XR	pBluescript (pBS)
	Zap Express	pBK
	lafmid BA	plafmid BA



pSport1	pSport1
pCMVSPORT 2.0	pCMVSPORT 2.0
pCMVSPORT 3.0	pCMVSPORT 3.0
pCR <sup>®</sup> 2.1	pCR <sup>®</sup> 2.1

- 5 Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are
- 10 commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS. The S and K refers to the orientation of the polylinker to the T7 and T3
- 15 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the f1 origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the f1 ori generates sense strand DNA and in the other, antisense.
- 20 Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University,
- 25 NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR<sup>®</sup>2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al.,
- 30 Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with  $^{32}\text{P}$ - $\gamma$ -ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25  $\mu\text{l}$  of reaction

mixture with 0.5 ug of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl<sub>2</sub>, 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific

to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

5     **Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide**

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

10    **Example 3: Tissue Distribution of Polypeptide**

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P<sup>32</sup> using the rediprime™ DNA labeling  
15    system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA  
20    expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to  
25    manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70°C overnight, and the films developed  
30    according to standard procedures.

**Example 4: Chromosomal Mapping of the Polynucleotides**

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This  
30    primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated 32 times followed by one 5 minute cycle at 70°C. Human, mouse, and

hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

#### **Example 5: Bacterial Expression of a Polypeptide**

A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp<sup>r</sup>), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan<sup>r</sup>). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.<sup>600</sup>) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1

mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The

origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

#### **Example 6: Purification of a Polypeptide from an Inclusion Body**

The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4°C without  
5 mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 µm membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive  
10 Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4  
15 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using  
20 a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant  $A_{280}$  monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the  
25 above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 µg of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL  
assays.

30



**Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System**

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector, contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("Geneclean" BIO 101 Inc., La Jolla, Ca.).

5        The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by  
10       gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five µg of a plasmid containing the polynucleotide is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method  
15       described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One µg of BaculoGold™ virus DNA and 5 µg of the plasmid are mixed in a sterile well of a microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 µl Lipofectin plus 90 µl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the  
20       transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

25       After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell  
30       culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then

resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

- 5           To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and
- 10           cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 µCi of <sup>35</sup>S-methionine and 5 µCi <sup>35</sup>S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).
- 15           Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

#### **Example 8: Expression of a Polypeptide in Mammalian Cells**

- 20           The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient
- 25           transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

- 30           Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109),

pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

5 Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing  
10 cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker  
15 is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the  
20 production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the  
25 CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

30 Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be  
5 modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

10 The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

15 Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five µg of the expression plasmid pC6 a pC4 is cotransfected with 0.5 µg of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are  
20 seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50  
25 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 µM, 2 µM, 5 µM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 µM. Expression of the desired gene product is analyzed,  
30 for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

### **Example 9: Protein Fusions**

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see  
5 also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create  
10 chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

15 Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion  
20 can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will  
25 not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

30

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAAACACACATGCCACCGTGC  
CCAGCACCTGAATTTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTACATGCGTGGT  
GGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG  
5 ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA  
CAAÇAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACT  
GGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCA  
ACCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC  
CACAGGTGTACACCCTGCCCCCATCCCGGGATGAGGTGACCAAGAACCAG  
10 GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGT  
GGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCT  
CCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTAÇAGCAAGCTCACCGTG  
GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCA  
TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG  
15 GTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

**Example 10: Production of an Antibody from a Polypeptide**

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells  
20 expressing a polypeptide of the present invention is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

25 In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell  
30 Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in

any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56 degrees C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

5           The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as  
10           described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

            Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method  
15           makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody  
20           whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

            It will be appreciated that Fab and F(ab')<sub>2</sub> and other fragments of the  
25           antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic  
30           chemistry.

            For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced



using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

#### **Example 11: Production Of Secreted Protein For High-Throughput Screening**

##### **Assays**

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at  $2 \times 10^5$  cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45

minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degrees C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl<sub>2</sub> (anhyd); 0.00130 mg/L CuSO<sub>4</sub>·5H<sub>2</sub>O; 0.050 mg/L of Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O; 0.417 mg/L of FeSO<sub>4</sub>·7H<sub>2</sub>O; 311.80 mg/L of KCl; 28.64 mg/L of MgCl<sub>2</sub>; 48.84 mg/L of MgSO<sub>4</sub>; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO<sub>3</sub>; 62.50 mg/L of NaH<sub>2</sub>PO<sub>4</sub>·H<sub>2</sub>O; 71.02 mg/L of Na<sub>2</sub>HPO<sub>4</sub>; 4320 mg/L of ZnSO<sub>4</sub>·7H<sub>2</sub>O; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H<sub>2</sub>O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H<sub>2</sub>O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H<sub>2</sub>O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H<sub>2</sub>O; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319

mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and 0.680 mg/L of Vitamin B<sub>12</sub>; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of

5 Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for

10 endotoxin assay in 15ml polystyrene conical.

The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degrees C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

15 On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the

20 polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

### 25 Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The

30 binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATS</u>	<u>GAS(elements) or ISRE</u>
	<u>IFN family</u>						
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
	<u>gp130 family</u>						
10	IL-6 (Pleiotrophic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrophic)	?	+	?	?	1,3	
	OnM(Pleiotrophic)	?	+	+	?	1,3	
	LIF(Pleiotrophic)	?	+	+	?	1,3	
	CNTF(Pleiotrophic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrophic)	?	+	?	?	1,3	
	IL-12(Pleiotrophic)	+	-	+	+	1,3	
	<u>g-C family</u>						
	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
20	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25	<u>gp140 family</u>						
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30	<u>Growth hormone family</u>						
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35	<u>Receptor Tyrosine Kinases</u>						
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., *Immunity* 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

10 5':GCGCCTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:4)

15 PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

20 5':CTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCT  
25 AGGCTTTTGGCAAAGCCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter  
30 molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not  
5 contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance  
10 gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS  
15 with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-  
20 2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

#### **Example 13: High-Throughput Screening Assay for T-cell Activity.**

25 The following protocol is used to assess T-cell activity by identifying factors, determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used  
30 in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately  
5 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells  
10 containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

15 During the incubation period, count cell concentration, spin down the required number of cells ( $10^7$  per transfection), and resuspend in OPTI-MEM to a final concentration of  $10^7$  cells/ml. Then add 1ml of  $1 \times 10^7$  cells in OPTI-MEM to T25 flask and incubate at 37 degrees C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

20 The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing polypeptides of the invention and/or induced polypeptides of the invention as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and  
25 resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells  
30 into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100,000 cells per well).



After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the  
5 assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophane covers) and  
10 stored at -20 degrees C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4 degrees C and serve as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the  
15 positive control wells.

The above protocol may be used in the generation of both transient, as well as, stable transfected cells, which would be apparent to those of skill in the art.

#### **Example 14: High-Throughput Screening Assay Identifying Myeloid Activity**

The following protocol is used to assess myeloid activity by determining whether polypeptides of the invention proliferates and/or differentiates myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is  
20 U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest  $2 \times 10^7$  U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing  
30 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM Na<sub>2</sub>HPO<sub>4</sub>·7H<sub>2</sub>O, 1 mM MgCl<sub>2</sub>, and 675 uM CaCl<sub>2</sub>. Incubate at 37 degrees C for 45 min.

- 5 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degrees C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 10 These cells are tested by harvesting  $1 \times 10^8$  cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of  $5 \times 10^5$  cells/ml. Plate 200 ul cells per well in the 96-well plate (or  $1 \times 10^5$  cells/well).

- Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37 degrees C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

20 **Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.**

- When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

- Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably

transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 5 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified 10 product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 15 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat- 20 inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine 25 protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% 30 confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as  $5 \times 10^5$  cells/ml.

- 5           Add 200  $\mu$ l of the cell suspension to each well of 96-well plate (equivalent to  $1 \times 10^5$  cells/well). Add 50  $\mu$ l supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ $\mu$ l of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the  
10           supernatant according to Example 17.

#### Example 16: High-Throughput Screening Assay for T-cell Activity

- NF- $\kappa$ B (Nuclear Factor  $\kappa$ B) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and  
15           CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF- $\kappa$ B regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- $\kappa$ B appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.
- 20           In non-stimulated conditions, NF- $\kappa$ B is retained in the cytoplasm with I- $\kappa$ B (Inhibitor  $\kappa$ B). However, upon stimulation, I- $\kappa$ B is phosphorylated and degraded, causing NF- $\kappa$ B to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- $\kappa$ B include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.
- 25           Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF- $\kappa$ B promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF- $\kappa$ B would be useful in treating diseases. For example, inhibitors of NF- $\kappa$ B could be used to treat those diseases related to the acute or chronic activation of NF- $\kappa$ B, such as rheumatoid  
30           arthritis.

To construct a vector containing the NF- $\kappa$ B promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF- $\kappa$ B binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC  
TTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

10 5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene)

Sequencing with the T7 and T3 primers confirms the insert contains the following  
15 sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGGACTTTCC  
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCC  
ATCCCGCCCCTAACTCCGCCCAGTTCGCCCCATTCTCCGCCCCATGGCTGA  
20 CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTA  
TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA  
GCTT:3' (SEQ ID NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-  
25 promoter plasmid (Clontech) with this NF- $\kappa$ B/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF- $\kappa$ B/SV40/SEAP cassette is removed from the above NF- $\kappa$ B/SEAP vector using restriction enzymes  
30 SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF- $\kappa$ B/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF- $\kappa$ B/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF  $\alpha$  (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

#### **Example 17: Assay for SEAP Activity**

As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15  $\mu$ l of 2.5x dilution buffer into Optiplates containing 35  $\mu$ l of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degrees C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50  $\mu$ l Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50  $\mu$ l Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

#### **Reaction Buffer Formulation:**

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5

19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

**Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability**

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used

instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO<sub>2</sub> incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO<sub>2</sub> incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10<sup>6</sup> cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10<sup>6</sup> cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca<sup>++</sup> concentration.

#### **Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase**

##### **Activity**

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine



Kinase RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also  
5 membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and  
10 non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of  
15 activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from  
20 Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or  
25 calf serum, rinsed with PBS and stored at 4°C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen  
30 Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na<sub>3</sub>VO<sub>4</sub>, 2 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub> and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4 degrees C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degrees C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg<sub>2</sub><sup>+</sup> (5mM ATP/50mM MgCl<sub>2</sub>), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl<sub>2</sub>, 5 mM MnCl<sub>2</sub>, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degrees C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mm EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degrees C for 20  
5 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degrees C for one hour. Wash the well as above.

10 Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

15

**Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity**

As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation  
20 (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other  
25 phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G  
30 plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any

of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degrees C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

**Example 21: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide**

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring

suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenin deoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

25

**Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample**

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

30

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in

- 5 Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled  
10 water to remove unbounded polypeptide:

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

- 15 Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale).  
20 Interpolate the concentration of the polypeptide in the sample using the standard curve.

### **Example 23: Formulating a Polypeptide**

- The secreted polypeptide composition will be formulated and dosed in a  
25 fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the secreted polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

- 30 As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will

be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the secreted polypeptide is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Compositions of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., Biopolymers 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release compositions also include liposomally entrapped compositions of the invention (see generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-

Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and 353-365 (1989))..

Liposomes containing XXX polypeptide may be prepared by methods known per se:

DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985);

Hwang et al., Proc. Natl. Acad. Sci. (USA) 77:4030-4034 (1980); EP 52,322; EP

5 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S.

Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal XXX polypeptide therapy.

10 In yet an additional embodiment, the compositions of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

Other controlled release systems are discussed in the review by Langer  
15 (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the secreted polypeptide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and  
20 concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both.  
25 Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

30 The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as



phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

10       The secreted polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any polypeptide to be used for therapeutic administration can be sterile.  
15       Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for  
20       example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

25       The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the  
30       agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

The compositions of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the compositions of the invention, include but not limited to, other members of the TNF family, chemotherapeutic agents, antibiotics, 5 steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the 10 combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

In one embodiment, the compositions of the invention are administered in 15 combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the compositions of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma 20 (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International 25 Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 30 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

Conventional nonspecific immunosuppressive agents, that may be administered in combination with the compositions of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

In a further embodiment, the compositions of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the compositions of the invention include, but are not limited to, tetracycline, metronidazole, amoxicillin, beta-lactamases, aminoglycosides, macrolides, quinolones, fluoroquinolones, cephalosporins, erythromycin, ciprofloxacin, and streptomycin.

In an additional embodiment, the compositions of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the compositions of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

In another embodiment, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the compositions of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl

estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., mephalen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g.,  
5 dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

In an additional embodiment, the compositions of the invention are administered in combination with cytokines. Cytokines that may be administered with the compositions of the invention include, but are not limited to, IL2, IL3, IL4,  
10 IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha.

In an additional embodiment, the compositions of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the compositions of the invention include, but are not limited to,  
15 Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PIGF), as disclosed in International Publication Number WO 92/06194; Placental Growth  
20 Factor-2 (PIGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515;  
25 Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E  
30 (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the compositions of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the compositions of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, 5 FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

In additional embodiments, the compositions of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

10 Example 24: Method of Treating Decreased Levels of the Polypeptide

The present invention relates to a method for treating an individual in need of an increased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an agonist of the invention (including polypeptides of the 15 invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering 20 to such an individual a pharmaceutical composition comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on 25 administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising 30 administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer. For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

10 **Example 26: Method of Treatment Using Gene Therapy-Ex Vivo**

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine

sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

#### **Example 27: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention**

Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411,

published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in  
5 the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be  
10 operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second  
15 targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added  
20 together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked  
25 polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place  
30 which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the



polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary  
5 phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl,  
10 5 mM KCl, 0.7 mM  $\text{Na}_2\text{HPO}_4$ , 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately  $3 \times 10^6$  cells/ml. Electroporation should be performed immediately following resuspension.

Plasmid DNA is prepared according to standard techniques. For example, to  
15 construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one  
20 non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The  
25 resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120  $\mu\text{g/ml}$ . 0.5 ml of the cell suspension (containing approximately  $1.5 \times 10^6$  cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is  
30 performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960  $\mu\text{F}$  and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their

genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer  
5 pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after  
10 having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a patient as described above.

#### **Example 28: Method of Treatment Using Gene Therapy - In Vivo**

15 Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter  
20 or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318  
25 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 94(12):3281-3290 (1996) (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The  
30 polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are

free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in  
5 Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain  
10 sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to  
15 provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and  
20 connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of  
25 the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely  
30 differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after

preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

5

**Example 29: Transgenic Animals.**

The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, *e.g.*,  
10 baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (*i.e.*,  
15 polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus  
20 mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, *e.g.*, Ulmer et al., Science  
25 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

30 Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to

quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

#### **Example 30: Knock-Out Animals.**

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (E.g., see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in

research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly  
5 administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (*e.g.*, knockouts) are  
10 administered to a patient *in vivo*. Such cells may be obtained from the patient (*i.e.*, animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (*e.g.*, lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the  
15 invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, *e.g.*, by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc.  
20 The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, *e.g.*, in the circulation, or  
25 intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, *e.g.*, genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and  
30 Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).



When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing  
5 for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders  
10 associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

**Example 31. Isolation of antibody fragments directed against polypeptides of the invention from a library of scFvs.**

15 Naturally occurring V-genes isolated from human PBLs are constructed into a large library of antibody fragments which contain reactivities against a polypeptide having the amino acid sequence of SEQ ID NO:Y to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein in its entirety by reference).

**Rescue of the library.**

A library of scFvs is constructed from the RNA of human PBLs as described in WO92/01047. To rescue phage displaying antibody fragments, approximately  $10^9$  *E. coli* harboring the phagemid are used to inoculate 50 ml of 2x TY containing 1% glucose and 100 micrograms/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU,  $2 \times 10^8$  TU of delta gene 3 helper (M13 delta gene III, see WO92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2x TY containing 100 micrograms/ml ampicillin and 50 micrograms/ml kanamycin and grown overnight. Phage are prepared as described in WO92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37°C without shaking and then for a further hour at 37°C with shaking. Cells were spun down (IEC-Centra 8, 4000 revs/min for 10 min), resuspended in 300 ml 2x TY broth containing 100 micrograms ampicillin/ml and 25 micrograms kanamycin/ml (2x TY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 micrometer filter (Minisart NML; Sartorius) to give a final concentration of approximately  $10^{13}$  transducing units/ml (ampicillin-resistant clones).

**Panning the Library.**

Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 micrograms/ml or 10 micrograms/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately  $10^{13}$  TU of phage is applied to the tube and

incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over  
5 turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log *E. coli* TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The *E. coli* are then plated on TYE plates containing 1% glucose and 100 micrograms/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as  
10 described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

#### Characterization of Binders.

15 Eluted phage from the third and fourth rounds of selection are used to infect *E. coli* HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtiter plates coated with either 10 picograms/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting  
20 (see e.g., WO92/01047) and then by sequencing.


It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above  
25 teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence  
30 listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>129</u> , line <u>N/A</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit  203105	Accession Number  August 13, 1996
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
EUROPE In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")     	

For receiving Office use only	
<input checked="" type="checkbox"/>	This sheet was received with the international application
Authorized officer  J. L. Smith Patent Division	

For International Bureau use only	
<input type="checkbox"/>	This sheet was received by the International Bureau on:
Authorized officer	

**ATCC Deposit No. 203105****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No. 203105****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>129</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 203081	Accession Number July 30, 1998
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
EUROPE In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer: Yvette A. Simms PCT International Division	Authorized officer

**ATCC Deposit No. 203081****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.



**ATCC Deposit No. 203081****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>131</u> , line <u>N/A</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>209090</u>	Accession Number <u>June 5, 1997</u>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>EUROPE</b> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")     	
<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application  Authorized officer <u>Yvette E. Simms</u> PCT Administrative Division	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on:  Authorized officer

**ATCC Deposit No. 209090**

### **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

### **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

### **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

### **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

### **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No. 209090****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.


**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>131</u> , line <u>N/A</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit  <u>209215</u>	Accession Number  <u>August 21, 1997</u>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>EUROPE</b> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer  Yvette E. Simms PCT International Division

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**ATCC Deposit No. 209215****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No. 209215****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>133</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 203027	Accession Number June 26, 1998
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) EUROPE In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application Authorized officer Yvette E. Simms PCT International Division	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on: Authorized officer



**ATCC Deposit No. 203027****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No. 203027

#### DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

#### SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

#### NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

*What Is Claimed Is:*

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group ;  
5 consisting of:
  - (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a  
10 polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - 15 (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X,  
20 having biological activity;
  - (f) a polynucleotide which is a variant of SEQ ID NO:X;
  - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
  - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
  - 25 (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.

5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide  
20 sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.

25

8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.

30

9. A recombinant host cell produced by the method of claim 8.

10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

5 (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;

(c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

10 (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

(e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

(f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

15 (g) a variant of SEQ ID NO:Y;

(h) an allelic variant of SEQ ID NO:Y; or

(i) a species homologue of the SEQ ID NO:Y.

12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.

25 14. A recombinant host cell that expresses the isolated polypeptide of claim 11.

15. A method of making an isolated polypeptide comprising:

30 (a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount  
5 of the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of  
10 claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to  
15 a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.  
20

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

(a) contacting the polypeptide of claim 11 with a binding partner; and

(b) determining whether the binding partner effects an activity of the  
25 polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

22. A method of identifying an activity in a biological assay, wherein the  
30 method comprises:

(a) expressing SEQ ID NO:X in a cell;

(b) isolating the supernatant;

305

- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

<110> Human Genome Sciences, Inc. et al.<sup>1</sup>

<120> 52 Human Secreted Proteins

<130> PZ032PCT

<140> Unassigned

<141> 1999-08-24

<150> 60/097,917

<151> 1998-08-25

<150> 60/098,634

<151> 1998-08-31

<160> 170

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

gggatccgga	gccccaaatct	tctgacaaaa	ctcacacatg	cccaccgtgc	ccagcacctg	60
aattcgaggg	tgacaccgtca	gtcttcctct	tccccccaaa	acccaaggac	accctcatga	120
tctcccgga	tcctgaggtc	acatgcgtgg	tggtggacgt	aagccacgaa	gacctgagg	180
tcaagttcaa	ctggtacgtg	gacggcgtgg	aggtgcataa	tgccaagaca	aagccgcggg	240
aggagcagta	caacagcacg	taccgtgtgg	tcagcgtcct	caccgtcctg	caccaggact	300
ggctgaatgg	caaggagtac	aagtgcaagg	tctccaacaa	agccctccca	acccccatcg	360
agaaaacat	ctccaaagcc	aaagggcagc	cccggagaacc	acaggtgtac	accctgcccc	420
catcccgga	tgagctgacc	aagaaccagg	tcagcctgac	ctgcttggtc	aaaggcttct	480
atccaagcga	catcgccgtg	gagtgggaga	gcaatgggca	gccggagaac	aactacaaga	540
ccacgcctcc	cgtgctggac	tccgacggct	ccttcttctc	ctacagcaag	ctcaccgtgg	600
acaagagcag	gtggcagcag	gggaacgtct	tctcatgctc	cgtgatgcat	gaggctctgc	660
acaaccacta	cacgcagaag	agcctctccc	tgtctccggg	taaatgagtg	cgacggccgc	720
gactctagag	gat					733

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

Trp Ser Xaa Trp Ser

1

5

<210> 3

<211> 86

<212> DNA

<213> Homo sapiens

<400> 3



2

gcgcctcgag atttcccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60  
 cccgaaatat ctgccatctc aattag 86

<210> 4  
 <211> 27  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 gcggcaagct ttttgcaaag cctaggc 27

<210> 5  
 <211> 271  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 ctcgagattt, ccccgaaatc tagatttccc cgaaatgatt tccccgaaat, gatttccccg 60  
 aaatatctgc catctcaatt agtcagcaac catagtcccc cccctaactc cgcccatccc 120  
 gcccctaact ccgcccagtt ccgcccattc tccgccccat ggctgactaa ttttttttat 180  
 ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240  
 ttttgaggc ctaggctttt gcaaaaagct t 271

<210> 6  
 <211> 32  
 <212> DNA  
 <213> Homo sapiens

<400> 6  
 gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7  
 <211> 31  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8  
 <211> 12  
 <212> DNA  
 <213> Homo sapiens

<400> 8  
 ggggactttc cc 12

<210> 9  
 <211> 73  
 <212> DNA  
 <213> Homo sapiens

<400> 9  
 gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg 60  
 ccattctaat tag 73

<210> 10  
 <211> 256  
 <212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

ctcgagggga	ctttcccg	gactttccg	ggactttccg	ggactttcca	tctgcatct	60
caattagtca	gcaaccatag	tcccgcct	aactccgcc	atcccgcct	taactccgcc	120
cagttccgcc	cattctccgc	cccatggctg	actaattttt	tttatttatg	cagaggccga	180
ggccgcctcg	gcctctgagc	tattccagaa	gtagttagga	ggcttttttg	gaggcctagg	240
cttttgcaaa	aagctt					256

&lt;210&gt; 11

&lt;211&gt; 3289

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

cdcggtcga	cccacgcgtc	cggcgaggac	cgcgccggc	gcagtcttca	atgagcagcg	60
cggaaactgc	acccagacc	cgagcctgct	gcgcgcccc	tcccagagct	cacctggtgc	120
caggtaacag	gcctggcctc	gccctgtgga	tgatgatggc	cttgcccccg	tgagctacaa	180
cctggccttc	agcaccgcc	cacctccaac	cagcaggatg	cggtgtgga	aggcggtggt	240
ggtgactttg	gccttcatga	gtgtggacat	ctgcgtgacc	acggccatct	atgtcttcag	300
ccacctggac	cgcagcctcc	tggaggacat	ccgccacttc	aacatctttg	actcgggtgct	360
ggatctctgg	gcagcctgcc	tgtaccgcag	ctgcctgctg	ctggggagcc	accattggtg	420
tggccaagaa	cagtgcgctg	gggccccggc	ggctgcgggc	ctcgtggctg	gtcatcacc	480
tcgtgtgcct	cttcgtgggc	atctatgcc	tggtagagct	gctgctcttc	tcagaggtgc	540
gcaggcccat	ccgggacccc	tggttttggg	ccctgttcgt	gtggacgtac	atttcaactc	600
gcgcatecct	cctgctctgg	tggctgctgt	ccaccgtgcg	gccaggcacc	caggccctgg	660
agccaggggc	ggccaccgag	gctgaaggct	tccctgggag	cggccggcca	ccgcccgaac	720
aagcgtctgg	ggccacgctg	cagaagctgc	tctcctacac	caagcccgac	gtggccttcc	780
tcgtggccgc	ctccttcttc	ctcatcgtgg	cagctctggg	agagaccttc	ctgcctact	840
acacgggccg	cgccattgat	ggcatcgtca	tccagaaaag	catggatcag	ttcagcacgg	900
ctgtcgtcat	cgtgtgcctg	ctggccattg	gcagctcatt	tgccgcaggt	attcggggcg	960
gcatttttac	cctcatattt	gccagactga	acattcgctt	tcgaaactgt	ctcttccgct	1020
cactgggtgc	ccaggagaca	agcttctttg	atgagaaccg	cacaggggac	ctcatctccc	1080
gcctggcctc	ggacaccacc	atggtcagcg	acctggcttc	cagaacatca	atgtcttctt	1140
gcggaacaca	gtcaagggtca	cgggcgtggg	ggtcttcatg	ttcagcctct	catggcagct	1200
ctecttggtc	accttcatgg	gcttccccat	catcatgatg	gtgtccaaca	tctacggcaa	1260
gtactacaag	aggctctcca	aagagggtcca	gaatgccttg	gccagagcga	gcaaacaggc	1320
ggaggagacc	atcagtgcc	tgaagactgt	ccggagcttc	gccaatgagg	aggaggaggc	1380
agagggtgtac	ctgcggaagc	tgcagcaggt	gtacaagctg	aacaggaagg	aggcagctgc	1440
ctacatgtac	tacgtctggg	gcagcgggct	cacactgctg	gtggtccagg	tcagcatcct	1500
ctactacggg	ggccaccttg	tcatctcagg	ccagatgacc	agcggcaacc	tcatcgctt	1560
catcatctac	gagtttgtcc	tgggagattg	tatggagaat	gtctccttca	gcctgtcccc	1620
cggcaagggtg	acggccctgg	tggggccctc	gggcagtggtg	aagagctcct	gtgtcaacat	1680
cctggagaac	ttctaccccc	tggagggggg	ccgggtgctg	ctggacggca	agcccatcag	1740
cgctacgac	cacaagtact	tgcaccgtgt	gatctccctg	gtgagccagg	agcccggtgct	1800
gttcgcccgc	tccatcacgg	ataacatctc	ctacggcctg	cccactgtgc	ctttcgagat	1860
ggtggtggag	gccgcacaga	aggccaatgc	ccacggcttc	atcatggaac	tccaggacgg	1920
ctacagcaca	gagacagggg	agaagggcgc	ccagctgtca	ggtggccaga	agcagcgggt	1980
ggcatggccc	gggtcttggt	gcggaacccc	ccagtcctca	tcctggatga	agccaccagc	2040
gctttggatg	ccgagagcga	gtatctgatc	cagcaggcca	tccatggcaa	cctgcagaag	2100
cacacggtac	tcatcatcgc	gcaccggctg	agcaccgtgg	agcacgcgca	cctcattgtg	2160
gtgctggaca	agggccgcgt	agtgcagcag	ggcaccacc	agcagctgct	ggcccagggc	2220
ggcctctacg	ccaagctggt	gcagcggcag	atgctggggc	ttcagcccg	cgcagacttc	2280
acagctggcc	acaacgagcc	tgtagccaac	ggcagtcaca	aggcctgatg	gggggcccct	2340
gcttctcccg	gtggggcaga	ggaccgggtg	cctgctgggc	agatgtgccc	acggaggccc	2400
ccagctgccc	tccgagccca	ggcctgcagc	gacctgaaag	gacctgccat	gtcccatgga	2460
tcaccgcttc	ctgcatcttg	cccctgggtc	ctgccccatt	cccagggcac	tccttacc	2520
tgctgcctcg	agccaacgcc	ttcacggacc	tccctagcct	cctaagcaaa	ggtagagctg	2580
cctttttaaa	cctaggtctt	accagggttt	ttactgtttg	gtttgaggca	ccccagtcaa	2640

4

ctcctagatt	tcaaaaacct	ttttctaatt	gggagtaatg	gcggggcactt	tcaccaagat	2700
gttctagaaa	cttctgagcc	aggagtgaat	ggcccttcct	tagtagcctg	ggggatgtcc	2760
agagactagg	cctctcccct	ttaccctccc	agagaagggg	cttccctgtc	ccggaggagg	2820
acacggggaa	cggtattttc	cgtctctccc	tcttgccagc	tgtgtgagtc	tggccagggc	2880
gggtagggag	cgtggagggc	atctgtctgc	catcgcccgc	tgccaatcta	agccagtctc	2940
actgtgaacc	acacgaaacc	tcaactgggg	gagtggggg	ctggccaggt	ctggaggggc	3000
ctcaggggtg	ccccagccc	ggcaccagc	gctttcgccc	ctcgccacc	caccctggc	3060
tggcagcctc	cctccccaca	cccggccctg	tgctctgctg	tctggaggcc	acgtggatgt	3120
tcatgagatg	cattctcttc	tgtctttggt	ggatgggatg	gtggcaaagc	ccaggatctg	3180
gctttgccag	aggttgcaac	atgttgagag	aaccgggtca	ataaagtgtg	ctacctctta	3240
cccctaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaagggcggc	cgctctaga		3289

&lt;210&gt; 12

&lt;211&gt; 2342

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

ggcacgagct	cagatctctc	ctggtacccc	ttccccacgc	ccttagataa	tccatctcaa	60
tctctcagtc	taattgagga	gctatggctg	caaggcacct	tccaggattt	cacacctaca	120
caaatctcct	ttttctcctt	ttgcctcttc	tgcttatggg	atattctgag	tccccacccc	180
caatcactga	cagctggggc	cccttcacac	gcctcacaca	ccacgtatta	agtcagtcac	240
aatckcccct	ctcmtctaac	tgctggattt	gtctttctac	acacacccaa	tgattcacgg	300
ctcttcgggc	tgamctactt	acatggacac	agtccaacgt	gagcttacac	atttcttact	360
tggctatacc	attcttggtc	gattcattcc	tgaamcgggt	tcataacctg	gaaactcagc	420
taaacacctt	tccttcaaac	tcagctctct	cagcatgggt	tcaggtaggg	ctgttgccct	480
ccttcaccta	atagcttctg	gactaacatc	catacaaac	aacacagcat	catctaaacc	540
accaatatgg	gggtaccatt	tctactcaaa	cttccttcat	atccccaccc	cccttatgtc	600
tcagccgaac	ctaccctaac	ccagcccacg	ccacaatggt	gggacagggt	ccccagtcct	660
tatgtggtct	tatttttacc	cttgcactcc	ctgtagacca	tcaattctac	accctaatta	720
caaaatcata	tccacctctg	cctggcgaaa	ggtgttatgc	ttttctggct	cgccctaccat	780
ccacacatcc	ctacacctca	ccaccggatc	ctcttttctt	tccttccatc	caattcctgg	840
cttccccgct	gccaactctg	ctctctatgt	ctccagttta	aagggtgcccc	ctggaaaaaa	900
tgtaacaatt	ccctcacctg	tgactgggtac	ctgacagcca	ccacaccggg	gcagcaatgg	960
ctaacggttg	acaaagacaa	tttctttctc	tctccaaaac	caaacagcct	tcatcaactc	1020
cctagccaag	actccctatc	aggcccttac	agggtgcgct	ctggctggga	gttacccaat	1080
tggggaaaac	gaaaataccc	tatcatggta	cctaccttca	cctacaactt	tgctgtcacc	1140
cccagttctc	ttttgtgtga	tacaactgat	atttkgccta	ccagccaact	ggtcagggaac	1200
ttgcaccctg	gtctttcagg	ctccaaccat	caacatccta	ccccctaacc	aaactattct	1260
aattttctgta	gaagcctcta	tctctctctc	accataaga	aataaatggg	ctctacatct	1320
catcacctcg	ctaacaggat	taggcatcac	tgctgcactt	ggcactggaa	tagcaggcat	1380
aaccacctca	atcacctcat	accaaact	attcacacac	ctttctaaca	ccgtagaaga	1440
tatgcacact	tccattacca	gtctccaacg	acaattagac	ttcctcgtgg	gagtcacctc	1500
tcaaaactgg	agagtcctgg	acctcctaac	cactgagaaa	gggggtacct	gcatatacct	1560
ccaggaagaa	tgctgtttct	gtgttaatga	atctggcatt	gttcatatcg	cagttcgtag	1620
gcttcatgac	agggctgcag	agctttgaca	tcaagtcgct	gactcctggt	ggcaaggatc	1680
atcccttcta	agatggatac	cctgggttgc	ccccttcta	ggaccctga	tcttctctt	1740
cctgttacta	atgattgggc	catgcatatt	taaccttgta	tcccgttca	tttcccaaag	1800
gctgaattgt	tttatccagg	caagcatgca	aaaacacatt	gataatata	ttcacctttg	1860
ccacgtctaa	taccagagcc	tacaggagaa	ccattcggaa	gctccagAAC	ccaggcccta	1920
atcacaacgc	ccctatccag	caggaagcag	ccagatgac	aacgacgccc	tttttctttt	1980
ttatactaaa	gtaagaaata	agaatgttag	cccaaactgc	actattttgc	agacccttac	2040
cattttacaa	actggtcaga	gtggaaaatt	ccaccagggc	ctgagctgtg	agaaacatcc	2100
tgtcaggcag	gtcccaggcc	taaccctgg	ctgcaactaa	ttccttcatt	atcagcagcc	2160
aaacacaccg	ccccacccc	attttcacaa	caatcccaga	cctctcctgc	ccgggactgt	2220
aactggtcca	gcctgtaagc	gggaaggggg	ctctggcact	agctggtacc	ccctctccgc	2280
aggtctttct	cccaataaat	ctgtgttgcc	attgaaaaaa	aaaaaaaaaa	aaaaaaaaactc	2340
ga						2342

<210> 13  
 <211> 1666  
 <212> DNA  
 <213> Homo, sapiens

<400> 13  
 ggtggagttc gcacctccag ctcgggccga tgtggaagct ttggagagct gaagagggcg 60  
 cggcgccgct cggcgccgct ctcttcctgc tgctcttcgc gctaggggtc cgccagctgc 120  
 tgaagcagag cggcgccgatg ggcttcccc cggggccgcc, ggggctgccca tttatcggca 180  
 acatctattc cctggcagcc tcatccgagc ttccccatgt ctacatgaga aagcagagcc 240  
 aggtgtacgg agagatcttc agtttagatc ttggaggcat atcaactgtg gttctaaatg 300  
 gctatgatgt agtaaaggaa tgccttggtc atcaaagcga aatttttgca gacagaccat 360  
 gccttccttt, attcatgaag atgacaaaaa tgggaggctt actcaattcc agatatggcc 420  
 gaggatgggt tgatcacaga cgattagctg taaacagttt tcgatatttt ggatatggcc 480  
 aaaagtcttt tgaatctaaa atcttggaag aaaccaaatt tttcaatgat gctattgaaa 540  
 catacaaagg tagacctttt gactttaaac agttaataac gaatgctgtt tcaaacataa 600  
 ccaatctgat ctttttgga gaacgattca cttatgaaga caccgatttt cagcacatga 660  
 ttgagttatt tagtgaaaat gtggaactag ctgccagtgc ctgagtcctc ttgtataatg 720  
 cctttccatg gattggcatc ctgccttttg gaaaaatca acagctgttt agaaatgcag 780  
 ctgtagtcta tgattttctc tccagactca ttgaaaaagc ttcagtcaac agaaagcctc 840  
 agctacctca gcattttgtt gatgcttatt tagatgagat ggatcaagggt aaaaatgacc 900  
 catcatctac tttctccaaa gaaaacctaa tttctcagt gggggaactc atcattgctg 960  
 gaactgaaac taacaaccaat gtgctacggt gggcgattct tttcatggcc ctttatccta 1020  
 atattcaagg acaagttcag aaagagattg atttaattat gggccctaatt gggaagcctt 1080  
 cttgggacga caaatgcaaa atgccttata ctgaggcagt tttgcatgaa gttttaagat 1140  
 tctgtaatat agttccatta gggattttcc atgcaacctc tgaagatgca gttgtacgtg 1200  
 gttattccat tcctaaaggc acaacagtaa ttacaaatct ttattctgta cactttgatg 1260  
 aaaagtactg gagagacca gaagtgttcc atcctgagcg atttctggac agcagtggat 1320  
 attttgccaa gaaggaagct ttggttcctt tttccctagg aagaagacat tgtcttgag 1380  
 aacacttggc tcggatggaa atgttcttgt tttttacagc attgcttcag aggtttcatt 1440  
 tgcattttcc acatgaacta gttccagatc tgaagcccag gttaggcatg acattgcagc 1500  
 cccaacccta cctcatctgt gctgaaagac gctgaaactg cctgggatgt tttcgggaac 1560  
 aagaatgtat atttgcctta tccctgaact tggtttaatc aaatcaatgt gtgtattaga 1620  
 ataaaagtca cagcatcaaa aagmcaaaaa aaaaaaaaaa aaaaaa 1666

<210> 14  
 <211> 2027  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (294)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1976)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1981)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1985)  
 <223> n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2021)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 14

ggcagcagtt	gggagcagct	ctgcgtgcgg	ggcctcagag	aatgaggccg	gcgttcgccc	60
tgtgcctcct	ctggcaggcg	ctctggcccg	ggccggggcg	cggcgaacac	cccactgcgg	120
accgtgctgg	ctgctcggcc	tcggggggcct	gctacagcct	gcaccatgct	accatgaagc	180
ggcaggcggc	cgaggaggcc	tgcctcctgc	gaggtggggc	gctcagcacc	gtgcgtgcgg	240
gcgcgagct	gcgcgctgtg	ctcgcgctcc	tgccggcgag	cccaggggcc	gganggggct	300
ccaaagacct	gctgttctgg	gtcgcaactg	agcgcgaggc	ttccactgc	amcctggaga	360
acgagccttt	gcgggggttt	tcctggctgt	cctccgaccc	cgccggtctc	gaaagcgaca	420
cgctgcagtg	ggtggaggag	ccccaacgct	cctgcaccgc	gcggagatgg	gtacttccag	480
gccaccgggt	gggtcgagcc	cgcagctgga	aggagatgcg	atgccacctg	ygcgccaacg	540
ctacctgtgc	aagtaccagt	ttgaggtctt	gtgtcctgcg	ccgcgccccg	gggcccgcctc	600
taacttgagc	tatcgcgcg	ccttcacagt	gcacagcgcc	gctctggact	tcagtccacc	660
tgggaccgag	gtgagtgcgc	tctgccgggg	acagctcccg	atctcagtta	cttgcatcgc	720
ggacgaaatc	ggcgctcgyt	gggacaaaact	ytggggcgat	gtgttgtgtc	cctgccccgg	780
gaggtacctc	cgtgctggca	aatgcgcgag	gtccctaac	tgccatagacg	acttggggagg	840
ctttgcctgc	gaatgtgcta	cgggcttcga	gctgggggag	gacggccgct	cttgtgtgac	900
cagtggggaa	ggacagccga	cccttggggg	gaccggggtg	cccaccaggc	gcccgcgggc	960
cactgcaacc	agccccgtgc	cgcagagaac	atggccaatc	agggtcgacg	agaagctggg	1020
agagacacca	cttgtccctg	aacaagacaa	ttcagtaaca	tctattcctg	agattcctcg	1080
atggggatca	cagagcacga	tgtctaccct	tcaaatgtcc	cttcaagccg	agtcaaaggc	1140
cactatcacc	ccatcaggga	gcgtgatttc	caagtttaat	tctacgactt	cctctgccac	1200
tcctcaggct	ttcgactcct	cctctgccgt	ggtcttcata	tttgtgagca	cagcagtagt	1260
agtgttggtg	atcttgacca	tgacagtact	ggggcttgtc	aagctctgct	ttcacgaaag	1320
cccctcttcc	cagccaagga	aggagtctat	gggcccgcgg	ggctggagag	tgatcctgaa	1380
gcccgtgctg	ttgggctcca	gttctgcaca	ttgcacaaac	aatgggggtga	aagtcgggga	1440
ctgtgatctg	cgggacagag	cagagggtgc	cctgctggcg	gagtcacctc	ttggctctag	1500
tgatgcatag	ggaaacaggg	gacatgggca	ctcctgtgaa	cagtttttca	cttttgatga	1560
aacggggaac	caagagggaac	ttacttgtgt	aactgacaat	ttctgcagaa	atcccccttc	1620
ctctaaattc	cctttactcc	actgaggagc	taaatcagaa	ctgcacactc	cttccctgat	1680
gatagaggaa	gtggaagtgc	ctttaggatg	gtgatactgg	gggaccgggt	agtgtggggg	1740
agagatatatt	tcttatgttt	attcgagaaa	tttgaggaga	tgattgaact	tttcaagaca	1800
ttggaaacaa	atagaacaca	atataattta	cattaaaaaa	taatttctac	caaaatggaa	1860
aggaaatggt	ctatgttggt	caggctagga	gtatattggt	tcgaaatccc	agggaaaaaa	1920
ataaaaaata	aaaattaaag	gattgttgat	aaaaaaaaaa	aaaaagggcg	gccgcnctag	1980
ngggnccaag	ctttacgtac	gcgggcatgc	gacgtcaagc	ncttcca		2027

&lt;210&gt; 15

&lt;211&gt; 2334

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2278)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2290)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 15

gggagtgtgg	ctgcagaacc	cagggtggcag	ggctttcctc	aggcccccta	ctcctgacct	60
------------	------------	-------------	------------	------------	------------	----

7

```

ggacgaggcc ggggcttddt caaggaggct ctctgactgc cacccttgcc tgcctgcccg 120
gccctgcaca acatgcagcc ctccggcctc gagggtcccg gcacgtttgg tcggtggcct 180
ctgctgagtc tgctgtctct gctgctgctg ctccagcctg taacctgtgc ctacaccacg 240
ccaggccccc ccagagccct caccacgctg ggcccccca gagccacac catgccgggc 300
acctacgctc cctcgaccac actcagtagt cccagcacc agggcctgca agagcaggca 360
cgggccctga tgcgggactt cccgctcgtg gacggccaca acgacctgcc cctggctcta 420
aggcaggttt accagaaagg gctacaggat gtaacctgc gcaatttcag ctacggccag 480
accagcctgg acaggcttag agatggcctc gtgggcgccc agttctgggc agcctatgtg 540
ccatgccaga cccaggaccg ggatgccctg cgctcacc, tggagcagat tgacctcata 600
cgccgcatgt gtgcctccta ttctgagctg gacttgtgta cctcggctaa agctctgaac 660
gacactcaga aattggcctg cctcatcggt gttaggggtg gccactcgct ggacaatagc 720
ctctccatct tacgtacctt ctacatgctg ggagtgcgct acctgacgct caccacacac 780
tgcaacacac cctgggcaga gagctccgct aagggcgtcc actccttcta caacaacatc 840
agcgggctga ctgactttgg tgagaagggt gtggcagaaa tgaaccgcct gggcatgatg 900
gtagacttat cccatgtctc agatgctgtg gcacggcggg ccttgggaagt gtcacaggca 960
cctgtgatct tctccactc ggctgcccgg ggtgtgtgca acagtgtctg gaatgttctt 1020
gatgacatcc tgcagcttct gaagaagaac ggtggcgtcg tgatgggtgc tttgtccatg 1080
ggagtaatac agtgcaacct atcagccaat gtgtccactg tggcagatca cttcgaccac 1140
atcaaggctg tcattggatc caagtctatc gggattgggt gagattatga tggggccggc 1200
aaattccctc aggggctgga agacgtgtcc acatacccag tctgataga ggagttgctg 1260
agtctgtggt ggagttagga agagcttcag ggtgtccttc gtggaaacct gctgcggggtc 1320
ttcagacaag tggaaaagggt acaggaagaa aacaaatggc aaagccctt ggaggacag 1380
ttcccgatg agcagctgag cagttcctgc cactccgacc tctcagctc gcgtcagaga 1440
cagagtctga cttcaggcca ggaactcact gagattccca tacactggac agccaagtta 1500
ccagccaagt ggtcagctc agagtctcc cccacatgg cccagtcct tgcagttgtg 1560
gccaccttcc cagtccttat tctgtggctc tgatgacca gttagtcctg ccagatgtca 1620
ctgtagcaag ccacagacac cccacaaagt tcccctgttt gcaggcacia atatttctctg 1680
aaataaatgt tttggacata gaaaaaaaaa aaaaaaaag ggcggccgct cttagggatc 1740
cctcgagggg cccaagctta cgcgtgcatg cgacgtcata gctctctccc tatagttagt 1800
cgtattataa gctaggcact ggccgtcgtt ttacacgtc gtgactggga gatctgctag 1860
cttgggatct ttgtgaagga accttacttc tgtggtgtga cataattgga caaactacct 1920
acagagattt aaagctctaa ggtaaatata aaatttttaa gtgtataatg tggtaaacta 1980
gctgcatatg cttgctgctt gagagttttg cttactgagt atgatttatg aaaatattat 2040
acacaggagc tagtgattct aattgtttgt gtattttaga ttcacagtcc caaggctcat 2100
ttcaggcccc tcagtcctca cagtctgttc atgcacataa tcagccatac cacatttgtta 2160
gaggttttac ttgctttaaa aaacctycca caccctcccc tgaacctgaa acataaaatg 2220
aatgcaattg gtggtggtaa cttggttaat ggagcttata atggtaccaa taaagcantg 2280
catcacaan ttcccaaata aagcattttt tcttgggaatt taaatggggg ttgg 2334

```

&lt;210&gt; 16

&lt;211&gt; 2608

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

```

ccacgcgtcc gcacagggct caggctgggc tcagaggcca ctgagatgcc agctccttga 60
gagcagtggg ggtgtcccag gccagagcag cctcttctc tcttggctcc agaaaaaccc 120
ttgcagtaaa tgggtggcctc tgggtgggta cttctagcgc aggttctctt cttctcteta 180
gtcctctccc gtgcccggg tgctgggtgc tggatggatg ggcgtctctt agctcctccc 240
ggtgccctgg gtgctgggtg ctggatgggt gggcgtctc tagctctctc cggtgccctg 300
ggtgctgggt gctggatggg tgggcgccac ggtgcaccct tgttgggctg cctgtgcccg 360
agtggcctct gcagctctta tgtctgcctc taatgggatg tgcgccctga ctgcctcggt 420
cttaagggca tagtgggtcg gctaagatct gatgcgcagg actgcgttct gggcaggtgc 480
tgggaaggcg gaaccaagcc gtccgtgccg ctaggagagc gagactgccg gaaagaagag 540
ggcaggaggg gggcgtgggt gtgcagcccc accccgggga ggggctttag gcaactggaa 600
ggaaagtcct gttggaggaa ttcggtgggt gttcacacc gcctcgctgt cttggagctt 660
tgatctgtcg tcggcgggtc gctggcacag gactaaacat ggctgaggct gggctccagc 720
cagatctggc tggggacagc accgcgtggg cccaggatcc acccagagca ggcaggcctt 780
gctggggccc cagtcaagtc cacttccagt gaggagagcc agccgggagg tcagtgccag 840

```

8

```

agctctggtg gagccagac cctgccttcc ctgagggccg cccctgtcgc cgccctgggg 900
tccctgtcct cctatcctga ctcctgcccc agggccacca cccctgaact gtgccctggg 960
gccccaccc tccacctggc cgactccatc tctgggcctg tcagtccacc tgggtcctct 1020
ctggggcctg atgcctggac cctctgtgcc aagcaccacc aagcaaaggg gatgaccttg 1080
ggcaccacca aggtgctgag actacagcca gtgagccct gctggggggc aaagtcattg 1140
agggtgcctg ggcccttcca acctggaagg aggaggggag agagcaggca gcagggcagg 1200
gggaagagga ggagtgcctg atctgcacag agccctacgg gccagagag cgccgcctgg 1260
ccctgctgaa ctgtagccac ggctgtgtg tgggctgcct gcacaggctg ctgggctcgg 1320
cctccagtgc cgacctgggc cgggtgcgct gcccgctgtg ccgcagaaga cgcccgctgt 1380
ggagtgggag atctgccggc tacaggagga gctgtctacag gccgacgggc ctcacgcca 1440
gccccgccga gagggccctg cactctatca ccgcaaccct gggccctggg gctccctgga 1500
gcaccgctac cagctgcgct tccctggcagg gcccgctggg gccgggggct gcctgccctt 1560
cctgccctgt ccacctgcc tgggtgcccg gctctggacc ctgctgggag ggggacctg 1620
tgcccgccgc ctggcgctgc tgagcctgct ggcccttgag cttctggggc tgtgtggt 1680
cttcacgccg ctctgtctgc tgggactgct cttctgtctc ctggaccgct ctggccgctg 1740
agcagagccc aggacagccc cgccgcaaca ggccaggggg ccagactgg ccacgtccc 1800
catgcctggg tgctgtgagg cctgatgacc aggtcgaaa aacccaaggt tgggtccagg 1860
gcagtggcct tcaatcaaga cctccattg ctgaaccac aaccagggt acccagaggc 1920
ctgaccctgc agagtccatg gctgactgc tgccagaca ctactgaac ccaaggacac 1980
cagcgcccaa ggacagctcc tggaggaggc cagcccagca ggaaagtctg tgagcaggac 2040
cccattcacc ctgcggcaga cgggoaccgt actggccacg ggctgacgcc ggccacactt 2100
cccctccgag ggccagctga gcacagcagg catgaaagca aacagagata cagcagttag 2160
tcagttcctt ggagagggca gggactccgc ccacctgtg ttcagataag ggccagtgtg 2220
tgtccctgaa ggtcaggcca gccgggggag ggggtccatgc tgcgaaaatt cagcctgcaa 2280
aggctcctct cccacttga tcaggcccag accaggtggg ggttggcgct ggcctgtgtt 2340
gcaggggaca agggcccacc caggccttgg aacataagct ctgcccctgc acaccctcat 2400
gtcaccacac ctgggatgga gacatcaggt ggcccagcga gagatggagg actgatcctg 2460
gaacgtgaag cagctttcaa taaaccagct cctggggaaa aaaaaaaaaa aaaaaaaaaa 2520
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2580
aaaaaaaaaa aaaaaaaaaa aaaaaaaa 2608

```

&lt;210&gt; 17

&lt;211&gt; 1291

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1279)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1286)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1290)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 17

```

aaacctcttc tataggtaaa gctggwacgc ctgcaggtac cgggtccggaa ttcccgggtc 60
gccacgcgt ccggaaagag gaaacataga ggtgccaaag gaacaaagac ataatgatgt 120
catccaagcc aacaagccat gctgaagtaa atgaaacct acccaacct taccaccaa 180
gcagctttat ggctcctgga tttcaacagc ctctgggttc aatcaactta gaaaaccaag 240
ctcagggtgc tcagcgtgct cagccctacg gcatcacatc tccgggaatc tttgctagca 300
gtcaaccggg tcaaggaaat atacaaatga taaatccaag tgtgggaaca gcagtaatga 360
actttaaaga agaagcaaag gcactagggg tgatccagat catggttggg ttgatgcaca 420

```

```

... ttggttttgg aattgttttg tgtttaatat ctttctcttt tagagaagta ttaggttttg 480
ccttactctgc tgttattggt ggatacccat tctgggggtgg ctttctcttt attatctctg 540
gctctctctc tgtgtcagca tccaaggagc tttcccgttg tctgggtgaaa ggcagcctgg 600
gaatgaacat tgktagttct atcttggcct tcattggagt gattctgctg ctgggtggata 660
tgtgcatcaa tggggtagct ggccaagact actgggccgt gctttcttga aaaggcattt 720
cagccacgct gatgatcttc tccctcttgg agttctctgt agcttgtgcc acagcccatt 780
ttgccaacca agcaaacacc acaaccaata tgtctgtcct gggtattcca aatatgtatg 840
aaagcaaccc tgtgacacca gcgtctcttt cagctcctcc cagatgcaac aactactcag 900
ctaattgcccc taaaagaaaa aggggtatca gtctaattctc atggagaaaa actacttgca 960
aaaacttctt aagaagatgt cttttattgt ctacaatgat ttctagtctt taaaaactgt 1020
gtttgagatt tgtttttagg ttggtcgcta atgatggctg tatctccctt cactgtctct 1080
tctacatta ccactactac atgctggcaa aggtgaagga tcagaggact gaaaaatgat 1140
tctgcaactc tcttaaagtt agaaatgttt ctgttcatat tactttttcc ttaataaaat 1200
gtcattagaa acaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaggcg gccgctctag 1260
aggatccaag cttacgtang cgtgcntgcn a 1291

```

&lt;210&gt; 18

&lt;211&gt; 3129

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

```

cggcacgagg ggagaccgaa tggcggggtg aggggagcta ctgcgtccgg ctttgcaggc 60
cgttcctttc agggttgccg cagcaatggc acttcatcct ggcagcagcc atttgctggt 120
tgcagtgcc a gtttcttggt ttcttttttg cattcctgga atcagtttca tcaactctctc 180
ttggagttac caagagtcgc cagtcagttt cctctccgtg gaaggctgag tcccaattcc 240
atagggctct gtgccagact tttagtctct gataactccg aaatccttcc ttctgtttcc 300
taagctctac aggtattcta gtttttctcg caatttttca ttactaggct acatcagtgt 360
ccttttcttt tctttctttt tctcactttt tatttgtacc tttagtctc caacaattaa 420
caattcccca tgttaaattc tctctgttta aacaactaac gggctctctg tattccaact 480
agatctcggc tgatgcaaat gccgtagaca atcaatcgat caatcaatca gcaaatccaa 540
tttccctcta gccactaaag ggacttaata taattgaatg attataaaat tctgatccca 600
tttatctcaa gccattgaaa atatttttctg tgtgattata agatacgtaa ttttaaaaaa 660
tgattttcta aatcaggatt cttaccagat tatttcacag cttattcact attgattgc 720
tttccctcca ccttctgttag cttttgtatg cctgtcctgc attttaaacc aaggtagag 780
gaggataaag taaatagtaa atggtagat agtggccaaa actttgaatc tttttcccta 840
ttcatcatct ttgtgcaagt tggtagggag gtccactaat tcagcctccc cattccctgt 900
cccttcaaag taaatcagaa gcttcaggta aaaagcagag aagctaccca agttcttttc 960
atatttttta tcatcatact aataattttg ctttctgaaa tgattactct tacaagcagc 1020
aaatatattt tttaggaggt tagaattatt gatgtcacgt tttcatattc agtatcccaa 1080
gtaaatatca ggtaaattca ggaaaataaa tacaaccatt ttcttggtta attgttttta 1140
aatttaatat atgttcattt attcttttaa atatatattt cataaatggg gactatgtat 1200
tccatgacat aaaaactaag tatttttagt gggaaattct ttgtggaatc actgaaatga 1260
taactagtgt tagcaatact tctgtttcat gagacgagtc actttatgaa gattgcaaat 1320
tattttttgt cccctctaga gttcatttca tgggaaattt aaattttatt gtcttaactt 1380
taatgttaaa catttcttcc acatatggta tggccagaca tgttttcttc tttgctgcta 1440
ttttacattt tagatgtagt ccattcagga tctttttttt ttttttttag gagtcagagc 1500
ttgaagtgat gcaacacttt cactctttat ctctgattta catgctaata aagacaactg 1560
atagacaata tgatatactt tggccctaga agatccttaa tgaggaaatt ctgtggcctg 1620
ttttctcttc tgtgctattt tttgtcttct aaggcaggca tttttatcag ttcagtttgt 1680
tgctgtgact agttcaaact tgttcagggc tttatgcctg aaataagtaa tcaattagcc 1740
tatatatata aatatataac caaccattac aattatctaa gaatgtcagg gagttgtgaa 1800
gtggaaacaa attaaagagt tgcaaaatta aaagagctcg tcttccaaca gaagtgtgct 1860
gacatatatt aaagaataaa tgaattaaag tcaccaaaaga ggaacaggag ttaaaaagaa 1920
gaccattgtt tagatggagt caggctgggt cactgccaaag gaaagcaaac aggtattctg 1980
aaggcttggg atgaatttca gcttgcatca tccgctgga ggggggtgatt tcccaatat 2040
catgttagat ccatagcttg ttcttgacag agtggcagta ccttctctcc actgcactct 2100
caccactagc atagatgtaa aacacagtaa gtactcagaa actacttgaa gagtgcagtt 2160
atcagtagag atgatcgaat catttgtttt tctagggaaat atttttgcct ttcttcttcc 2220

```



10

agaatcctct	ggttataatg	tgctcactgc	taggtcacca	gtcataaaac	attatgtaga	2280
ggttactggt	cattatccta	atatatttat	caaaaaatct	ggagtatatg	aaactgcctt	2340
tcattgtaac	attagacaaa	aacattttatt	ctatcaaata	cagacttaaa	actgccacca	2400
aattggaaga	atatgatctt	aaatttaaaa	aaaccccata	tacttaaaaca	caataatcta	2460
tctttatcta	ccttcctaaa	cattaatgca	tgagacttct	cttaatatca	tgaatatgca	2520
ctatgatttt	tatgtttacat	ctttttcttg	tttccattta	tgctagttaa	atattattagt	2580
atccttcatt	gaaacactat	tattttccat	ggaggaaaat	attttatttc	ctttatgaaa	2640
agggtgcttt	acttctgaaa	tacagaattc	attttgtttc	ggattctggt	tgtttgtttt	2700
cacatcaact	tcattctaaa	tgttcattca	aaattatttt	tcagacttga	gttcagggtga	2760
aaatgttaat	ggaattaaaa	agtagttagt	caaattgaaat	ttcaatatat	aagtcaaatt	2820
tgaagaaaac	tgaatttaata	aaggattcta	agtttataaaa	gaaatcaaaa	ttatgctttc	2880
aaaaatatta	tgtaactgga	gaaagtaatt	ttatttttgag	atctctcatg	attctttaat	2940
atatattttt	tcttattagt	atgpaacctt	ttggtaacat	atataggaga	cagttaattt	3000
tggttaagact	acataattca	tacattcttc	atcttgacat	acatgatcaa	atactttata	3060
acgtccacta	gccttgctct	tgctccactgc	aatggaaaaa	aaaaaaaaaa	aactcgaggg	3120
ggggcccg						3129

&lt;210&gt; 19

&lt;211&gt; 3629

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

ccacgcgtcc	gccagaggct	gggaaggaaa	aattattccc	atcttaaaat	atcttaaaaa	60
tatgtacaga	tttattgtaa	atcaaaattt	tcctcataac	tacagcccct	tcccacacat	120
aaaattcccc	ctaaatcagg	aacataatgt	acagcctggt	tttaacttgt	atttttccat	180
tcacactatg	tcacaaaaaa	attctcatgg	tgatccatga	tttcacaggt	cctgtccatg	240
tgtttctctga	gaagacagtc	ctggaatgga	attactaatt	caaaaggcat	ttttaaggct	300
gttacacata	ctgccaaatc	ggccttcagg	aggatgacac	aaattccacc	tatgtcatct	360
gagcgtgcc	ctatctgctg	ctattgcccc	atggttgagg	ttttgtcctt	tctcctcatt	420
ctcatgggca	actgtcaaca	tcaccaccac	tgatgctctg	aaaagaaagg	ctcagaagac	480
atagacagtc	tgggattttt	ttttttccct	acctttcttt	gactgtagg	gaaggagtga	540
gatgtggctg	caggattggg	gaatagctaa	gtcagaaatg	agcagcagtg	caactgtggc	600
aatcttgagg	gggggtctca	gccatgaaag	aaaacagggg	gtgccttgcc	agctggaggc	660
tgacagaatct	tagcttgagg	gttttgccct	ctccgtggaa	ctgacagatg	acaggaatga	720
cgtggagtat	tctcatcaag	cgtctcagca	tagtcagctc	ctgagatggc	aaactttagg	780
caaagacacc	tgggctgaaa	ggteccacat	ttaaaacttg	tcagtcagat	aactgcccac	840
caccaccctc	tgtaagatac	ctagatatgg	gaaccccaca	ctcttcccga	ataagactct	900
tgtatgactg	aaagactttt	tggttaaaaa	catattctaa	tctaacctgt	attttccctg	960
ctgcaatttc	agcctttttc	tatctttttc	tggtgaatga	acaccacaca	ggctttcaga	1020
tcagacaact	tctggctccg	tcattttacta	gctgtatcag	ttaacaccag	ctgctgactt	1080
cacttctgca	tatgcattcc	aaaatttagtg	tcttaaaaata	acaactactt	attacttttc	1140
acaagcctaa	aagttggctg	aatgggtctg	ctgatctgac	ttgggttaac	ttggctggac	1200
actgtctgat	tcagcatagg	ttgactgtcc	tctgcctcat	gtctcttata	atccataaat	1260
tgcccaggct	tgcttgccctg	aaggcagaag	agttcccact	agcaggagag	ggcaagcact	1320
tttcaatgca	caagcacctt	tcaagcctct	atgtgcatca	agtttgctaa	tgctccgctg	1380
accaaagcaa	atcatgtggc	ttacttcgga	gtcagtggtg	gtaggcatta	cccaaagaaa	1440
gtggctatag	ggaagcatga	agatctagag	ccatttagta	aatcaatcta	ccacattaac	1500
tttgtagcct	tgggcaattc	ccttagcctc	tctgagcttt	aactctataa	tctataaaaag	1560
tgacagaattt	ttttgggatt	cttttgctata	atgtatgtaa	aatgtctgct	acacaataaa	1620
catgccatat	ttattaattt	tctttttctc	tctttttcat	catttgaatt	ttttcttaaa	1680
tgcatatata	tatgtgcata	tatatatgtc	catctcgata	tatatatatg	tatatatata	1740
tatatatatg	acttactctg	tcatccaggt	gagagtgcag	tggcatgatc	acagctcact	1800
gcggcctcga	cctcccaggc	tcgagtgatc	gtcccacctc	agcctcccag	gtagctggaa	1860
ctacagatgt	gtgccaccac	acccagctaa	atttttttgt	atattttata	gagaaggggt	1920
tcattatggt	gctgaggctg	gtctttaact	cctgggttca	agcaatccac	ccaccttgac	1980
ctcgcaaaagt	attgggatta	caggcatgag	ccaccatgcc	tggcccttaa	atgcatcatt	2040
cttatacatt	cctttataca	tacaaagctc	gaagataata	ataataaaca	ttattcccac	2100
ggcttctggt	gaatcccca	ctcctagcct	gtgcctaata	aagtgtgccc	tcaattcatt	2160

11

```

ccaatggtct ggaatttgaa gactaaagga atcaaagggt atctcttccc tttgttcctg 2220
ggaccatcaa gttgtgaaat gacgtcttgg ctttctgac tgggtccact accacaaatg 2280
aaatagaaag gagttaacac atttgcatta actgatgatt cttaatagag gcaataaaat 2340
gagaaacccc catttcctaa tttaaatgtg gctttgggga taggataagt atcctggccg 2400
ctccttagct aatcagggtc cagaaggggac attaatgca gccagcgcat cctgctgggg 2460
agacccaagg ctgccagctc cttgctgag aatggaatag aattttaatt ggctgtgaaa 2520
gcatcatggt aggaggtaca ggaagaggt ggtctgctcc cccaaagcag gttagggaag 2580
ctccaaggga aaggactcaa acagggtgat ggcagcaaga acctcccatg ctgaggggcc 2640
ttcccagctg caacaccaca gttccaattc cagggtgga ctccagccta ggtgaaggac 2700
cttctcacac ttgggcgatt agtactggag cacggagcct gtgaatcttc tcatctctgt 2760
gcttccccca acttccatga gagaagcacc acatagttac tgtcacttgt attgtcaatc 2820
ttcgacttca gagaggtctc caaatataag ctcaactccc cagtgtactg tgtgtgtgcc 2880
attagtccat gtgtatggtc tgaagcaatg gcacagcatt cctgtactgt tgtttcagag 2940
caatgtgaat ttatttcact cacaataaat taatctaact atgcttaggg caaagtgtgt 3000
ctcatggtaa tgagtatgtc ctttctgcca tggcaatttc tggacacac atatagacaa 3060
gaatgacatg tgaagggtcaa tagatgagac tatgaaaaac aagacaatat agcttttttag 3120
cataaatgta caatgatgca tgtggttttt ggagattgtt ggacaaatta cttaacctct 3180
tggtgcctta gttttctaatt tattaataatg agggcaacag tattgctgat ctaataaggc 3240
ttttggaaag aataatgaga tgtaaagagc ttagaatagt gcagagtctt aataaacatg 3300
tcttcgtgat tccttctagt gaagtgacat agagaagtgg gaccagtcac tgacccaccc 3360
aagctctgtc aaacctagtgg tatagtaaat ctggtggcaa aaagcaaagc agaacaacaa 3420
aaaaacgcaa gcaaaatcat tgctttcagg aaactatttt gtgagacaga agcttgataa 3480
aaattcaaaa ttgacagtga actctacttc attaataact taaaactcac agcaataaac 3540
cattacttag aacaagtaag cagttttgtt ttatttgtgc attttgatat ttattttcaa 3600
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3629

```

&lt;210&gt; 20

&lt;211&gt; 1144

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

```

tcgacccacg cgtccgcgct tgacttgctt ccagacaaag gttgtctcaa gtttgttgc 60
caaaccgagt tctggagAAC gccatcagct cgctgcttaa aattaaacca cagggtccat 120
tatgggtcga cttgatggga aagtcacatc ctgacggcc gctgctcagg ggattggcca 180
agcagctgcc ttagcttttg caagagaagg tgccaaagtc atagccacag acattaatga 240
gtccaaactt caggaactgg aaaagtaccg ggtattcaa actcgtgtcc ttgatgtcac 300
aaagaagaaa caaattgatc agtttgccaa tgaagttgag agacttgatg ttctctttaa 360
tgttgctggt tttgtccatc atggaactgt cctggattgt gaggagaaag actgggactt 420
ctcgatgaat ctcaatgtgc gcaatgtacc tgatgatcaa ggcattcctt cctaaaatgc 480
ttgctcagaa atctggcaat attatcaaca tgtcttctgt ggcttcagc gtcaaaggag 540
ttgtgaacag atgtgtgtac agcacaacca aggcagccgt gattggcctc acaaaatctg 600
tggctgcaga tttcatccag cagggcacat ggtgcaactg tgtgtgcca ggaacagttg 660
atacgccatc tctacaagaa agaatacaag ccagaggaaa tcctgaagag gcacggaatg 720
atttcctgaa gagacaaaag acgggaagat tcgcaactgc agaagaaata gccatgctct 780
gcgtgtatgt ggcttctgat gaactctgct atgtaactgg taacctgtc atcattgatg 840
gaggctggag cttgtgattt taggatctcc atggtgggaa ggaaggcagg cccttcctat 900
ccacagtga cctgggttacg aagaaaactc accaatcatc tccttctgt taatcacatg 960
ttaatgaaaa taagctcttt ttaatgatgt cactgtttgc aagagtctga ttctttaagt 1020
atattaatct ctttgaatc tcttctgaaa tcattgtaaa gaaataaaaa tattgaactc 1080
atagcaggaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaact 1140
cga 1144

```

&lt;210&gt; 21

&lt;211&gt; 1443

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 21

12

```

ggaaccattg gcctataattg ggttggatct attattatga gtgttggtgt ttttgtgcca      60
ggaacattg tagggaagta tggpaacacga atttggcctg cttttttctt aagcaaacca      120
tatacttgtc ttcctgtctg ggctggtttc agaactctata atcagccatc agaaaattat      180
aattaccctt caaagggttat tcaagaagcc caagcgaaaag acctgctgag aagaccattt      240
gatttaattg tgggtgtgtg tctcctcctg gcaactggat tttgctgtt cagagggttg      300
attgctttgg attgcccac tgagctctgc cgattatata cgcaatttca agagccctat      360
ctaaaggatc ctgctgctta tctctaaaatt cagatgctgg catatatgtt ctattctgtt      420
ccttactttg tgactgcact gtatggctta gtggttcctg gatgttcctg gatgcctgac      480
atcacattga tacatgctgg aggtctggct caggctcagt tttctcacat tgggtgcatct      540
cttcatgcta gaactgctta tgtctacaga gtccctgaag aagcaaaaat ctttttttta      600
gcattaaaca tagcatatgg agttcttctt cagctcttgg cctatcgttg tatctacaaa      660
ccagagttct tcaaaaaaac aaaggcagaa gaaaaagtgg aataaaaaata ttacttcatg      720
ttcctccttt ctaaaattact aacttttgtt atactggtag tgatattttg tcccatttca      780
ctctcttctc atacgtgagt acttaagaat atgtacattc ttgctctgca ctgtatgtgt      840
gagctatatg gtattgtgta aatttttttt gaaggaaaat ggaaattctt gagaaacagt      900
ttgtttaaag aaatatattc aaaatcattt gtgaataaac ttgatcatcc atctcaatat      960
tgtttgacat ataaaaataat tataagtgtg aaaaattaca atttagtgcc aacagtagtg      1020
agcatgaaat gaaactattc aaaagagaat atggcctgtg catattaaaa aattcaaaac      1080
agtgaatgca gactggagga gtaacttttg caaataagat gaatatgctt cattattaaa      1140
ctcaatataa aaggcaaate atcagaatat ttttaaatgt tgtttgaaaa atgttttccc      1200
aaggaaaagt tattatttgc tgctgtttca agaaaattac ttttactaaa tttttttgtg      1260
tgaattttaa cagctaaata gggatcagta actttatctc tatccttaat gaacatttgt      1320
tttattgggt gotggaaata tttctattgt atttctgtgt atatttttaa taaaattatt      1380
tttggcctct taaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaactc      1440
gag                                                    1443

```

&lt;210&gt; 22

&lt;211&gt; 1053

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 22

```

cttcgctggt gggaagaagc cgagatggcg gcagccagcg ctggggcaac ccggctgctc      60
ctgctcttgc tgatggcggt agcagcgccc agtcgagccc ggggcagcgg ctgccgggccc      120
gggactgggt cgcgaggggc tggggcggaa ggtcgagagg gcgagkcccc tgtgtctctca      180
gccatcccaa gaagggtttg ctggtccctc ctttcccccc gtcccacgag gccacctggg      240
ccagccccct gtccctctgcc ttctgctggc agaggagcag ctggactggg gcctttggca      300
cagcagccgg tgtctcctgc gccgcctcc cccatggccc catgcagccc caggggcttc      360
ccccctgccc atggagtaga gcccgagatc ctggccacta tgccagttct gacctcgcat      420
ccccctaccc cgagcccatg cagtctggga acatgccgcc ttctctccag cctctgtgccc      480
tttgttccag gtggtctcac cctctgtcc ctgggtgggc taggtgggtc tgtccaggct      540
cctgcagcgc cccctcact ttgacactgg actaggatgc agcctccctt ctgtgtcccc      600
ttgagggtag cctgggtccc ctcatcaggg gcagaggcat gaaagagtcg gggctggatg      660
gccaggggct tctgggcccc acgcctagtg cagcccctgg ggtcgtggtt tgacatttgt      720
ctgcctgggt caaacaagga atccttgctt ttaaggtagc agggcctcca caggcttcca      780
gacttgaggg aaaaggttta agaaagaaaa caaaaccaac agttagtggc cggtaggggc      840
ccaggtggtg cagcgtcccc tcttgacac ccaggggccc ccttttctgc tggagtcccc      900
tgtgtcctcc accaccccc gccgcccagc atcctacctg gactgcgggt ctacgagggc      960
ctgccccgct ttgctgtgtg ccaccctccc tgtaagtcta tttaaaaaca tcgacgatac      1020
attgaaaaaa aaaaaaaaaa aaaggcgggc cac                                                    1053

```

&lt;210&gt; 23

&lt;211&gt; 741

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

```

gtagacttct ggaccttgra tgtcactgag ttttaagagaa aggtagtccc aggactctgg      60

```

13

tctagcagct	gagtaaggct	caggccttag	aatggcttag	cttgetccat	tgcaatgcat	120
catcctggat	gttaaaatcc	agctgtctct	ctgaaamcta	aatatgaaag	actgagattt	180
agtcaacttt	gctgagattt	aatttacata	taataaaatg	aatcatttta	actgtgcagt	240
tcagtaagtt	ttgaaaaatg	tctatacaga	ttcatgtaac	tgccaccaa	attgagaaag	300
gacacttcca	ccatctcaaa	agattctgtg	tggtcctttg	tagtaagtct	cttctacccc	360
atccctagac	aacagctgat	gtgctgtcat	tgtacacata	tattagcttt	gcctgtccta	420
gaacttcatg	ttaatgggaa	gcacccgtga	tgtactgttt	tgtgtctggc	ttcttcagtg	480
tatttttgaa	atztatccac	attgttgtgt	gtatccaaag	tgtgttcttt	ttcattgcca	540
aataatgttc	tgctatatga	atatattaca	aaatatttgt	ctgtttatct	attggtggat	600
ttttgcattc	gttccagttt	ggggctatta	tgaataaagc	ttctgtgaaa	attcaaaaaa	660
aaaaaaaaaa	aaaatgacct	tcgagggggg	gcccggwacc	caaaacggag	tatttccttt	720
tttccccccc	cccgtcccc	g				741

&lt;210&gt; 24

&lt;211&gt; 946

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 24

ggcagcagcg	aaagcctctc	tcttaacaac	ggtgccgcac	agctttgccc	ttgaaagcat	60
ctctactgga	ccggaacaca	ctcatgtgcc	ccgctccctg	acccagccaa	ggctgccctt	120
tcacttccaa	ggctgagatg	ttgccgggtg	tcccatgaga	gcctgcccac	gggctcaggt	180
gcccctttac	cttctgctgg	atggacatct	ggctgtgagc	caggctgggg	tcattggccg	240
ggtagagcga	ggcaggggtg	gacggaggct	tcgagggccc	atcactagta	gggtcattac	300
ctcttgccaa	cagccggggg	tgggagtcct	ggtctcgctc	aggccagagc	ttctcaacct	360
ggagtcctct	ggggtggctg	ccaaagggtg	gtatgacaag	cacgtatccc	tggacatttc	420
cggggagagg	tctggggctt	tggtcacatt	ctccaagggc	tgctgggctt	cggagcagtc	480
cccccccatg	tctcagccac	tacagggtcc	ctctctctcc	ttgcacccca	gacctccgc	540
tgccctggta	atgagcagaa	ggaaagtctt	ggggtgtgct	caaagtcagg	agagcaaaat	600
atgccaggca	aaagctcccc	ggaaaagccg	gaggagtctg	gggtggccac	cgggatgtgg	660
agcagcgagg	gcaaagacgg	tgaacacagc	cctccagctg	tctgagcctc	agttttctaa	720
tctgtagaat	ggggatgatc	atacctgect	cacaagaatg	ttgagacaat	tcacagagac	780
gttctggagc	ccctttcccc	cgagaccggc	attcatgagt	ctgctgggac	cagaaaaacc	840
atctcagggg	cccagcgggg	cacccaggag	agtctggcgg	tgcaagcgct	gtataaacca	900
caagcgttct	caaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaa		946

&lt;210&gt; 25

&lt;211&gt; 831

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (5)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (10)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (11)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

14

<221> SITE  
 <222> (15)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (27)  
 <223> n equals a,t,g, or c

<400> 25  
 catcnacggn naccnctact ataggtnaag ctggtacgcc tgcagggtacc ggtccggaat 60  
 tccccgggtcg acccacgcgt cggggggaaw tcccagtcga tttttccaag cagtactccg 120  
 ctctcctggat gtgtttgtct ctcttggtcg cactggcctg ctctgctgga gacacatggg 180  
 cttcagaagt tggcccagtt ctgagtaaaa gttctccaag actgataaca acctgggaga 240  
 aagttccagt tggtaaccaat ggaggagtta cagtgggtggg ccttgtctcc agtctccttg 300  
 gtggtacqctt tgtgggcatt gcatacttcc tcacacagct gatttttgtg aatgatttag 360  
 acatttctgc cccgcagtgg ccaattattg catttggtgg tttagctgga ttactaggat 420  
 caattgtgga ctcatactta ggggctacaa tgcagtatac tgggttggat gaaagcactg 480  
 gcatgggtgg caacagccca acaaataakg caaggcacat agcagggaaa ccattcttg 540  
 ataacaacgc agtgaatctg ttttcttctg ttcttattgc cctcttgctc ccaactgctg 600  
 cttgggggttt ttggcccagg ggggtgaactt tatttcattt ccmcagggtg aaactgaatg 660  
 ggcagttcat gktaaaatcm cttttcatgg aaagagctct atgtaadagc ataataaaac 720  
 tgsctaccta gcagcaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaagggcg 780  
 gccgctctag aggatccaag cttacgtacg cgtgcatgcg acatcatagc t 831

<210> 26  
 <211> 1294  
 <212> DNA  
 <213> Homo sapiens

<400> 26  
 ctgcagaatt cggcacgagg ttatttcacc tctcttgccc tcagtttctc tgtgaaatca 60  
 ggaattaaca tgggtctctga gaccccttcc tgatggtgaa tgtgtggttt ggtgattttg 120  
 tggccctgca tcatgacctt atttagttct ctttcaacag gggatgtttt actgccttgt 180  
 aaaatcctcg tgggactgcg tgtctttata ggagccaggg tgtaaatgaa cagaattcag 240  
 attggttcta atatatattta cctctaaaag aaaggccatg gggaggccat gaccttaaag 300  
 cagggtttttt ctgttgtctg tgaagcctgt gatgattgag agtggctggg actggcggga 360  
 cgatgttttg gtggaagagg gaggccatct cgatgcgccc cgtcccgggg aggcacccag 420  
 cctgtaagga ggtgatgtct atctacactg agcgcaagga ccctgaaccg ggggaggctg 480  
 aggcggggcc tcttgattcc caccctgtcc cccagtggct aggcctagtgt ggcccgggaa 540  
 atgacttcca tctctccctc caggcatatt taataagagg ccagtatttt cagattctgc 600  
 cgcttctgga cgaatgtctc agagagctgg gaggcgccct ggaggatgga acccttccct 660  
 gagcgttggt gaggtgtgtc ggggggtgcc tggcacaggc cccctccctt ggggggcatc 720  
 actgttccct tgcctctgcat ccccgctgtt tccccgtccc ctgaacaggc gtggagatgt 780  
 gcacgggaca ctcgagggcc ggatgctcaa cagagtggag tgcgcgacg gtgtggccgc 840  
 agcctggctg tgccttcacg acgcagctgc aatcagagga gctgtgggac gctgtcccat 900  
 gtggacacag cccactcact ggggtgtgct cctgtgtggt gcgctgact tttattgtcg 960  
 ttaaaaattt atattaagat gcgccggggc atggtggctc atgcttgtaa tcccagcacc 1020  
 ttgggaggcc gagacgggag gatcacgagg tcaggagatc gagaccatcc tggcttacat 1080  
 ggtgaaaccc cgtctctact aaaaatacaa aaaaattagc cgggtgtagt ggtgagtgcc 1140  
 tgtagtccca gctactcggg aggcgtaggc aggagaatgg cgtgaaccg ggaggcggag 1200  
 cttgcagtga gccgagatcg tgccactgct ctccagcctg ggcgactgag cgaaactccg 1260  
 tctcaaaaaa aaaaaaaaaa aaaaaaaact cgag 1294

<210> 27  
 <211> 1656  
 <212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

```

ggcacgaggt tcacagcacc tgatttgcaa ggcagctata caagttcctg gactcttgta      60
gttccggagt gtttcacctg accttaagcc caccatcc atctttaatc aagaaaccat      120
gtgctttccc gcatgcctgt gttccccctt cacgtgtctg ctgtctgtgt ggaagcctgg      180
cctggcgcat gctgtggtgc actgcatgct ggaaccctg gagtttgac gcgtggtaca      240
gtatgaggcg ggtcacgttt tgtagtgtgt gccgtgggct cccgagaaac aagttaaagt      300
gtgtgctgaa atagatttta ttgacataaa ataagcctta ttgctaaatt taagagaatg      360
tgttacaaat gttttttgct aaacatcagt attgattatt ctacatgatg tacttattga      420
cataacaacc tgaattctt gatatttagac aatttctcct caagttgatt cagctgcatg      480
actctcagaa atcagtcatt ttttattgta gattgctggg tttcttcctc tagtttgtat      540
cgtgtatttt cctcctgtgg agaaatgtg gttggcaaga aatgccatat tttaaagctg      600
tatcgtggct gttaatgcag aaaacaccag tgtactgcag gctgtttggc agtggggctg      660
gggctgagtg tcctgccctc agtggcctgt gtctgtgctc ttgttcgctg acatgcagat      720
acaggggcag atctgagggt ttgatggagt gcagaaggcc acacgtgtgg ctttctgtaa      780
atgcagaaac atggaatcct tgagcagaca cttgtcttct ggagcacctt gcatggattt      840
cgctcctga tgcttcattg ccgttaatag agtgggtggg gttgtgttat gagaaatttt      900
gtctaacctg gcttctgaaa tttctcaaac taaatattca tgctgttttg tgtttttctt      960
aatgactgag gctagtgata ttactcagaa aagtaacagt aacttgggtc ttctgagcgt      1020
caggatgttc accatttaac ttgtttctcg ttagtgtcta gtacgtcggc tttcggtagt      1080
gtagggtgtg gttctgtgct ctttcccgtg tgtgcctgca ctagtggcag cctctgcttc      1140
caggtcagtt tagagtagac tggctctggg attgctagca agtagttgct gttaccagat      1200
gtagccatga agcccagctc cttggatctt gacatatatg ttccaggcaa agtacgtaat      1260
ccagacgttt ctaactcttt ctatgatgatt gcaattgttc tccatgttgt ctgtaggcg      1320
ttatgttaat tctcgatcta acagtgtgcc tgtaacatat atggtagtga agagacatca      1380
catgcagaga ccgttttctt tttatcaact acaggtccgc tatcgacgag agcacctttc      1440
tgctaggcag tcacctact tcccgttgtt ggaggatttg atgagagacg gcagtgatgg      1500
tgctgtcttc ttagctgtga ttactatta ttgccagag cagatgaaac tggatggtga      1560
gtggagaatg cttcctgaaa cagatccgaa aaggcttaaa ggaaaattat agtgtacatt      1620
gatccacata tatattaaaa aaaaaaaaaa aaaaaa

```

&lt;210&gt; 28

&lt;211&gt; 1350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

```

gtgacttcta tattcaatag atttttgtaa atgttaaaac atctatattt aaatgttaaa      60
aactaaata tagagagggg ctttatttca atcatagagc aacaacaaaa ataatgctta      120
tagctaaact gcctgttcta gaaagcatct gctttttcat gttattccta aatcctcttg      180
tcatactttt gtcattgaac aatgctctcc ctctcgtctt ccatcctcat tcagaatttt      240
tagaagacca caatcgtgga gatacactac ccagtattgt ttgatacatt tttatttgat      300
aaacattcag tgcaggaaac tgtgatttgc tatatgttta tgtatataat cttattctgt      360
agtcatcaga atgttaatgt aaggtagatt tgatttttat tttttacatg tgtagttttc      420
tttcttcaca gtcaaagcat ttatattatt ggggggtggg gcagggaatt aagttggtgg      480
gctcgaaaat ccattcatat gtatctgtct acaaattgtc ggggataatt taaatttgaa      540
acctaagtta tatatagttt ggcaatgtct ttcttcaata tttacaataa taggatgatc      600
tacaagaaaa taagtttctt ttgcaaatt tttatcatac taaagttgtt cttttaattt      660
agcatatcta aaataggatt tagttcagtt tagctcacac aggtgtttgc tgacattcat      720
tggccattta atacagtgtt gagtgttct ccgtaaaagt ataagtgtta aactacgaa      780
gaaatgcaca cgatcattct tgctcacttc tataacaaac ttacataaaa tggatttaaa      840
aattcctact cacagcctaa aacttctgga gttcactacc tttttttcaa attcatagta      900
agatcacctg tgtattttat attttagtaa agccaattat gaagtacaag tatcatacac      960
gtacttttga gctactatta tttgaaaaaa atctgccaaa tagcatcttt aggatataat      1020
tacattttca ctcatctaaa aagtatacaa aaataaaaag tggaaaaagg tatcttctga      1080
atgttcaaga gcacacctata gtgccaaata ataaagcacc atttttttct tcataaccag      1140
gattaaaatt catatatact gcagggcaga catacatatg atagcttgtg ctgattaatt      1200

```

16

taacccatt	tgtaaacaga	tgaaaatttt	atcttcttat	ttcatttata	agatggctca	1260
atgtattggg	aggcttcttt	tttattacag	aaagtgtata	ttgggtatata	ataaatgaac	1320
ttttcaaatg	aaaaaaaaaa	aaaaaaaaaa				1350

&lt;210&gt; 29

&lt;211&gt; 1766

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1743)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1748)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1749)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 29

gctcagcgcg	ctgcccggct	ggggacccgc	gcacctgcag	cgcccgcctgc	tgggccctgc	60
atcctgcctg	ggcatcctgc	gcccggccat	gacggcgcac	tcattcgccc	tcccggctcat	120
catcttcacc	acgttctggg	gcctcgctcg	catcgcgggc	cctgggttcgt	gccgaagggg	180
cccaaccgcg	gagtgatcat	caccatgctg	gtcgccaccg	ccgtctgctg	ttacctcttc	240
tggtcatcgc	ccatcctggc	gcagctgaac	cccctgttcg	ggccccagct	gaagaatgag	300
accatctcgt	acgtgcgctt	cctgtggggg	tgacccgccc	cccccgacc	aggtgcccag	360
ctctcggaat	gactgtggct	ccactgtccc	tgacaacccc	ttcgtccgga	ccctccccc	420
cacaactatg	tctggtcacc	agctccctcc	tgctggcacc	cagagacccg	gaccgcgagg	480
cctgcctcgt	tcctggaagt	cttcccagtc	ttcccagcca	gcccggggccc	tggggagccc	540
tgggcacagc	agcggccgag	gggatgtcct	gctccaatac	ccgcactgct	ctggagtttg	600
ccctctttcc	caaggagatg	ctgctgggga	gctgggtatg	gtggggctct	tccctttaca	660
gacggggcag	atgccaggac	tcagcccatc	ctgaggagga	cacgtgtcct	catggagagg	720
gtgctccggc	ccaggcgggg	gagtcgggtg	ccagtcagca	gctctgccac	catcctgctg	780
ggaactgggg	gggcctctat	tgggttatag	gcaaggcctt	ttctctggca	tgggaattgtt	840
aattttctga	cacgtctaga	tgtgaaattt	ctgaaaatgt	tgaagcagag	aaacattcac	900
acacaaaaag	caacatagtc	atgtgggtcc	agatggcctc	agtcctagat	gttggcaccc	960
tttgctgtgt	ctcctcagag	tatcctgttc	cgctcctgc	cacctggacc	tccctcagtg	1020
gatgtcttcc	ctccccgac	cccagcctgt	cagtcggagc	acagtgcagg	tttggctctg	1080
acttgggcct	ttggctgcag	tgggggtgga	tttcagagcc	tctcatggca	gcattctaagt	1140
gaccagagct	gggatgagag	aggggaaggg	gcaatgtgag	tggcgctatg	ggacgggcca	1200
gccctgctcc	tgagccagcc	cgccctctg	ccccctggcc	ctgggctctg	tgctagggat	1260
ggtgaagaat	gggggcgtgc	carcctgcag	gagtggggaag	caacacgcag	gggtcccggg	1320
cctctcmagc	cttgccttya	cgcttatccg	agctcccagt	gtgggttagca	cagagctcac	1380
ccaccttgcc	tggtctccag	ctggggcctg	tcctcactgg	tgctccaggg	gaagaaacga	1440
cagcctcact	tctgtatgga	ctgctgatgt	ggcctgccat	cctgttcagc	gggcattgtc	1500
tttggagcag	caggagacta	ggatgcctct	cactcacatg	ccagttcctg	gctggccagc	1560
tgctcagggc	tcaggctggg	gcctcccat	gacatcctcc	ccctacactc	cctctctgag	1620
cctccgctgc	ccctcctgtt	gggtaagggt	gttgagtgtg	acttgtgctg	aaaacctggt	1680
tcatatataa	taaataatgg	tgatgaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1740
aaaaaaanna	aaaaaaaaaa	aaaaaa				1766

&lt;210&gt; 30

&lt;211&gt; 2790

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

ccgtcttggg	ctactggcag	ctcctctcct	tgggctcctg	gctgccaggc	gttggtgcca	60
cttctttaaag	gcctggaacc	agggaggaga	ggaaatgcta	ttgttgtggg	ctttctccgg	120
gggtctgtgct	gtgcctgcta	gagcaacccc	tgtacccagc	tccttttgtc	cccaggggccc	180
ctccctctgc	cccaagcagc	cagccagctc	tgcttaggcp	aaatgcacaa	gctcagaata	240
gatctgatgg	tgagctggga	agctgtactc	agagcagagc	aaatgaggga	gggggcgctc	300
aggacccagg	ccctccatgg	gctaattgtg	gtggcagcca	tgctcatgc	cacaccttct	360
tcgcaaaactg	atggaccggg	tgggcctggc	ctgagctggg	gccacaaatc	aaatcaaggg	420
ctccagcatc	cagcctgtgt	gttctgtaat	ggaactgacc	ccctcccctg	aaaacgaaag	480
ggccccgggg	ctggcaatca	gggaaagctc	cacggtgcgc	ggctgtggca	caaacttctg	540
gaaggctggc	tgactggaat	gcagggaaaa	cggcagtacc	tgggaaagga	cccaccatc	600
ttcctgctgc	tgtaactgct	gagccactcg	cagtcccagg	atccgctgcc	accagctctg	660
ccaggcccat	ctcaggtgcc	actccctgag	ctttggggac	agttggcaga	gaaggcctct	720
tgtgtcacg	ctcccccgca	gtccccagcc	cttctgcctt	tctccccga	cactgctgca	780
ccagagtga	agggctatgg	caagggggtg	tcatctgagg	agtattaaga	atgcagattc	840
ctgggcctgt	cccccaaggt	tttggagtca	gtaggtccaa	gggccatact	tttgagaggg	900
gtttgggtta	agtatgaggt	gaaaaggagg	atggctcagt	tgagagaggg	tgacccact	960
caccagggtc	cgcaccagct	gtctctgccc	ttgggcatcc	accagtgct	gccatgcpac	1020
tgccaggcac	ctggcctgct	gggaaccccc	cagcccgtga	agcagtgcct	cgaggcaccg	1080
gcgctgcagg	tacttctctc	tgatggccaa	gagcatcgtg	acccttcagg	gccagaagga	1140
gggcagagcc	atgggcctgg	gcctgctttt	ccaggatcct	gcaggaacga	gcactggcca	1200
gagagggccc	agctgtagcc	atggctcagg	caagcccctc	agcccttgcc	cccatccctc	1260
ggaccacca	aactgcacac	acagctcctc	ttaccgtagc	ctccgtttat	gggccttgct	1320
ttgggctttg	caggctctgg	gctcagggct	ggagtgcgct	cttggtccct	ggctccctcg	1380
ccacaggggc	aggcctggga	cccagctact	ctgtccaggc	cactgtggcc	agagctggaa	1440
ggcagggcag	agggaatgtt	ccctgcaccc	tggaagggg	agttgagtca	caagaggtta	1500
agggtgggtcc	aggaaaggcag	ctgctcttag	tgcccgccta	ggagttgagt	acagttagga	1560
gggtggaggga	agggtgctgag	cttagccttg	tgccctgccc	ccatctcccc	aggcctccag	1620
cctctcccgg	ctgcctgcc	cccaaagaga	aatcacaggg	gcggggcagg	aatgcaaagt	1680
gttttctcag	aacagctgaa	acattccgaa	gagggaatgg	atggggagaa	tggtcaatac	1740
acataagacc	gtgtcccaag	gagctgattt	ccaggccctc	gaggactgga	gaccgcttca	1800
cccctgcact	tcagacaccg	tttgtcccc	ggggcaagg	ctccttactc	tgagcccagg	1860
ccgttccctt	tggcttccct	cgtccaccca	ggctgcactg	cagtgatggc	gcgggaggca	1920
ccagctctgt	ggcctgtgtc	cagcagctgc	gggtctgaag	gaatagccag	agaggagcac	1980
ctgaacccca	tgggcttgga	cttcttggg	ccccgctggg	atttcttcgc	tgctctagct	2040
ggcaggacac	atcccgccct	cttcttccac	ccattccccc	atgtggctga	agacattcca	2100
acaatgggggt	ggggccataa	tagttagccc	tcagtcagtt	cccggagcac	agccctggga	2160
gggggctatt	tctctcccca	ctgaaaacat	ttcaaagctg	agttacttgt	ctgaggcctc	2220
atccctcgga	agccgtctga	ctccagagtc	tgagcccccg	gctagtaccc	tatagagagg	2280
gggctctcca	aaggggctgc	tggggcatgt	gtgcctgtgg	cagaaaagag	gagaccctgg	2340
aattcagcac	cctgggtgcc	attcccagcg	tttagtttct	agaggcctca	gtttctccat	2400
cagcttatgg	gaccttgctc	tttactgaca	agaatggaat	agaaatgtaa	aagtactctg	2460
aaaagcaatt	gccctgtaac	ttatctagaa	agaaaagacc	ctgagactcc	agaatctgct	2520
gttgccatag	ccccatatgt	gtgaattctg	caactagcca	aggctagttc	ctttcaattc	2580
catttaaaaa	acaaaaacca	gcagggtgtg	tggctcatgg	cgtaatgggc	ctgcccattg	2640
ctttgggagg	ccaaggcagg	tagatcgctt	gagcccagga	gtttgagaca	agccctggca	2700
acatagtgag	atcccatttc	tacaaaaaaa	aaaaaaaaag	gaattcgata	tcaagcttat	2760
cgataaccgtc	gacctcgagg	ggggggcccg				2790

&lt;210&gt; 31

&lt;211&gt; 1417

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens



18

<400> 31  
 tttttttttt attaaaaaaa tttaaaaaat tataaaatga tgcctatat gagtttata 60  
 catgacgttg gaggagcata gagatagacc tagactaggc atgtgtatgt gtgtgtgtgc 120  
 atgtgtgtat, gcatgcatgc ttatgcatgt gtgtgtgcat gcatgcttgt gtgtgtgtgt 180  
 gtgtgtgtgt gtagagcctt ggcatccccg acagagcaaa gacacaggag ggtggcacat 240  
 ggaagaacaa gtgactccac cctcccttgc acagttaaaa tctggccaag tgagagggga 300  
 gatgggagag gggagagggg agaaaggaga agaggcactg actggagggg ctgaagcttt 360  
 gtccctcctg ggcaggcgtt ctccatccac acccctcttc ttgtagatag aggataagca 420  
 ggccaaagat gcacgaaacc tgagttccac ttagctcca gacttctaga aaagtcaaca 480  
 gcccctgtat ctctagctga tctctgttg ttcaatgtct gcattaccgc actgggagac 540  
 acttgacaga ttgggcctgc cgcaggccat agcagacatt ggcagccct agaacgaagc 600  
 tgactgtcct tggaaatgtgc cacaggggtg tgacgccccg gccaactcca gtgtgccta 660  
 aaatggcctc, ttgcaacatt cccctctctt catcttaaat cagggaactg aagccacaaa 720  
 atggcaata cacagttctg gcagtdgtt tgagtattgg agaaatcgct ctggccatct 780  
 gttttgtctc cagcatgttt ctacggaat atccacggat atatccatgg atataacaga 840  
 catcctgcga aggcagagct tggctcttga gaactcggca agctcagtc ttgcctggat 900  
 tctgcctca tgtcccatcc agtgtttgga gaaaagctct gagagaaaga tgaatgtctg 960  
 aggccacaca gcctagaagt agtcaagagc acaggctcta gaactagccc cactgagggt 1020  
 gaaatcccg caccagcgcc tgcgggtgt gtgatgtagg agagcttctt accagctctg 1080  
 tgcctcactt gtctcacttg taaaatgaga ataagaattg gccgggctcg gtggctcacg 1140  
 cttgtaattc cagcacttcg ggaggctgag gtgggcggat cacttgaagt caggagttca 1200  
 agaccagtct ggccaacgtg gtggaaacc cgtctctgcc aaaaatacaa aaattagcca 1260  
 ggcggtggtg cgggcacctg cagctctcag tactcaaaag gctgaagcag gagaatcgct 1320  
 tgaacctggg aggtggaggc tgtcagtgag ccaagatcac accactgcac tgcagcctgg 1380  
 gtgacagagc aagactctgt ctcaaaaaaa aaaaagg 1417

&lt;210&gt; 32

&lt;211&gt; 1906

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (617)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (940)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1461)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1901)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 32

tctggagctc caccgcggtg gcggccgctc tagaactagt ggatcccccg ggctgcagga 60  
 attcggcacg agcccagctt ctgggtggtg ctggcaatct ttggtgttcc ttggcttgta 120  
 gatgcatcat cccaatctct gccttcattt tcatgctgca ttctccctgt gtgtgcatgg 180  
 ctgtctctgt gtccaatttt tcccttttta taaggacacc artcatattg gattagagcc 240  
 caccctaatt acctcatctt aacttgataa tctgaaagca ccctatttcc aaatagggtg 300  
 acattcarag ggtgttagga cttcaatatc ttttgagggg acacagttca acccataaca 360

19

```

cctaccaacg gtttctggaa tcattctctca aaataaacta tctaaactct aatctttgwt 420
tcaggggtcag cttctagcag aactcaatgt aagacaccct tttaaagatg gtacctcaga 480
ggtacagaca gattgtgaac cttccctagc acagtgtaac aagtccttg caaaaatcct 540
gatttgagtc aacattgtaa tttcttgctt aaatctaaga atatgcctyc cagcttcttc 600
caagactatc tgggggnagt tgkttctagg gtggctcaat ttattcgttt tgaacgctga 660
tggccgtgca ccacaccgtg gctcaatggg tgttgaggac acgttcacat gcaataaatg 720
cacagggctt tgccattagg tggcattagg ggaaggacac ttcagcagata ctgcagaggc 780
attctcttcc agcctgggtt cctgtcaagc acctgctgag atgactggcc cagacggaga 840
ttgtgggggt gctgagccag ccctgtcccc tccttatctc cggaagatgg aataaatgcg 900
tgtcagaagg gaggggtgct cccatctcgg grgcaccggn gttgcecttc agaaaaacat 960
tgccctgcaca tttttagtgc ttgaaatgaa tctgagtggc aattcaagcg ggcagagctt 1020
gttttggatt ttagacagtt ctacctgctg gctcctcttc tctgctccag ctctgacatc 1080
tcggctccac atacagtggc ctgaagtggc atacgggacc tgagaagagg agaggcaaag 1140
gwrvcagctg tggctggccc ctctgccyt ctggtctcct tggctgggtg gggaagaaca 1200
aacacagttg tgtttccagg tcacagctgc cagggctcca cctgtggggt gggtagctcc 1260
agttctgttc tgagttaaga aatgctgcaa atacgttctc cttraaagag mcctaggaat 1320
tgccatttct ttctgcagct ttctgtagcc aataagcatt tgaggaactg ragaagggtt 1380
cagccctgaa ttgcaaggga aaactgtgtg agtgtgtttt agttaagaaa aaagttaatt 1440
ctagttagac ctgcttgttt ncaaaacaga tgtataggcc agacagatgt acagggatga 1500
ccttgacttt cttttgtcat tgcaggaagg tgggggtatgt atgaccctcg gtttaagacca 1560
ataggaggcc gggcgagtg gcttacgcct gtggctccag cactttggga ggccggggcg 1620
ggtggatcgc ccgaggtcgg gagtttgaga tcagcctggc caacatggag aaaccccgctc 1680
tctactgaaa aaaaaaata cagaattggc cgggtgtggg ggcattgcctg tggtaggggc 1740
tgctcgggag gctggggcag gagaatggc tgaacccggg aggcggagggt tgtggtgagc 1800
cgaggtcgcg ccattgcact ccagcctggg caacaggtgc ggaactcggg caaaaaaaa 1860
aaaaaaaaa raactcgggg ggggmaccga acccggtcgc nacatt 1906

```

&lt;210&gt; 33

&lt;211&gt; 543

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (367)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (376)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 33

```

ggcacgagaa aatattgact cctatctggc cttcatcaac tgacctcgaa aagcctcatg 60
agatgctttt tcttaatgtg attttgttca gcctcactgt ttttacctta atttcaactg 120
cccacacact tgaccgtgca gtcaggagtg actggcttct ccttgctctc atttatgcat 180
gttttgagga gctgattcct gaactcatat ttaatctcta ctgccaggga aatgctacat 240
tatttttcta attggaagta taattagagt gatgttggtg gggtagaaaa agaggggagtc 300
acttgatgct ttcagggttaa tcagagctat ggggtgctaca ggcttgtcct tctaagtgcac 360
atattcntat ctaatnctca gatcagggtt tgaaagcttt ggggggtctt ttagatttta 420
atccctactt tctttatggt acaaatatgt acaaaaagaaa aaggctcttat attcttttac 480
acaaatttat aaataaattt tgaactcctt ctgttttaaaa aaaaaaaaaa aaaaaaaaaa 540
aaa 543

```

&lt;210&gt; 34

&lt;211&gt; 1452

&lt;212&gt; DNA

20

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (283)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (596)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (607)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1275)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1284)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 34

gaattcggca	cgagattaag	ttgtgcactt	taattgggtg	aattgtacat	gtragttata	60
tatctractg	tagttgtwat	taaaaaacia	caggaggcca	tgtgggctgc	taggagtagc	120
aatgtctgtg	cccagccagc	aggtagagac	cagggctgga	cagagkagta	tgggctgtgc	180
tgcagattat	ttgtggtacc	caactgttgc	ataaaacagg	gtgtgatctc	ttgcattgct	240
atgcatgagt	ggattcccag	taaattgtgc	caggctgcct	gantgatgtg	tggcttgtgc	300
tttgatcgt	aatgcttacc	tatgctactt	aagttacata	ccctgtggcc	tttgtggcca	360
ggactgtggg	ctactacctg	kagtgattcg	ttaggggaaa	ggacccacag	cctgtgcagg	420
aggaaaaaag	catctctgag	tacagggtgg	atgagctgga	tgagctgccg	ggcaagagcc	480
acgcacaccc	aggtggtgag	tcttaaggat	aaggtggaat	ttgccccata	gctgtcctgg	540
acagaaaactg	cccagagaag	aatgaatgga	ggacataggg	ctctgtggtc	ccaccntttt	600
ttggganacc	tgtgactggt	cctgttacc	tgtcaactta	gccccaaacc	catctctgat	660
tgacttggtt	gcttattttg	gcacattctt	gctccacaca	gccacataca	tactggctgc	720
tcctcsaagg	ccaggcagat	gcagcagctg	ttgggccagc	aaagagggaar	gtcctggaag	780
gttctggcct	gaacgctgca	tctgttgtgt	gacagccaca	actgctcagg	cttccttgtc	840
tgtgggtgca	ctgtggggag	gagtgttatg	ataagaacat	tggctctcag	tctccctggg	900
gagaagtgtg	gcctcacgtg	ggatttgggc	gttgccctta	ggaaggctct	ctgcatgtct	960
agttccagtt	tgtactggga	agaattaaaa	aagtctgcc	gcttctttag	tttgtcctgt	1020
cttttgtgat	gattctttct	gagatcccc	cctatcagct	caggagtggg	attttctgga	1080
gaaggaaagt	gtttttctgt	tcctcactgc	tcaccttggg	gcattcagga	acatgggcct	1140
gatgaatttg	cttgaaggca	gtctgtaatc	ccatcacttt	gggagccaaa	gargcggtac	1200
atttgagctc	aggagtgtga	gaccagcctg	agcaacgtga	caaaaccctg	tctccaccaa	1260
acaacaacaa	caacnacaac	aacnacaaca	acaactacaa	caaactgggc	tggatggcac	1320
gctcctgtaa	tcccagctac	ttgggaggct	gagatgggag	gattgcttga	gcccaggagg	1380
tcaaggctgc	agtgagccat	gattgcacca	ctgcattcca	gcctgggtga	cagagggaga	1440
ctgtttcaaa	aa					1452

&lt;210&gt; 35

&lt;211&gt; 2908

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<220>  
 <221> SITE  
 <222> (1653)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1655)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (2850)  
 <223> n equals a,t,g, or c

<400> 35

gagctctaga. ctgatcgtag taagttttgt gacttaacaa tgaaagaaag tagaaagatg	60
ctttggggtt tcaaaatggt gttttttaaa attgttcttt ggggtgaattt actcagtgtt	120
gctttgagtt gtatacagaa acaaatgttg ggaattgctc ctcaaaaatg tgttcctaag	180
ttgtgttttc aactttacat catgaggtga agcattaggg aaagagattc tttcgatttg	240
tttaataatc taattataca gaccaaagtg ctttacgttt ctgctctatg ttaatgtbtt	300
agaatgggtga ttttgctgat taatttagac ctggcaattg aaataatgtg ctcagaataa	360
taacatgggt atagtctctg tatgataaag tattcaattt cagaatagtg ttggaatcct	420
gagtttgaat aatgtgttgt atttagaaac atagccctat ctgtttttaaa caaataattt	480
gttgccgat tgtccatggt tgagcatgac aaaaatacct cgtcgaagg caagcttagg	540
taactgctgg caaaacactg ggtgcactat ttttctggat aaaatttata gttattttct	600
atattaccct tcaaaaggga tctcttcagg ttaaaaatca cgcttatgct gaagcttta	660
tctggtgtta actaaaaatc tcatatgggt tcataaccga ggctactaat aattcatttt	720
tactacttgt aaaaatttgc tcaaaattcc aaaaaatat tgatttggtt tttagtgtt	780
ttgcaggctg accccaacct aagttttgat aacatctggt aagtcagtat agttctgtga	840
cttcatgtt taactaagaa ggaaaattca tagatattcc tattagattt tataaacctt	900
caaaagtctg aaacttaatt ttgagtctaa attttctgac actggcccct ttaaatattg	960
taagtctttg ttcactttct taagtataaa aaacatttaa ttactagtta gcccttaact	1020
gggaaactca ggtaatgaac tgcgtacttt tctaaagtcc ttttaactgat caattctgta	1080
tagagggata tttatctaac cactttccgt attttacaag tgccttttct aaaaagggaat	1140
aactattata gctctatttc cccaatctct ataggactca tgagagattg cttgtgtaaa	1200
tataaaagca ccatatgtgt tcttaactcc tatggctgct tgaagctcat gatgaaaaag	1260
tctttttgtc agttttaatt gttaagtaca gaacaaacaa ttgtttggtg atggcctggt	1320
tgaaagagag catataaata tatcccagtg gaactcacca aagaagacca cacctcagaa	1380
attattgcat tttctcatta tgtgttggtt ttgatttgct tttgttttta atgcagctct	1440
tttaataataa agattcttga tacagtgaat tctcttattt caagtgttaag ttattcttca	1500
cccaccctt cccctgccat tgtatttccc atctgtttca aggagtcca acaatttaca	1560
ttgcatcgta tgcagtaggt actgcttttt cagaaagacc tggaaaacat acctgctatg	1620
aatattttgt tcagatgtag ccatttaacct ggntntcaag gttgccttct gtggagagga	1680
tcttagacaa aaatcttct tgtatttact tgggttaagt gaagtccaaa tctttacagt	1740
atgctatttc aggatttctg atattaaaaa agaaaaaaca aaatctttat atctcttatt	1800
aacacttccc ccaagaaggg ttgtgctgtt atttattttc tattataaga aaagtccatt	1860
ctttaagtag tttcttttac ctctaactca atttcatacc aaatacctga tcaatagaaa	1920
tgatataatt aagcagcaaa gattcctaatt ccatacattat gaaaagtgtc agcatactta	1980
gtagtgaaca gataaagtca atttgaatat aattccactt tgtttttaga gactaaatta	2040
agattcaatt aacattatcc tatgaattct gaatgtgata atgtgattca aacagtcaaa	2100
ttttattaag ctcttagtaa ctgagtagg cattccatac taacctcaag ttagcaaaac	2160
aaattagtta aacagcttgg ttcttagcag actgcttaaa agatcaagaa aattttctca	2220
tcttttcttt ctacttagaa acattgcaag aaaccttggg cagtcttcac cagacctgcc	2280
atgattttat aagattttagg cctcagtgac atgctcctga aagtctctg ccagccatcc	2340
aaactaagca tccactcatt ccacttccc aaagtcactc accgataaag gtagcatcct	2400
taagttcatt tttgaaaggg ggaggaggat ctccccctgc ccaaaggaat ttttttatca	2460
gaataccttg aaaggggggt atataaattt ggaaaactta atttcttggc tgtgtttgat	2520

22

aacagttcct	atgcatgggt	tttaatgtga	ggtaaatttg	tttctttctt	cagaataacct	2580
ctctccaccc	cccaccttat	tcttctctct	taatgaata	ttttatggga	gctcaaactc	2640
catgacttac	gtgctcacta	agttttcttt	tttccccttg	tttactctgt	ctgtatgtat	2700
gtcaaaagct	ggcaaaacct	ctaaaactgt	caagaaaatg	cgtgaaagtt	gatttgatcat	2760
agtgcgaatt	catgataaaa	tgtcttaatg	ttatttggat	atgtagtaca	tagaaacaga	2820
aaaataaagt	cattttttata	acttaaaatn	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	2880
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa				2908

&lt;210&gt; 36

&lt;211&gt; 953

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

aattcggcac	gagaaccaag	gtactttcag	ctgcagactg	accatacccc	tgtccacctg	60
gcgttgctgg	atgagatcag	cacctgccac	cagctcctgc	acccccaggt	cctgcagctg	120
cttggttaagc	tttttgagac	tgagcactcc	cagctggacg	tgatggagca	gcttgagttg	180
aagaagacac	tgctggacag	gatgggtcac	ctgctgagtc	gaggttatgt	acttcctggt	240
gtcagttaca	tccgaaagt	tctggagaag	ctggacactg	acatttcact	cattcgctat	300
tttgtcactg	aggtgctgga	cgtcattgct	cctccttata	cctctgactt	cgtgcaactt	360
ttcctcccca	tcttgagaaa	tgacagcatc	gcaggtacca	tcaaaacgga	aggcgagcat	420
gacctgtgga	cggagtttat	agctcactgc	aaatctaact	tcatcatggt	gaactaattt	480
agagcatcct	ccagagctga	agcagaacat	tccagaaccc	gttggtgaaa	aaccctttca	540
agaagctggt	ttaagaggct	cgggcagcgt	cttgaaaatg	ggcaccgctg	ggaggagggtg	600
gatgacttct	ttacaaagga	aaatggtagc	agcttcagtg	agaaactgcc	cttacaacaa	660
gtcccttctc	tgctgtcaat	ccaatactgc	tcccaaactc	tgttttcagt	gttcatttcc	720
ctcaaggcag	gcgctgggct	cccacgaccc	ctcaggacag	atctggccgt	cagccgcggg	780
ccgctgggaa	ctccactcgg	ggaactcctt	tccaaagctg	acctcagttt	tcttacaaga	840
acccagttag	ctgatgtttt	attgtaattg	tcttaatttg	ctaagaacaa	ggtaataagt	900
aaatttttaa	aaagcctttc	tgctgggttg	gattaaaaaa	aaaaaaaaaa	aaa	953

&lt;210&gt; 37

&lt;211&gt; 3864

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 37

acccacgcgt	ccgtttctga	agcaatgtta	atcctactag	ccaagcatat	cacttagtcc	60
ccactgtgag	atgagggata	tgtgcttaaa	ttgtgaaaca	aatatatgag	tcagggtattt	120
ttcctttgag	tccaagtggg	ttatgacttt	ctttcctgtg	ttctttgtat	atgtgggagt	180
tttataattt	tttatcaaga	atgaaagggt	ggcctgtgtt	cttactgggtg	caggctgtca	240
cattttctctc	tggtgcccag	tcagggtgcta	tggtcgtgtc	tgcttctggc	gtagtgtact	300
ctgtggatgt	accagcatgt	tcttcaagggt	catgactgat	ttccagacc	tttgggaattg	360
agataaatgt	taaatttgta	gctatctctg	aatttcttcc	agatactttt	cttcatttgt	420
ttgtttgtag	ggtaaacata	cctgatagca	gcaatttaag	cataccctta	gaatgaccat	480
gtatggccag	tgcacctgaa	tgtgtgttcc	aaggtaggga	atccaggaat	ggccaactcg	540
gagattcatt	ccttactatg	ataaatatct	gagccccctg	ctcatcctgt	ggaacatggg	600
cttatttgagg	attaaggccc	tgagtttttag	gttaaatgaa	ggttaccaga	tggagggtcat	660
tagggggagg	gtgttaaatg	aaaatgcctt	ataaaactgca	tgctgtttgc	aagcagttgc	720
agttttcctg	cccagcccgc	agccactggc	catgcagtca	tggtgtccag	cctgcccga	780
ctggaccatt	tctgtacata	aggcagttct	cctgtccgcc	tgccaccagt	tctccactct	840
ctccccatat	gtaagccctt	agtaaacccc	atgtctcatt	tgctgcctct	gggtcttttc	900
ttcagcctct	tgaacctagt	gccttccctg	ctgaggttaa	taggggtaca	gcacaacagt	960
gttgtaacac	agaaagtgat	atttacaggg	atatctctct	cacaatatct	cttaggaaag	1020
gtaaaataaaa	tgttcacaac	ttgtaggtga	gtaattcctt	agataagttg	ttctttaact	1080
tgggaggagt	ttgggaagga	acctaagcag	gctgcagagg	ctgggcagtg	gagcttgtca	1140
tggctggaag	ttgaaatggt	caactccagg	cagatctcct	ggggcaagc	agcctccacc	1200

23

```

accagtagcc cttccttttt gtctttcata cccactgct ccatctgaag cctgaacccc 1260
ttccagaaaa ttgatggata gatttttttt ttccggtata tatagtttta gaggttagaa 1320
ctagatataa ttcagtctag aagatttctc cttcccaga aatgattgtt tttgtgcaa 1380
gccccgcaa, aatagtacgg agacttagac tgagtctact catcactaac aattaacttt 1440
ataaacattc aacaagtagg acaactatta ttactgttac tcagaacctc tcgctctgta 1500
tatacagttt gatttaagat gccacattta catggcattt tcaaccttca aactctagca 1560
gattttaaaa ctaggtggat gaaaatagaa tcattctaata aaatgtagtg tgtcagattt 1620
gaaaaatcat ttggtagca ggatctctgt aaagtatat gggccacgta tacaagacgt 1680
aactgaagaa aattaattca acagagcatg ccgtacttga acgacataga gatttactcg 1740
aactgaacta actcaagctg caggaactccg agcaagcctg gattgtaaaa gtctgggtga 1800
aaatagatgg agtatgccct gactgaacct ctgtactgcc ccacatgctt atacaggtgg 1860
gggattggat ggctgttagg tgatcattgc attctctttt ggatccctat tgagaagaaa 1920
tgataagaga, gggaaaggat atggggcaag aacagtctga aaaagaaagg ataaagttct 1980
cagactctct tcacactcta agaagaactt tctgaaaagc ttggattagg tctggcaatg 2040
gatataataa gcaaaagact cttggaatgt gtctctggct cttagcccca cctctgactt 2100
tgagcaaaac agctgatttc tctgcctgta aaataaatag cctctgata ttaatactta 2160
cctcatgagg ttatttagag gatagtgttg gtaataatgc cttgtgttta catcattcct 2220
ttcacagaga, gctcaaagca ctttacatgc attgagagag aagcttctcg tgaagagtaa 2280
atagaagtgt tcactttttg gaaatgaact taggcataa gagcctgaat ttaatgcatt 2340
gcaggaagaa atatggtaca tagtgacca gtgggtcaac tgaatttttt gttccactaa 2400
gagtcacctc ctggctcctt gttttgtgaa ttgaggaata tggtagtcc ctacacctgg 2460
atgggaaatc ccacatatgc aattggaatg gtctctcacg acacatgcag agattgaaga 2520
acagtctgga caftttttga taacgttctt tgggccttgg tagtagctga aagacacctg 2580
agaaatctta gtcagagct acagaatgac actaatggat ccagaaata gaaatgtaga 2640
tggtagtgtt tttatctgtt tatttcacct caattcaacc aatactcctt gagtgccttt 2700
tatatacatg attttgatg atgtggagaa ttaaaagagc ccacatgct caggaaagtt 2760
aaccctggtt ttagcaagga aaagaagtag gatttccaaa tagataagtg caccgggtat 2820
gtggaagttc agaaaagctt ccagtatatt cagcccatct acttggccat tctcaaccat 2880
gtattactca tgtaccaagc agtatgctgt tcacagagag atccaatctc tgccttaggg 2940
atccttgggg aaaacatgta caaagagata gttttagcac attctagtaa aggcagtgat 3000
caagggcacc tagccttacg tggcaattta gggaaggtag cttggaggat gagacttctc 3060
ctaaagtctt aagaattgaa aagaacatgg gaagggaatt ccaggctggg agagtagtat 3120
gttcatacgc ctcagtgtt taaccttctt tgaacaaaaa aatggccaac tacagaaagt 3180
ttggtcttat ttagcctaa attgtactta ggggtacgag tgagaaacag ggattaagat 3240
aaaggacctg ttttgctgtc ttgtttactg ttgaatagta gtatgaagta ggtcctgaaa 3300
aactatgttt ttggggaaaa aaaaaaaaag actgaatgat atgttgggtt taagtcttg 3360
caggcaggct atccaggtaa ataaacatgg aagtgatgg gaggtaatct gggctggaaa 3420
tacagatttg gaagtcaccc catatcagtg gtgtttaaaa tcaagagcaa atgaaattgc 3480
acaaggagaa tatatagaat gaacaaatta ccatgggtga agccttgagt aatacagaca 3540
tttaagaagc aaacaaaaga caaggaaccc atgagggaga ctggaaagga aaaaacagag 3600
aaataagaaa aaatgagagg agagaattga tacattttcc tcaggtgtgg cattatggag 3660
ttcactggtg tctcatcaga gaagtttcag tccagtggcc agggcagaat gacattgtgt 3720
cttgttttaa agtaaatggg taggtaaga aagttgagaa gggtagcac aaacctctct 3780
ttcaagtcac ttgccttaga agagaaggaa gggtaggtt tctggggtgc aaccagggtt 3840
caagaagcaa aaaaaaaaaa aaaa 3864

```

<210> 38  
 <211> 1411  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (1395)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE

&lt;222&gt; (1397)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1401)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1408)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 38

ccgggtccgga	attcccgggt	cgaccacgc	gtccggcgtg	aaccaccgtg	cctggccgga	60
agtcttttaa	aaataaagt	attctactct	tctaagctta	cagagaccag	accagggtgaa	120
tgtaactggg	gaaaatcaag	atgggtacctc	tctgcattat	cccgccagac	actgtatttt	180
atgcattcat	gtctaggata	cagtgtgaaa	attaaaaagt	ttagagggca	gatgcaattg	240
tggcaagtga	cctgccaaata	aagcagggtc	agctatagaa	gctggcatag	gtatatcctt	300
aatgggtgctt	tctccctggg	cttgtctttt	tggtgttttt	ttcccctata	ttcagagctc	360
cttgagaagt	gataaacacc	tccagctttc	taacatcctc	cccacaccat	ctcaccatat	420
ccatctccca	gcattccatct	gcattcagct	aagggcggga	aactgacctt	gtgcctgtgt	480
tgcagaccat	ttctgaggtc	tccaccatcc	aaggaggcac	agccgtcatt	actgtcctcc	540
atgccttcag	cagcccccct	cacagctaag	gtacatacca	ccccttctgc	cgcgccctca	600
cccttggcac	caaggctctt	tgtgtcttat	gtctaaaggg	atcacctata	tttaactgcc	660
tcagtgcact	aacctctttc	ttctcatgtg	ccagatgtta	agatgaagga	ggaatacmac	720
acatactcaa	gcctcagcct	gttttagttgt	tttctactgg	gctcgctttt	ctgggacggg	780
atattattat	agactggcaa	gcctaaactcc	atagggtttac	aggaagtagg	gatattttta	840
taaaacaatt	gtgtcctccc	cacattttgc	tatgttaata	tttgcttcta	acaatttgca	900
gctgtttcac	tttttctcca	tttgtctcta	agttgaaggc	tttgttggag	gggacagagc	960
acaggaacag	ccttgacagt	ctgtaattat	tgtacagata	ttttaatagc	atataaataa	1020
gtatattcct	tttattttga	aacaaaaatg	atcagacact	gccttttgtg	tgtttgtctg	1080
ctgtggcatc	ctttttttaa	aagactgtta	catattaaaa	tagtgtacat	atataaatat	1140
tacctctttt	gctgtacagt	tgtgatagag	actgaagatt	ttattttttg	tgtgcttttt	1200
ataagaaaaa	aattaataca	ctaaagaatc	ttgctgatgt	gattgtaatg	tacctatgta	1260
acttattttac	ttttgaatgt	tcttctgtat	ctttaaacct	tttattaaat	aagggttttaa	1320
aaattcaaaa	aaaaaaaaaa	aaaaaaaaag	gsgggccsct	ytaraggatc	caasccttgcg	1380
tacgcgtgca	acganancag	ngtcgagngg	t			1411

&lt;210&gt; 39

&lt;211&gt; 1182

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (496)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1162)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 39

ctaagctggg	tcgcctgcag	gtaccgggtc	ggaattcccc	ggtcgaccca	cgcgtccgat	60
aagcacccat	gtctttgaa	atgaatgtat	ttgtaaaaata	ccacgtttca	tgtgtgaaata	120
tgtgctttta	ctgtacatag	tgtattgtg	caataggtct	tatgctgttt	tcactcaatg	180

25

tgtgctaaga	tctagcccca	ttgactcttc	tagaaatgca	gtattgcttt	gacctgccat	240
gtggcactcc	acaatgtcaa	ttgcagttta	cacacattgc	ctaaagtggg	ggacacctgg	300
gtgcccctga	ccccttggca	cgggatacag	gccacgataa	acatcctttc	gtgtgttccc	360
ttctgtgctt	gtgtggcatg	tgtaccagag	atgggcctat	aggtcacaga	ggtcagtttc	420
tctttgggtt	tccagatttt	ctttagaacg	gtgactgacc	ctcctacttg	aggccgcctt	480
ttctccttat	ccttgnccag	acttgtattg	ccagactacc	taatttttgc	cagtctcatg	540
ggtagatagt	ggtscagtgc	tttamcatac	attcatctga	tcagcattaa	tttggggaat	600
tttttcactt	agcctttctg	gtttcccttc	ctgtgcattg	cccatcttct	catggagttt	660
cttatctttt	ttggtttatt	ctcaggagtt	gcttgtacat	tcttgggcaa	ttgcagataa	720
ttccaagaat	gcatatttgg	gctgggtatg	gaggttcact	ggtaattcca	gcactttggg	780
aggscagggc	agaaggatcg	ctgcagccca	ggagttcgag	actagcctgg	gcaacatagc	840
gagacctcgt	ctctacaaaa	aaaaattaaa	aagggggcct	tgggaggcca	aggcgggcag	900
atcatgaggg	caggagattg	agaqcctcct	ggccaacatg	gtgaaacccc	gtctctacta	960
aaatacaaaa	aattagctgg	gcatgggtgg	gcacacctgt	agtcccagct	actctggagg	1020
ctgaggcagg	ggaatcgctt	aaaccagga	ggcggagatt	gcagtgagcc	aaggttccac	1080
cactgcactc	cagcctggcg	acagagcaag	gctccactca	aamaaaaaaa	aaaaagggcg	1140
gccgctctag	aggatccaag	cntacgtacg	cgtgcatgcg	ac		1182

&lt;210&gt; 40

&lt;211&gt; 2457

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1622)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1713)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 40

gtcgaccac	gcgtccgccc	acgcgtccga	ttataataaa	aagtattatt	aggacaggta	60
aactcatcca	tctactgtga	gctgctggct	tggagaacga	cttttgagga	gccagtttcc	120
tgaggagaaa	tgcttttata	aagcactcat	tgctttgtaa	aaggagacaa	aactgatcct	180
aaatgaccac	tccagggttg	ctgattttgt	ttctggctca	tgtctgccta	gtaaaccacc	240
agcaagctgc	tgaaccaggc	tggaaacaac	attgttgcaa	ctgggaggga	catagagtac	300
tgtgaaagca	gcgttccaac	acatcttcac	ttttacaaag	ggataggcag	agacttccaa	360
catagagggt	cttaaaactt	gaggggttac	aatccccttg	caaataattt	tatagctatg	420
gaccctctcc	tcagatctac	agccaacatt	ttcagtgcac	cttaggaggt	cattttagtc	480
cacgcctctg	aaaaagctgc	actccaacgg	ctgaagagca	agccatacgg	ccgagaatgg	540
ggctcccttt	gccttcatga	aagctacttc	ccccaacact	aagactcagc	tgtacgtttg	600
cttagctcag	tcacattttac	attcttcttg	gtgaactgta	ccttttgagt	acctgccttc	660
attttctaga	atcagaccta	acaagggtcag	tagaagcctg	ggcagcagcg	ggcctggaaa	720
gacgaggcag	ccagcatgaa	ctgctgttct	ctccctgacc	acaaggcgtc	gtcttctctc	780
aggggtcaagt	aattgttctt	gttcgccttc	acaggaatgt	ggggaggaag	gacgttaaca	840
ctataaaatg	ctgcgtccta	ccttaaactt	gtactgtgag	aaagctggaa	acttccacct	900
gtacagtggg	tctggtttgc	atgttttagtt	tcatttgtgg	gaactgcttg	tccaagagtg	960
agctcaggtc	agggcagttt	gtgcctatga	gaattagctc	agctatcagg	cagggttttta	1020
gacacctttt	taaaatgtgc	tcgtgttttg	tttgttttgc	ttactgtcag	tcctgggtca	1080
atcaaagggt	tgtaagggtg	agaattttta	tgcactgcta	tatcgcaagt	gcttaaaaca	1140
gagactggcc	cagatgaggc	attcttcaaa	tgtttgttga	aatgaatgga	caaactctta	1200
ggataaaatc	taatttgttg	gcaactgtta	tttgatttta	gaaggcaaac	tgattttatt	1260
ttagagaggg	gaagggggagg	ggaggctcat	tagcctcttg	gtagaaagag	gactatttct	1320
gcaaataaat	aggtttccac	cttaagtagt	gacagtcctt	aacttcttat	tatggagtga	1380
gtcttgaccg	ctttccaagt	tcaatagaag	ttcaagattg	cctctcagtg	attagggaaa	1440



26

ttgaagcttt	taaagctcct	ggtctcagta	attcctcaga	ataaacctct	ttaaaaggga	1500
tattgatgga	aatgtacaat	taccagtaat	tgagggttta	tctgagggga	tggagatgat	1560
gaaatgggtc	cttcttgga	gttggtggca	ttttggcttt	atttttcaca	aataaagtga	1620
anccatttaa	aacgattgac	aacgattata	tagtgccatg	tggaatata	tagatattaa	1680
tttgtggttg	gtttttctgc	ctgctttaa	tgnaatgtat	tatgtttctg	ggttcctttt	1740
ttagctgtaa	aaataacttcg	tactaaagc	atgaaattta	atcagcagtt	gttcttcaag	1800
ttcctgaaag	ctatarragt	ttctcatgac	ttgagtgggt	ttttccctgc	ccaccagagg	1860
agaaagccct	tgtagaattc	tgagtggtta	caagtgttcc	ctacaaaaac	tgaaccatc	1920
agctcctctt	taacaagttg	gcttttttaa	agcacgtaat	tacaatttaa	tggattctg	1980
taaagtgggtg	ctctaggcat	aattttaaat	ctttttaatg	actatatttc	ttcaaaactt	2040
tgaagaaaa	atgtgttctt	tttgcgtcat	cctttgtaag	aagactgcca	acagaggaaa	2100
aaggacttta	caaattaaga	ccatcttggt	ttcatttcca	caaagatgag	aacaaatcat	2160
ggtgttagga	aaggatcctt	agaagaacac	agaatttga	aagcccttgg	tggttatcac	2220
tactatattt	catatttcca	cagaagtgc	ttagccaagc	tctgcatttt	gagcctgctg	2280
actttcattt	aaaaggaatg	aaaggctgaa	aatccaggct	gctgtgtctg	tagataaagg	2340
tcaaaccatg	tttgagttct	tactgttgt	gtccacctaa	ataaaactga	gtaagtaatg	2400
aaaaaaaaa	aaaaaaaact	cgaggctgac	ggtatcgata	agcttgatat	cgaaatt	2457

&lt;210&gt; 41

&lt;211&gt; 1847

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1279)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 41

gactccttag	ctaagcatat	aaggtaactg	gccctgcctt	gctgttttgc	ctcacctgct	60
aacgtgcaca	tgctctgcc	cgtgtacctg	ctgctagtgg	gacatgctga	mttctcagtg	120
ggcctcatgg	gccagaggaa	gttgcggtgt	tccataaact	cagctctcag	aagtgcgtgt	180
tctctgcct	ggaattcgtc	tatctgcttc	aatagctgaa	tgctgcttat	ccctcaagac	240
tcaattctgg	tggcacctca	gttctgacct	ttccccagtc	tctctttccc	ttctccccac	300
aggtgtgtct	cccttgatat	tttgccctgct	tgctacccat	gtccttttct	cctggtgaca	360
gtgctacagc	ctttccttgg	ggaatcaacc	ttgtctgaga	gtggacagaa	attatccacc	420
cctccatcca	gggatgaacc	aatgacctgg	cctaatcacag	tcgctgtgat	tggctcctggg	480
acaggcagga	gactcaagct	aggccacaga	gcacagctgc	tgagactgaa	actactggga	540
aaacaatatg	ctctttctgc	ctgaggcgcc	tagtggtatc	aatggaagcc	ttgaggcagc	600
tgatcatctt	tgcttggaga	gaggtggctg	cctaagaagg	aagccaattg	agaaaaagca	660
ggcagagaaa	aaaaacagga	acagatacaa	caggcaagac	tcagctgatt	tctcatgtca	720
gcaaatataa	gagaaggaga	ctggctcaaga	tagagatgtc	tgatgatact	gagctcccga	780
ctccagccac	tcctgaagtc	attcctaagc	ctctcgttac	acaagccaat	acatgatctt	840
gktagattaa	aaatagtttg	actttggtgt	tgccacttgc	cgtagaaagc	atycacgatg	900
atacactcty	ccgtgtgcty	ccacacccta	atttaaccty	ctycatagca	ctgactcgga	960
ggccctatga	gactgkgaac	tctytgaggg	caggaaatgt	atcttattcg	tctccaaggc	1020
cacgagactt	ggtgtatgat	agatattcaa	tacagggttg	tgaataaaa	aatgaaggaa	1080
tgtttatcta	gaaattaatg	aagcttttca	tttacttttt	tacttcaggg	cctttctgcc	1140
aagaactctt	aagatgcctt	aggatcttgg	ttgcagccag	gtggctgtcc	tccccagagc	1200
ccttgttgtc	agtcctctga	agtcattggw	cctggagata	acagggaggg	caggtcccga	1260
ctgtctgaaa	agtcacagnc	cccggatcct	accattacca	tgctacctgt	tacttgggc	1320
ttcaccagag	gccacactca	tctccgtacc	ccttcccaac	agtgggtgag	ggcaggagca	1380
cctggacatc	aagaatcgag	tgcatgcctg	aacctgtcca	ttacctatgc	ttctgcagct	1440
ttgtctatgc	tttcccttcc	ctgaaacgct	cttcccttgt	ctacttaatt	attcaaagct	1500
cagaggaaaa	aaaatcacct	gtcccaaac	gtcttccccg	aacctgtcag	ggaaccaa	1560
tcaaaagcca	tcagaaggcc	gaggcgggcg	gatcacttga	ggttaggagt	tcgagatcgg	1620
cctggccaac	atgggtgaaac	cctgtctcta	ctaaaaatac	aaaaattagc	caggcatggg	1680
ggcagggtgc	tgtaatccta	gcaactctgg	aggtgacgca	ggagaactgt	ttgaaccag	1740

27

gaggcagagg ttgcagtbaa ctcagattgc accactgcac tctagcctgg gaaacagagc 1800  
 aagactagggt taaaaaaaaa aaaaaaaaaa aaaaaaaaaa actcgag 1847

<210> 42  
 <211> 2597  
 <212> DNA  
 <213> Homo sapiens

<400> 42  
 ggcacgaggt tacacctcac cccctacttt cgctgactcc agaccccaga ggctgagggg 60  
 cagtagatat gctgcctgcc attgcctctt tccttgtagc tgtctgagtg gtgccagcct 120  
 gatcagattq tgttgcaagt ccccgctgtt gccacaatgt cggtcgcgtt tcttatccaa 180  
 cgctgtttct gcttctggtg gtttctgcta aatgcatttt caatcccaag tggtagagag 240  
 aagaaaagga ttgtctttaa aaagtggctt tgaatgtgtc gaaagcaagg ttgcctgctg 300  
 tgtgtttttg acaattgcct cccagagga gcctttgaaa acctcacttt tcaagacgct 360  
 tcaactccta ttgaaaagtg acttaataaa aattgtctct ctgaccatcg cggctctgga 420  
 tgccaggcta gtgattatgt gtctaactct agtctccagg gctagaactt cacaccgtgt 480  
 cttttcacta cctcccaga ggaggtgaga ggaggtgaat agacacttgt cagattcttc 540  
 tgctgtgtgt cactgtgggg ggatgcacac acactggagt ataggcttgc taaagagaga 600  
 ctgcattgct atttcaacta gacacgttta cttttccgc aaatgattac tgagcaccta 660  
 tgggtgtgcc ggtaccatgt tagggcaaga gatctacctc atccttgoat tgtaaacptg 720  
 tagcatccgt ggcaatgttt tacaccactg cctcccagg gtccctgcaa gctgactcct 780  
 tttattttcc ttgtgcatgc tgatgtgttc aaggattttc agtgggctac tgtgagacta 840  
 tctccccatc aggatctctc aaccttgcca ctaggaaat gttgggaaaa gtaggtctct 900  
 gtttaaggag gaagaggggc tttcctgtgc gtgataggat gcttggcagc atccctgtcc 960  
 agtagcacca catctacctc agttgtgaca accaaaaatg tctgcagaca ttgtcaaatg 1020  
 tctcctccct attttacatt taccaaaagt ccttttatgt cctttacagt taaggaaact 1080  
 taggaccagg aagaaacttg ctcaagtcaa acagttaggt ggcggtagag ctttagctac 1140  
 aatttaagtc ttgtcttccc aaccagtggt ccatcttatg gatccattca gtggtttgct 1200  
 cctaaatgct taataaccaa ttatctggga taaacgaaag ccaaacaac ttgctgggag 1260  
 cgtgggacaa tttccatggt gtaataactt ccaccatggc tgggtttcaag ctacttccat 1320  
 ggcgtcactg aatgtgaagt tgggaaggga tctgtacagt tggatctggt gagctggtgc 1380  
 cagccagctc agaatgcccc caagtccatg tgtgctgctg atcacatatt agtcttgcctc 1440  
 tcctctggac cttgttctgc cccccccag acacatgccc ttatcactgt gatgtgctgt 1500  
 ggggaaagca gtcacctggt aagagttttt gataacttct ctggataaac ataaagcagt 1560  
 tagagtggag ttatccccct cttctgttaa atgaaagtgt cttttcacag tctttccatt 1620  
 gaaggaatta gaaaagtcat gggatacctc tggctcttaga gcagaaatga agttccccag 1680  
 aagggggtcc tggacagaat ggtctcttgg gcagggacta gatgctgtta cttaccatca 1740  
 gatgtgtcta ctgtaagatg cttgggaggg atcaaagtgg tgacagtggg gggctgttgg 1800  
 attttaccat tctggaatat aagagaaggg gtgtggctgg tgttttagtc cagatagcag 1860  
 ctgtgaccac tgggagcctc attttccaca cctgcaaaag gagaaatagt gatccccctc 1920  
 ctagtgttgg tgtaaagagt agatgaacct ctgcctgagt ctcagatgcc tgctgccac 1980  
 ttggttgctt agtggatcgg ctggctgtca acagtgtgag cttatcaagc ctaaatactt 2040  
 agttggtctt gactttttca cctcgaccat gactcagtgt catgtgctcc tagtcattct 2100  
 gtctgtgggc caacgttagc ctgggaagca cctgggactg agggagaagc cctcagctag 2160  
 ttattcagcg atccaggttc tcctcctgcc tttgatata cctcattata taacctggg 2220  
 aaacactttg gtgtgactgg aatgtggata ctcccagggg aagggttaga gcactggtag 2280  
 gcatttggtg tttatcatga aggtggtagg aaatacttga agggttttta gcagggatga 2340  
 cacatcatca aatgtgtgtt ttgaaaactt ttttctgcag aggacagggc cacagcccaa 2400  
 gcaggatcaa accagttagg agattgatgt aacagtcccc tcagaaagtg atgaggtagc 2460  
 tgggtgctgg ggtcatgcc tgtaatccca gtcaccgtgg gaggtgaga ctggcagata 2520  
 gtttgagacc agcccttggc aacatggtga aaccttctct ctacagaaaa aaaaaaaaaa 2580  
 aaaaaaaaaa aaaaaaa 2597

<210> 43  
 <211> 3116  
 <212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

ggtgataatg	aaagtgggtg	tggatgatgt	ggtaatactg	gtgggtggga	cattgggtgt	60
ggtgggtgatg	gtgggtgatac	tggatgatgt	ggtgatgggtg	gtggcgctgg	tgacctgac	120
atgggggtcca	gtagcagtga	cagtggatgc	aggctcctgg	tgactgagga	gcattctcagg	180
ctgrggaggc	acctctgac	cccgccactg	ctccttacct	cctacagtct	ctcagcaaac	240
ctgctgggag	acagcggact	cagatgcctt	ctggaatgtc	tgccgcagtg	cccatctccg	300
gtttgcttga	tctgagtcac	aacagcattt	ctcaggaaag	tgccctgtac	ctgctggaga	360
cactgccctc	ctgcccacgt	gtcggggagg	cctcagtga	cctgggctct	gagcagagct	420
tccggattca	cttctccaga	gaggaccagg	ctgggaagac	actcaggcta	agtgaagtga	480
gcttcgggccc	agagcacgtg	tccaggctgg	ccaccggctt	gagcaagtcc	ctgcagctga	540
cggagctcac	gctgaccag	tgtgcctgg	gccagaagca	gctggccatc	ctcctgagct	600
tgggtggggcg	accgcaggg	ctgttcagcc	tcagggtgca	ggagccgtgg	gcggacagag	660
ccagggttct	ctccctgtta	gaagtctcg	cccaggcctc	aggcagtgtc	actgaaatca	720
gcattctcga	gaccagcag	cagctctgtg	tccagctgga	atttctctgc	caggaagaga	780
atccagaagc	tgtggcactc	aggttggtct	actgtgacct	tggagccac	cacagccttc	840
ttgycgggca	gctgatggag	acatgtgcca	ggctgcrca	gctcagcttg	tctcaggtta	900
acctctgtga	ggacgatgat	gccagttccc	tgtgtgtgca	gagcctcctg	ctgtccctct	960
ctgagctgaa	gacatttcgg	ctgacctcca	gctgtgtgag	caccgagggc	ctgccccacc	1020
tggcatctgg	tctggggccac	tgccaccact	tggaggagct	ggacttgtct	aacaatcaat	1080
ttgatgagga	gggcaccaag	gcgctgatga	gggccttga	ggggaatgg	atgctaaaga	1140
ggctggacct	cagtcacctt	ctgctgaaca	gctccacctt	ggccttgctt	actcacagac	1200
taagccagat	gacctgcctg	cagagcctca	gactgaacag	gaacagtatc	ggtgatgtcg	1260
gttgctgcca	cctttctgag	gctctcaggg	ctgccaccag	cctagaggag	ctggacttga	1320
gccacaacca	gattggagac	gctgggtgtc	agcacttagc	taccatcctg	cctgggctgc	1380
cagagctcag	gaagatagac	ctctcaggga	atagcatcag	ctcagccggg	ggagtgcagt	1440
tggcagagtc	tctcgttctt	tgcaggcgcc	tggaggagtt	gatgcttggc	tgcaatgccc	1500
tgggggatcc	cacagccctg	gggctggctc	aggagctgcc	ccagcacctg	agggtcctac	1560
acctaccatt	cagccatctg	ggcccagggtg	gggcccctgag	cctggccagg	ccctggatgg	1620
atccccccat	ttggaagaga	tcagcttggc	ggaaaacaac	ctggctggag	gggtcctgcg	1680
tttctgtatg	gagctcccgc	tgtctagaca	gatagacctg	gtttcctgta	agattgacaa	1740
ccagactgcc	aagctcctca	cctccagctt	cacgagctgc	cctgcccctg	aagtaatctt	1800
gctgtcctgg	aatctcctcg	gggatgaggc	agctgccgag	ctggcccagg	tgtgtccgca	1860
gatgggcccgg	ctgaagagag	tggacctgga	gaagaatcag	atcacagctt	tgggggctcg	1920
gctcctggct	gaaggactgg	cccaggggtc	tagcatccaa	gtcatccgcc	tctggaataa	1980
ccccattccc	tgcgacatgg	cccagcacct	gaagagccag	gagcccaggc	tggactttgc	2040
cttctttgac	aaccagcccc	aggccccttg	gggtacttga	tggccccctc	aagacctttg	2100
gaatccagcc	aagtgatgca	cccaaatgat	ccacctttcg	ccactggga	taattgactc	2160
aggaaagaag	agcctcggca	gggcgctctg	cactccacc	aggaggaagg	atactgtgt	2220
cctgctgcag	tcctcaggga	gaactttttt	gggaaccagg	agctgggtct	ggacaaagga	2280
gtacctgca	ttacgtggga	tatgtgtgat	caattgggga	catgcgacac	acaatgaggg	2340
tgtcatgaca	atgcatgaca	cgtacgggtta	tatgtggcag	tgtgacctct	tgacatgtgg	2400
cgttacatga	aagtcagtgt	ggcacgtgtt	ctgtggcatg	ggtgctggca	tcccaagtag	2460
caggatacat	gattgtttgt	ctatatatga	cacatgacaa	atgtccatgt	cacaggactc	2520
atggctggcc	agatgacctc	aggctggccc	aagatcta	ttattaattt	ttaaagcaaa	2580
tacatatatta	tagattgtgt	gtatggagca	gctaagtcag	gaaaagtctt	ccgcccagac	2640
tgggagggga	gagtgtccat	gcactgacca	gtccaggggc	tcaagggcca	gggtctctgga	2700
acaagccagg	gactcagcca	ttaagtcccc	tcctgcctca	atcctcagcc	tacctatcta	2760
taaacttgat	gactcctccc	ttacttacat	actagcttcc	aaggacaggt	ggaggtaggg	2820
ccagcctggc	gggagtggag	aagcccagtc	tgtcctatgt	aagggacaaa	gccaggtcta	2880
atggtactgg	gtagggggca	ctggcaagac	gctaagctag	gctactgggt	ccagctacta	2940
ctttggtggg	attcaggtga	gtctccatgc	acttcacatg	ttaccagtg	ttcttgttac	3000
ttccaaggag	aaccaagaat	ggctctgtca	cactcgaagc	caggcttgat	caataaacac	3060
aatgggtattc	caaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	ctcgag	3116

&lt;210&gt; 44

&lt;211&gt; 3460

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 44

acgcgtccga	tgaacttgac	eggctaaatg	ccccactttc	tcagatggct	tctaacgact	60
ttcaggatta	gggccagctg	tgggtctact	ccttgttgga	gcccattctca	cctgggatgc	120
ctgcagccag	ccctccctcg	tgatttgtct	caccttgagt	aggagacatg	cttctccccct	180
aaccttttcc	tttctgccat	aattaacata	tgctcctttc	agtaagtcca	tgctctctggc	240
aggggatgaa	gaagtactca	ctggtaatta	gctaccatct	ttgcagcagc	cctggtaact	300
tgaaaaattt	gggtctgggt	ctgttcatga	gtctttgtgt	aactgcaaaa	gcaggaaaagg	360
aagtcaagac	tcctgttgcc	tcgtgcttag	caaagcagtc	cttatccttt	atactctgtt	420
cttgggtttt	gtttttgtct	tgttttatac	caggcaaat	gcttagtagc	aaagggacca	480
aactgaaaag	gtgacaatct	ctaacttcta	aaagcagaca	ccaatcggat	gtcatttaga	540
ggtaatatgaa	gatgccattc	ttgggtggcct	ctgcacccaa	attgcattctg	gaaagaacta	600
gggtctcatt	cagaatgtcc	aaaaggaaat	tcttaagagc	ttaaattcag	atttgtgtct	660
cattaatgca	gtgaacaatt	caaaaccaca	cagattcctt	ggcaggaagg	ataatggaat	720
aacagtgttg	atgagacctt	tttagcttca	aggtttcgga	gtctaaacaa	atggatgatt	780
catttggaat	gaaactcaca	atgcaagtag	aaggacctct	ccaaatcagg	ccagttgggt	840
tatcctggct	tggaatctgg	tgtgaaacca	taggtcttaa	cactctggag	cagcacattg	900
ctgtggatat	gtccaggaga	ccttagatat	ggcttaaagg	ctttcaagat	gaggacagaa	960
attgcttaca	attgctcagt	ttctcaacag	aaagactcat	aagagtgcc	gcatggggta	1020
catggagtga	agctgggtgg	gaagcatcat	ctgcacagtc	cctgtcctag	tcgaggactt	1080
ttctctgtat	gttttcatat	catgggattt	ttggatatca	gtgtattttg	gttcttgaaa	1140
tagcctaata	gctgctcaca	cattgggttag	gaatattata	ccaatgtcat	ccccaaagga	1200
agggtagct	gaatggaaat	taagcccagt	cattttat	gatctattag	ctctgttatt	1260
agtgcatgat	caccagatc	accctcctca	gcccacacag	tgctgaacca	tcttccctcc	1320
tgttctccat	ggctattaat	agtatagcta	aatttagagt	gcagagccag	atataagtat	1380
tttggaatta	tctcccagtt	tgtggtagaa	gctgactgga	atacaggttg	agtatctctt	1440
atccaaaatg	ctagggacca	gaaagggttc	agattttttc	agatttttga	atacttaaca	1500
gttgagcacc	ccaaatctga	aaggcttctg	aacgtcatgt	cagcactcaa	aaaagtggat	1560
tttgagcac	ttcaaatttc	ggatttttgg	atttggtgatg	ctcatcctgt	gtaggagagg	1620
ctactcgatt	ccattttaatg	actgtcctag	tcataatcat	ccaaagataa	aagccaggta	1680
gatgttgaaa	gctctttcca	gggctgaaaa	agtgttctta	cgttctctgc	atgtgactag	1740
catcactgtg	gaaattaatg	ctctgttctt	cactagaatg	tagtaagtgg	ttaaactgag	1800
ctatccccca	cctgatgact	attggcatcc	atttgcaagg	ccaatggcct	ggattaaggg	1860
ttaggattat	ttgtagctag	aaggtaattt	tatttctgtg	aaactaattg	gtcatattt	1920
gaggttaggt	gtggccttga	ccttaccagt	acatttatac	ccactaccag	ttgactagcc	1980
cagataattg	ttaaattggtg	cttcttttct	gcttctcagt	agacttccat	gccattacaa	2040
aggaaatttg	aattacctag	tgtttgtata	ttccatgata	actatgtata	acttctgtta	2100
cacagcttat	gtattgttaa	catttaagt	ttaaccatgc	cacagctaac	acttaaaaat	2160
gaaaactaat	tagttcttgc	ttagggaaaa	tgccaggat	gaagtatggc	atatacttga	2220
cactgtcctg	tgtaaccctt	tactttgctc	aggctttcaa	gattgagtct	tttttcccc	2280
aaattaggtt	aacatgcatt	tgaccccaac	ctgtgggtt	tgagtaagct	ggaaatctgt	2340
gacggtaggc	tttctagtgt	cacgaggtgg	tggtgactga	aggaaaagct	gggatcacag	2400
gttccttctg	atggagagga	aggtttattt	ctatgccctt	cccaccaccc	tccacctaga	2460
gtcaccccaa	gcctgctcca	gtcccagggg	caggccattc	tgcaaaaagc	ggacctcaca	2520
gaaacaaagg	ctgggttgag	gtcacccctt	tcagagttgg	ttcctggcca	gatgggttaag	2580
aggcatttgt	aattttaaaa	atgtgaaact	tgggttttgt	gttttcttct	aagtgcctaa	2640
ataagcaagc	caggctgttg	atatttttagc	cagagaaatc	ggcaagccaa	gattaacccg	2700
aatctgaagt	ttagaatctt	gagtttgc	ctgcatcata	tcatgctgtt	ttgatgagga	2760
aacatttgcc	actgaggagt	tggagggagg	gcaagacgac	agtgttaagt	cagatcattt	2820
aatggtttcc	cctaagccct	ggaaaaatat	ttgaaagaat	ggcagcaaaa	agggttaagaa	2880
agcaagccag	atttactgca	caatatgcag	taccagtagc	tactttaaat	cccaagagaa	2940
cagtgtgatg	tctaatatat	acaggtctat	gaaaataactg	tggaataaagc	ccaggaaggt	3000
tagatgtgtt	tgcaaaataag	ttgcccataag	ggtccccctc	taagtaaaac	aaatatttcag	3060
accacaggct	ttaatgtaaa	ctgtcaaaaa	gtgggatgtg	gaggattttt	gttaagtgtc	3120
aatcgaagtt	aaaaagcaag	ggtttttggc	caggcgtggg	gtcacgcct	gtaatcccag	3180
cactttggga	ggccgaggcc	ggcaaatcac	ctaaggtcag	gagttcgaga	ccagcctggc	3240
caacatgggtg	aaaccccgct	tctactaaaa	atacaaaaaa	attagcccg	tgtgggtggca	3300

30.

agtgccctgta	gtccccagcta	cttggggaggg	tgaggcagga	gaactgcttg	aacccgggag	3360
gtagagggtg	cagtgagcca	cgatcatgcc	actgcactcc	agcctgggca	acagagcaag	3420
actccatctc	aaaaaaaaaa	aaaaaaaaagg	cggccgctct			3460

&lt;210&gt; 45

&lt;211&gt; 2622

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 45

aggttccctc	ctttggatat	ttggcagttg	tactttatgc	agttcagcca	agaactaaac	60
acaggccttc	aggtgtggtc	agagtacagt	gggccagagg	tcctttcttt	tcttcatgtg	120
gtcatacacc	tctgttaatt	caacttaagt	ttatattact	tttgggggtg	ggagtgggga	180
agaggatcac	attgttgact	tgctggaagc	attatgtcaa	ctaaaatcct	tcagttctta	240
ttttcatcat	gctgttgggt	tccccctatg	ttgtttcttt	ttaaaaacac	caaatgcaga	300
acttcccttt	tatactgctt	ctatttcate	ttgttgactt	gtagcctatc	agagtatgat	360
tctttattgt	catctaaggt	attctgatat	aatcactgca	tgtctaaatp	ctagttattg	420
actaaatgct	acttgagacc	attctcttgg	ttgtcacatt	atttttacagt	tgaagaaact	480
gaaccttaga	gagggttaagt	accttggttaa	aggccacctg	gctggaacat	aacatcctgg	540
tcattttaac	tactaatgct	gtcatgaaag	ggagtttagcc	agggcaggtg	tgaatagagc	600
attctagcat	aagcagccaa	caattagagc	aactgtgtat	ttttagtgtg	tggatagcaa	660
atctttgctg	tcttggaagt	tactatctta	gaagcatggt	attaagctct	ctattcaatg	720
tggtgttcac	atcagtagca	ttggtttcat	ttgggagctt	tttagaaatg	cagagctcag	780
gccccactcc	aggcctaagg	attcaatctg	cattttaaga	agtgacccat	atacacatta	840
aaggtttaag	aagcattgct	gtagtatcct	attttccaca	ctttagtcac	ttgcatatca	900
cttgtgcaac	tttggttcta	tctgcatact	tcctgaaggg	ggcctgcccc	tccacacctg	960
tgggtatttc	tcgtcaggtg	gagatgaaag	aatgagaaaa	gaaataagac	acaaagtata	1020
gagaaagAAC	agtgggccca	ggggaccggc	acactcagca	tgcgaggacc	tgcaccagtg	1080
ccggtctctg	agttccctca	gtatttattg	atcattatct	ttactgtctt	ggcgaggcga	1140
gtgtagcagg	gcaacaggtg	gcgagaaggt	cagcagggaa	acgtgagcaa	aggaatctgt	1200
atcatgaata	agttcaagga	aaggtactgt	gcctggatgt	gcacataggc	tagatttatg	1260
tttcacttta	cacaaatata	taactagcag	agagcaacaa	agcagtattg	ctgccagcat	1320
atctcgcttc	cagccacagg	gcgggtttct	cctatctcag	aatagaacga	atgggaatgg	1380
tcgactttac	actaagacat	tccattccca	gggacgagca	ggagacagaa	gccttcatct	1440
tatctcaact	gcaaagaggc	ctccctcttt	cactactcct	cctcagcaca	gaccttttac	1500
atgtgtcggg	ctgggggatg	taaggtcttt	cctttccac	gaggccatat	ctcaggctgt	1560
ctcagttggg	ggaaacctgg	acaataccca	ggcttttctg	ggcaggggtc	cctgcccgtc	1620
tccgcagtg	attgtgtctc	tggttaatcg	agaatggaga	atggcgatgg	cttttaccaa	1680
gcatactgcc	tgcaaacata	ttgttaccaa	ggcacatcct	gcacagccct	aaatccatta	1740
aaccttgatt	caatatagca	catgtttctg	ggggcacaga	gttggggcta	aatttacaga	1800
ttaacagcat	ctcaaagcag	aacaattttt	cttagtacag	atcaaaatgg	agtttcttat	1860
gtcttccttt	tctacgtaga	cacaggaaca	atctgatctc	ttttcccccac	acttccata	1920
tggttaatgta	ttttaattgt	ctgttagctt	ttatgtcttt	ccaacttgaa	ataatttcac	1980
ttacaaaata	gctgaaaaaa	tagtaaagat	aattcctgta	ttttcttcac	ccacgtattc	2040
ccaaatgtgt	aacacctaca	atcagcaaaa	atcagcacat	taatattatt	acctaattcta	2100
cagacctcat	tcacattttg	ccagttgttc	tgctgggtgtg	ctgtatgtag	tgcaggatcc	2160
agtctaggcc	cgtatgttgc	attttgctgt	catgtctctc	tagtctcctt	tcattctcagt	2220
ggctttttct	tttccatgac	tatgacagaa	atgatttgta	tccttcatat	atctgcagtc	2280
acatgaagtc	tatttgtccc	actactggtg	acgtaaaattt	tggttaacttg	gttgaggggtg	2340
gtaccgtcta	ggtttctcca	tattaagttg	ctatttttcc	ttgtgtattt	aataagtatc	2400
ttatagggaa	atcctttgag	actatgtaaa	tatcatatta	catgtatttt	gcctatttta	2460
gcatcttttg	aagataattg	cctgaaacat	tttttactgt	gatgattttc	taattctatc	2520
attctgcatt	tattagtttg	gaattcgatt	gtaagggaagt	atattccctt	gtttcctatt	2580
attttattaa	agaaattttac	atccaaaaaa	aaaaaaaaaa	aa		2622

&lt;210&gt; 46

&lt;211&gt; 1984

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (106)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 46

```

agttgttgct atgaatgatt ttaagccttt tttttaagc ttggcabaaca tcccagctaa      60
tcaaaatagt catattcctg agaagtagga aactaaaact tctttncata taattgtag      120
aagggttgtt tcccaaacta ccatagttac aaagggtgaat agccaaattt taggaacaga      180
atcaaaagaa taaaaatctg tgaagagatc ctactactct tccttctatg ttttggtttt      240
ggtttctatt gtccctatca tttcagcaag tggaacagca gcaagttttt cagtgcatat      300
gctgcacaag aacaaaatat aaatctgtat ggcacaaaaa atcaaagtga aaaccaaac      360
aaaaacccaa acaccctatg taactatcgg aggcataatac gtggtataaa tgactgtagc      420
tgtgatacac acatggctac ttgtcacatc actttccata attatttact gcaaaatgat      480
tgagaggctt ttggtgcagg cagccgttaa cctcctgctt cctttgttac ctctggatta      540
ctttgcagta aattgcaggt cttttaagag atttaagctt cagttttctc aaaccaaac      600
aattatcctg tcttatctga agatgcaggg ttgtgggcaa aagaggctgg ttataataat      660
gccctcatat tgagtggctt gtaaacggct gcacacttca ggcactgtag ttgctgaaga      720
tgctttgtta aatgtgacct tgactggctt tacaggggtg tagaatgtaa tctacacaag      780
gtgactttgc atctatcttg ctcttgaggt ggatgaaatt gagaagctgg agtgtgtgaag      840
ccatgcacat aagtattctt cactgtaaat tttgttttca tttttaaccc aattatggta      900
ctttatccaa tgcacaactg atctctcagt agatattcat ttgaaaatag tgtggccttg      960
atcagtgaga aagggaagga gaaaagtgac ttttttgctt atgtagaat gactcatttg      1020
ctgagagttt gtctttctgc agcactcttg gtataatggt agtgatcggg ctcctttttg      1080
attggggaaa gttaatgttt ttgaccctgg agttaattca gttgagttat cttatatttt      1140
taggaagtat cagaattgct ctgatgaata acaaagttga ctgttttgat gtccaatctc      1200
aggtttttaga atatagtggg gtaaagtccc actattttta attcttaaaa caactttaat      1260
ttcgtacacc ctaaaagtca catgcataag gcctgttcag agagcagagc ctccatcttt      1320
ttgctccttt tctactttgt acttcacttg raaaaatata aagtgacttt acatkgtata      1380
tttccattgt aaccctgaca tttctcaaag ataaagcact ttttgatcat gaaatacatg      1440
aaatctttgt gtgatgtgga tcatagtttc tcaggctccc ttagataatt gcttatgaat      1500
attgttctaa ctctgtgtaa gaagagtaga aatctttgct aatgttagaa ggtttgtatt      1560
attgatccag aatgcatttt gctagtttcc aatggatggg agagtaata atgctgcatt      1620
cacaatttaa taagttactt tcccttgagc ctttaaggtaa ctttttcttt tctgtcaact      1680
acagcactga agttatagta agtgaatgag attatcagtt ttcagggttg gtttttagagt      1740
actgtaaate aattagctgt cttcctaaag agttacaact ccatttcagt atactggata      1800
atgggtgtgt ggggtgggct ggggagggcg ggagatagtt tgtagaaaag aaaaaagaaa      1860
aaaaaaaaaa acacattttac cttaagaaaa tacagacaaa aaaaaaaaaa aagggcgcc      1920
gctctagagg atccaagctt acgtacgcgt gcatgcgacg tcatagctct tctatagggt      1980
cacc                                                                 1984

```

&lt;210&gt; 47

&lt;211&gt; 1987

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (442)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (444)

&lt;223&gt; n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (473)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (493)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1011)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1025)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1111)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1119)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1169)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1234)  
 <223> n equals a,t,g, or c

<400> 47  
 ggcacgagac cggcgctagt cttgcgtcgc acagcaggat tgcgctgctt tctgatgacc 60  
 ttcgacatgg atggcgatgg taatgaacgc ttttcccgtg acgccccacc cgcaaccgac 120  
 ccaggccgga atgacctga tcgaggtcct ggtatctgtg ctgatcctcg ctgtcggcct 180  
 gctaaggggc gcagtcattc agctcaatgc attgaaatac accgacagtt ccaggatgac 240  
 cagtcaggcc agtttcattg cctacgacat gctcgaccgg atccgcgcca attcgggtgc 300  
 tgattactcc tggggccagg gtgaacgcgc gccctccacc acctcggtcg cgagtgtgcg 360  
 tgatctggac ctgcacgact ttgaagcgaa tatcgtcggg ttcgccgggg aaagcgccaa 420  
 ggggtccggt gcggtcaatc ancnagaagt gaccatcagc atcagttggg acnactcccg 480  
 tggagcgaaat gcncaaggca cccgggaaac attcacctcg accagccggg ttgcagtcga 540  
 tccgaggggtg ttgccatgag aggtccagtg tgcggtttca gcctggtgga gatgctgctg 600  
 gcattggccc tcggcctgat gttgatcctg ggggtgaccc aaattgcact cagctcccga 660  
 accacttatg ccagccagag tgcggcttcg ctgttgagg atgatgcacg gttcgccctt 720  
 ggcaagctga ttcaggaaat acgccaggcg ggcattgttg gctgcttgtc cgctgcatca 780  
 atcagcaacg ctcccgcagg ttttgatcgt cccattggat ggagtaccac cggcagttcc 840  
 cggctccctga cgctgggtgac cgccgacgtc ggggaggggtg gcagcaagcc ggactggacg 900  
 gtgctttccg attgcaccgg ctctgcccac gcctatgttg gaagcccgcg ggcagcgaaac 960  
 gcccgggcaa atccacttcc cacttgcgca aagctgacct aacacctttg nagggcgggc 1020

33

aagcnggaag	ttaagtaacg	ctggcggccc	ccgagcaaag	cggtggttgg	tggataacgt	1080
ggggcattcg	atatcagttt	tcggcgtggc	ngacaagcnt	ggctcaacgg	ttgtcagccg	1140
atatgacccc	accccccggc	atgagtcgnt	catccgcagc	gtgcggattc	tgctgacact	1200
tcaggatcca	aatgggttgg	tgaaagacca	ggcntacagc	gtggtcgcgg	cactacgtaa	1260
tcgcctggag	tagcgtgccc	atgggttatt	acctctcccg	ttcgaggcag	gcaggcatgg	1320
ttttgctgat	cagcctggta	ttcctgctgc	tgttggcact	cctcggagtg	tcttcgatgc	1380
agggagcaat	ctcgcaagaa	aaaattaccg	gcagccttcg	gcagcgcaac	cagtcgtttc	1440
agcaggccga	aagcggcctc	aggcttggcg	agtctttggt	gcaggcgta	ggtttcgccc	1500
tgcgccttg	ccactcgacg	gctgcgtggc	cgccacctgc	cgaatcgggt	tcggtagtgg	1560
ggcggggac	gaaccccgta	tcgactgtga	cctggatagg	gatgaaagat	ggcgtctacg	1620
gtattcaaaa	cctggggccg	ggaacgggtt	tggtcaactc	ccggcagagg	cccaggccac	1680
ggtctatcgc	gtgacatcag	tgggcgtcag	tgggcactcg	cgttcggtcc	tggagtctgt	1740
gtatgcccgt	gtgggcagcg	ggcccggcga	gcgtttccga	cgaatcatgt	ggcgacaact	1800
tcaataggtg	agcagcacga	tgggcaagga	ttgcacaggc	ttcacctga	tcgaattact	1860
gacgcctgtg	gcctcgtgcc	gaattcggca	cgagattaat	tccccaaaa	atctttgaaa	1920
tagggcccgt	atttacccta	tagcaccccc	tctacccccc	ctagagccaa	aaaaaaaaaa	1980
aaaaaaaa						1987

&lt;210&gt; 48

&lt;211&gt; 2113

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

ggggaatatg	gatggctttc	ttctttgccc	tctttgttat	tttctttggt	attgttggtc	60
agatggagtc	tcactccggc	ctgggcaaga	agagcaaat	tctgtctggg	gggcaggggg	120
aagaggatata	ttttttggat	taaaatggaa	tgctttcctc	aaactgttaa	aattaggtag	180
tgttacaagt	ttaaagaata	ggtttagcca	gtgtgattat	tgagtgtaa	taaaaggat	240
cccagggtggc	acacatgcta	gtgtggtgcc	cagatgctca	actgaagtga	aggataacat	300
gcgtttcaac	atcttctata	tactaggcag	ttttctttaa	gtcttggttt	attgtgatat	360
aatacagtaa	aatttactct	ttatacgtat	gattccatta	atattaacaa	atttaaacag	420
tcttacagct	aaatacaatc	aagatacaga	acatttccat	cactacagga	agttgttttt	480
tgctaattta	actaaacttc	ttaccatgct	caacccccca	ccccaccct	gagaaccact	540
gctttgtttt	tctgatcata	tagtttgccc	ttttcaaawa	tgtcatacag	ttggaatcat	600
atagtatgca	tgaatcatgt	agctgaagtt	ttcatttcac	ttaggtaaac	acctaggaat	660
aggagtgtcg	ggttatactc	taagtgttaa	acttcataag	aagctgctaa	actgttttcc	720
aaagcatctg	taccattttc	ctttcccacc	agcaataaag	cattcattag	tctacgtact	780
caccagtgtc	agtgtggtca	gaatgtatag	ttttaattat	acacattata	atagatgtag	840
agtggatatc	catcgtgatt	ttgccttttc	ctaataaata	tctttcctta	catatttgct	900
attagtgtgt	cttctttagt	aaatggctct	attgttttgc	ccatttttaa	aagttgagtt	960
ttcatattgk	tcaatttcga	gagttcttca	tataattctg	acacaagtc	tttgtcagac	1020
atgtgatttg	caaataatct	ccccagagt	ttttcctgtc	ttgtcagtc	cttaaatagt	1080
tttttaggaa	ggagaacagc	tgcttttaat	tttgataaag	cttgcaattg	tttttctttt	1140
atggatcatg	cttttggtgt	agtatctgag	aactctttct	taaaaccagtc	acacatgtct	1200
tctgttttct	ttgaagcttt	acaggttgag	gtacattttg	gtccatgac	acttttgagc	1260
tactttttat	atataggtag	tataagatat	ggtttgaggt	ttttgttttg	ggtttttttt	1320
cttcctgcat	atgaatgttc	agttgttcca	gcaccatttg	ttgaaaaaac	actatccttt	1380
ctctagtaat	ttgctttttt	acttttgcca	gttgactatt	tgtgggtctg	tttttaaaact	1440
ccattttata	ttatgtctac	attatgaact	ttatagttag	tcttgaaatt	mggtaattgtg	1500
ggtcttctca	cttgtgcttt	tcaaaattgt	tttggctctt	ctaattcttt	cactttttcc	1560
atataaattt	tagaaamagc	ttattgattt	ctaccacttt	cccccaaaaa	gccacttggg	1620
aatttgacta	agtttacatc	gaatctaata	aatgattttg	gagagaagtg	gtatcttagc	1680
aatacagctc	tttctgataa	cacttcctgt	tttagttctt	tttctcattt	aataattttc	1740
caacattaaa	accttggaac	aatttagatg	tgtatctgaa	taattccagt	tttggtgcca	1800
tttaaaatgg	tacataatcc	ttattttata	gatgagaaaa	ttaagtaact	tgcccaagtc	1860
acacagttac	taaatgacaa	agctagattg	aaatctatga	gcttataaac	taatgtcctg	1920
ttttgagcta	cagtactagt	tttattaaat	tgttggtggt	aaactgaggt	gggaggatag	1980
tttgagcacg	gaagattgag	gctgcagtga	tctttgatcc	tgccattgta	ctccagcctg	2040



34

gcaacagagt gagaccctgt ctccggaaaa aaaaaaaaaa caccatggaa gcaagcaaaa 2100  
 aaaaaaaaaa aag 2113

&lt;210&gt; 49

&lt;211&gt; 3465

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

agaaaaatgaa agcaccaaag agccttcttt gctacaatat ctttgtgtgc agtctcctgc 60  
 aggattaaat ggtttcaatg tacttttatc tggcagtcac accccccta ctgtgggccc 120  
 gtcctcaggt, cagctgccgt ctttcagtgt ccttgcagtg gtcttaccat ctccacctct 180  
 gggccctttt cctgttctct attctctctg aatgccgggc ccggtttctt ctactcttgg 240  
 tgctctccca aacacaggac ctgtgaattt cagcttgccg ggccttggat caatagccca 300  
 gcttctcgtc ggccccacag ctgtgggttaa tccaaagtgc tccacactcc cttctgcaga 360  
 cctcagctt cagagtcagc cctcactaaa cctaagtcga gtgatgtcaa ggtcacacag 420  
 tgtcgtccaa caacctgagt cccccgttta cgtgggacat ccagtctcag tagtaaaatt 480  
 acatcagtcac ccagttccag tgaccccaaa gagcatccaa cgcacacatc gtgagacgtt 540  
 tttcaagaca cccggcagcc ttggagaccc tgtcctgaag agaagagaaa ggaacaatca 600  
 cgaacacacca gtcgagccca gaggagacta gaaatcccca gcggcggcgc tgactaacct 660  
 gccgctttgc caggtggggg tgggatcaaa cgccctgaga gtcccggatg tccgaggcgg 720  
 gatgcaaaccc atcccgtcct gagcacgggt ccttcctctc tctttcatcc acacttctgt 780  
 taacttccca ccaccatcaa tcatctgatt tccctgaaagt aattaattgt gcatttaata 840  
 ccagttagag ttccgactct gcatgggtgc acagtgaag cgccgactga cttatggttt 900  
 tgattcaaga atcgtcttat tgcgtggaag agatctgaat aggataccgg agccttgttt 960  
 ttctaagggg gggcgtctgc tagcacttaa ctagggtgag cattcttaac atgtatttcc 1020  
 acttgccctg agtaaatctg tgggtgagaga agcttctctt ctgcagttta aaaaagctac 1080  
 tgcttctctta ggcttcatca ggaagccatc ttcagttgtg aatcctatgg tgttatttat 1140  
 tttgttctctg aaatgggatt tagtgcaaaa agtttacaac tacagtcttt aacacatttt 1200  
 tttcagggta tgacgacttg aatgtttata cttttattct ataatttgcc ctgcacttat 1260  
 tttacaacct agtaataatg tggataaatg tatctacatg acacatgtca agaccaaatt 1320  
 aactgtgaat gacacacctt gctgtaaatg aactgtgcta accctgactg tgggcttgag 1380  
 aacaaagatg aactctagaa ctctagcagc ctaactgctg cttctcaaat aactgtgtga 1440  
 acagtgagat attactgttt gtttctaaaa atcctactgt gccoagtttc cttcactaca 1500  
 tgccctgcat tttttattta aatatttagc tgtagcgcca tcagatatgg atgccttcta 1560  
 acaattgctg tttgtaaaaat aaatcaggat ggtagaaagt gattttatgg aaaattggaa 1620  
 cctggatgag accttttctg tgaattctga agagtaatga tgcgaaaatt gatacagggc 1680  
 aagagatgat tcttttgttt ttcttctact tcatgtccag aagagtaaga gggaaaatgg 1740  
 acatatgttt catatccaag ggtattcaaa ctgtagttag ttggtacctc tgaaaaatga 1800  
 gaatggtag cgcacggggt ggttgttcta gcatgaatc aattctggaa actgttatgc 1860  
 aatttccctt ttttaaccca cattacttta ggggtgcatt aagtcgcca actatactag 1920  
 ttctttgtat tcctagactt gctgatattt acctctctct tgtctcttca gagttaaagg 1980  
 ttccctctct ttcttctac ttcttctcat tctctcttcc ttccctctct cctacttctt 2040  
 ttcttctctc ctcttctct cttaaaaacta tcttagatgt agaactctgg ttaggggttt 2100  
 tattttattt ttattttttg acccaataaa atgttatatg aaagaatgaa aatattaatt 2160  
 taagagactc tgggagctg aataaagtag ctttatatta actacaggat aatattagcc 2220  
 ttattacccc cacaagattt tttaaaactt gaggtaggtg gctacattaa ataaatttgc 2280  
 tacttatata aaaattttta tcaacactaa acttttaaag tttacaagtt ttttttttct 2340  
 tttttacagt cttctataga gttaggttaa aaatgtggtt ctaaccatca acaattgcac 2400  
 ggttaaatga cctgaacta aaactgatgg gttccctatc aaaacaaata aaaatatacc 2460  
 tttttcaggt ttcaatctgt gcagggtata tgcagtgtta ttctaccatg ctttaagaact 2520  
 tccacaaaaat atttcatgga gaggtctgca tttagacgga aacagaaatt gcttttcccc 2580  
 tcaactgttc tgaatgctct atacttgttt taacattttt gctatctttt tttattattc 2640  
 tgatcatgat atgaccattt aacctcagaa ttcataatc ctgaggggtg ttaagaagca 2700  
 gtcccattgg tgaggatatt atgacttggg gaccattctt aggagtagaa aaccaaggac 2760  
 aattgcttct gtattcagta tccacttctt aatgtggctt tatatgtaaa aataataatg 2820  
 cagtgggtgt ttctgtcagg aaaataaatc ttacagaaca actggtggaa ttgaagctgc 2880  
 tgcgctagac ttggatattt tgggtagtga agaagcaatg gcaatcttga gtctattatt 2940

35

gtataattta	gtaaaagaaa	aaaataatcg	ttgggtgggtcc	tactaagaga	atgcagcttt	3000
tttgagttgt	cacagaggct	gtgtgtgccc	tacactgacc	agggtttgta	aaaccctttc	3060
attctggtag	aagagtcggg	ggtataactt	ttatacttga	atctacctac	caagtttaca	3120
tttctcaatt	cctttttgta	aggtgctatt	tctgtattta	aataactttc	ttttaacgta	3180
aagctgcttt	ctgcttatct	tattgcactg	ctagttgtat	gtaggattta	attttattgc	3240
tgcttactgc	ttttgttttc	ttattattta	gctctgctct	ttttccta	ggctatat	3300
tctatagcta	tttacttgta	actgtactac	atgtaaactg	attttttggt	ctgatttttt	3360
ttctaataatt	tttaggaaaa	tattaagctt	tataaaatag	caataaaaaa	taattcattt	3420
aaaaaaaaaa	aaaaaaaaact	cgtagggggg	gcccgtagcc	aattc		3465

&lt;210&gt; 50

&lt;211&gt; 1237

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

atggggccagg	aatgggggtcc	ccgggcatgg	tgctgggcct	cctgggtgcag	atctggggccc	60
tgcaagaagc	ctcaagcctg	agcgtgcagc	aggggcccac	cttgctgcag	gtgaggcagg	120
gcagtcaggc	gaccttggtc	tgccagggtg	accaggccac	agcctgggaa	cggctccgtg	180
ttaagtggac	aaaggatggg	gccatcctgt	gtcaaccgta	catcaccac	ggcagcctca	240
gcctgggggt	ctgcggggcc	cagggacggc	tctcctggca	ggcaccagc	catctcacc	300
tgtagcttga	ccctgtgagc	ctcaaccaca	gcggggcgta	cgtgtgctgg	gcggccgtag	360
agattcctga	gttgaggagg	gctgagggca	acataacaag	gctctttgtg	gaccagatg	420
acccacacac	gaacagaaac	cggatcgcaa	gcttcccagg	attcctcttc	gtgctgctgg	480
gggtgggaag	catgggtgtg	gctgcgatcg	tgtgggggtg	ctggttctgg	ggccgccgca	540
gctgccagca	aagggactca	ggtaacagcc	caggaaatgc	attctacagc	aacgtcctat	600
accggccccg	gggggcccc	aagaagagtg	aggactgctc	tggagagggg	aaggaccaga	660
ggggccagag	catttattca	acctccttcc	cgcaaccggc	cccccgccag	ccgcacctgg	720
cgtcaagacc	ctgccccagc	ccgagacct	gccccagccc	caggcccggc	caccccgctc	780
ctatgggtcag	ggtctctcct	agaccaagcc	ccaccagca	gccgaggcca	aaagggttcc	840
ccaaagtggg	agaggagtga	gagatcccag	gagacctcaa	caggacccca	cccataggta	900
cacacaaaaa	aggggggagc	gaggccagac	acggtggctc	acgcctgtaa	tcccagcagt	960
ttgggaagcc	gaggcgggtg	gaacacttga	ggtcaggggg	ttgagaccag	cctggcctga	1020
acctggggagg	cggagggttc	agttagccga	gattgcgcca	ctgcactcca	gcctggggcga	1080
cagagtgaga	ctccgtctca	aaaaaaaaaa	aaagcaggag	gattggggagc	ctgtcagccc	1140
catcctgaga	ccccgtctc	atctctgtaa	tgatggatct	cgctcccact	ttcccccaag	1200
aacctaataa	aggcttgtga	agaaaaaaaa	aaaaaaa			1237

&lt;210&gt; 51

&lt;211&gt; 1397

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1383)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1396)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 51

ggctccctagg	agttgagcag	gaacaggcat	ctgtgggtta	cggcgacctg	gctctccgcr	60
ggccacgtgg	gtgggtgagg	cacacgagtg	ggaagcggca	ccgacgtgtt	tctccccgac	120
cgtggccttg	ccaaagactt	ttaatagcat	tttttaagtg	caaaacgtct	aggtaaaaat	180

36

ctttatcatc	agtgacaaaa	ttagaatgta	tttaatatag	taggtggttt	aagaactggt	240
ttaacgtaag	acaaactgat	agcaacattc	tgttgtttta	aaggaggtgg	gtccgtgaca	300
ttctgcagct	agtcactac	tccaaggtaa	ctatcgactt	ggtttcagtg	aatctatttt	360
gttttttaact	acagtgattt	attagctcag	tatctagaaa	ttacgtatat	tttgtgctac	420
tgtcatcgat	gtgtaaactc	tgtttttatt	tgtatttatg	cacttggttc	ccatttggag	480
cctctggtct	tttctgggat	aagtgggtgc	tgccgagaca	tctcccggtt	gtcagtggtc	540
aggagcagct	gagctctagt	ctgccagctg	ctctgctctt	tctgggaagg	aggtggcgcc	600
cgccctcag	ggtgtctcca	gggtcagct	tccgggtgg	tagagctggg	gagccccagg	660
ggtgggggga	cagctgggag	atggaggtgg	cacctgctcc	cctagatcag	tactggctct	720
gaggacaggt	gagcagtggt	aagaccaaag	aatggctggc	agcgtgccca	rggttggaaa	780
tgggggcaag	atcctggggc	tgtgtgccct	ggggcctccc	tcacctgtct	tgggtggcat	840
ggctcaggg	atggctccta	ggtggctgag	gcacagcagt	ggctggaagg	tgccccgtgg	900
aggctgaggt	ggaggcgccg	ccagcagctc	ccccctgtgg	ccatggcggg	cacgggscgt	960
aggagctggc	tggcgccgg	ctctgcatgt	tcttgttgcc	tgtcgtctgt	aactctagt	1020
ttcgacattc	gccgtgatac	agtgggtgtca	cgacgtgtgt	aactgtggtc	agcagacctt	1080
gttccgcgtg	gacgcctcaa	gtggattaat	ttctggaagc	ctcaatctgt	atgtttgagt	1140
atttacatga	gaatgttatt	tgaatggaat	tttcttaacc	cagaaggtag	tatttataat	1200
catttacttg	tagcgaactg	tttaaagtta	acacttgttt	aaattttttt	acactatagc	1260
atttatgcaa	tgttttacag	aattcatgga	gttattttta	tcagtatggg	aattaattaa	1320
aaccttgaat	cttaaaaaaa	aaaaaaaaag	ggcgccgct	ctagaggatc	caagcttacg	1380
tangcgtgca	tgcgana					1397

&lt;210&gt; 52

&lt;211&gt; 2271

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

cggcagcagc	ggcagcagta	tggaggggct	ttggaccatg	aggaggaggc	cctgtcatcg	60
ggcagtggtc	aagaggcaga	agccatgtta	gatgagcctc	aggaacaagc	ggagggtctc	120
ctgactgtgt	acgtgatata	tgaacactcc	tcacttcttc	cccaggacat	gatgagctac	180
attggggcca	agaggacagc	agtgggtgcg	gggataatgc	accgggaggc	ctttaacata	240
attggccgcc	gcatagtcca	ggtggcccg	gccatgtctt	tgactgagga	tgtgcttgct	300
gctgctcttg	ctgaccacct	tccagaggac	aagtggagcg	ctgagaagag	gcggcctctc	360
aagtccagct	tgggctatga	gatcaccttc	agtttactca	accagacccc	caagtcccat	420
gatgtctact	gggacattga	gggggctgtc	cggcgctatg	tgcaaccttt	cctgaatgcc	480
ctcggtgccg	ctggcaactt	ctctgtggac	tctcagattc	tttactatgc	aatgttgggg	540
gtgaatcccc	gctttgactc	agcttctctc	agctactatt	tggacatgca	cagcctcccc	600
catgtcatca	acccagtggg	gtcccggctg	ggatccagtg	ctgcctcctt	gtaccctgtg	660
ctcaactttc	tactctacgt	gcctgagctt	gcacactcac	cgctgtacat	tcaggacaag	720
gatggcgctc	cagtggccac	caatgccttc	catagtcccc	gctggggtgg	cattatggta	780
tataatgttg	actccaaaac	ctataatgcc	tcagtgtctg	cagtggagag	cgagggtggac	840
atggtgcgag	tgatggaggt	gttctctggc	cagttgcggt	tgtcttttgg	gattgtctcag	900
ccccagctgc	ctccaaaatg	cctgctttca	gggcctacga	gtgaagggtc	aatgacctgg	960
gagctagacc	ggctgctctg	ggctcgggtc	gtggagaacc	tggccacagc	caccaccacc	1020
cttacctccc	tggcgagct	tctgggcaag	atcagcaaca	ttgtcattaa	ggacgacgtg	1080
gcatctgagg	tgtacaaggc	tgtagctgcc	gtccagaagt	cggcagaaga	gttggcgctc	1140
gggcacctgg	catctgcctt	tgtcgccagc	caggaagctg	tgacatcctc	tgagcttgcc	1200
ttctttgacc	cgctactcct	ccacctcctt	tatttccttg	atgaccagaa	gtttgccatc	1260
tacatccccc	tcttctctgc	tatggctgtg	ccatcctctc	tgtccctggt	caagatcttc	1320
ctggagaccc	gcaagtcctg	gagaaagcct	gagaagacag	actgagcagg	gcagaccttc	1380
cataggaagc	cttccctttc	ggccaagggt	ggcggtgtta	gattgtgagg	cagctacatg	1440
gggcctgccc	gaatgactta	aatatttgtc	tccagtctcc	actgttggct	ctccagcaac	1500
caaagtacaa	cactccaaga	tgggttcata	ttttcttctc	ttccatttca	cctggctcaa	1560
tcctcctcca	ccaccagggg	cctcaaaagg	cacatcatcc	gggtctcctt	atcttgtttg	1620
ataaggctgc	tgctgtctc	cctctgtggc	aaggactgtt	tgttcttttg	ccccatttct	1680
caacatagca	cacttgtgca	ctgagaggag	ggagcattat	gggaaagtcc	ctgccttcca	1740
cacctctctc	tagtccctgt	gggacagccc	tagccccctg	tgtcatgaag	gggccaggga	1800

37.

ttgggtcacct	gtgggacatt	ctccctcact	cccctccctc	ctagttgggt	ttgtctgtca	1860
gggtgcagtct	ggcgggagtc	caggaggcag	cagctcagga	catgggtgctg	tgtgtgtgtg	1920
tgtgtgtgtg	tgtgtgtgtg	tgtgtgtgtc	agaggttcca	gaaagtcca	gatttggaaat	1980
caaacagtcc	tgaattcaaa	tccttgtttt	tgcaattatt	gtctggagag	ctttggataa	2040
gggtattgaat	ctctctgagc	ctcagttttt	catttgttca	aatggcactg	atgatgtctc	2100
ccttacaaga	tggttgtgag	gagtaaatgt	gatcagcatg	taaagtgtct	ggcgtgtagt	2160
aggctcttaa	taaacactgg	ctgaatatga	attggaatga	tacaaaaaaa	aaaaaaaaaa	2220
actcgagggg	gggcccggta	cccaattcgc	cctatagtga	gtcgtattac	a	2271

<210> 53  
 <211> 2769  
 <212> DNA  
 <213> Homo sapiens

<400> 53

ctgccaccgc	cgctccccc	ctcccgtgc	cctcgggccc	ggctgggctg	agctgcgatg	60
ccctcggact	tcattctcatt	gtcagcgcg	gacctagacc	tggaaatcgc	caagtccttc	120
tactcgcgag	attctctgaa	gttacacca	tcacagaatt	ttcatagagc	tggactattg	180
gaaggtcagc	aaagtaccat	tttaaagcat	attgtgttag	tatgcaaagg	ctttctctga	240
aatggagttt	aaatgggaaa	gtatgaaact	ttgcaaactc	acacaatgta	gttttctcct	300
aaagagtctg	atccttcttt	tagagcagct	gaatgtttca	atgggttttg	ttgctgcgtt	360
tgatgtgctt	gttggctgtt	ctatctgctt	tgagaaacat	tgaaaaaaaa	atcattttaa	420
attatgtgtt	gagtgttcta	ctttagaatt	ccagaccttc	attttagctt	tttatacaca	480
aaatagaatg	tgaaaaggcc	caaacattaa	catgaatata	aaagagtgc	accaaacaca	540
taattagttt	gaaaaattcag	aatttagaac	atttatatgc	tgagcatttc	aaaacttgtt	600
aattatttat	tgcaaaagca	ttgatgccat	tgttattgca	ttataactta	gggtaaggaa	660
agccaagtaa	tagcgatgat	gattgagcta	tgcttagaag	aaaggggttt	agaaagttct	720
ggaaatgtta	atagagtctg	tagattagtt	taaacagctt	acacttactt	tagaaaactt	780
ttggctttgt	gtgtgtgtgt	gttttaacat	tttaagacat	aaaatagtct	tgtgtttttg	840
tagattttga	agtgatttca	catatatcat	ctttaaatct	ttataatttt	ggaattggac	900
agcaaaggta	ctaatttccc	cattgtagag	aagagaaact	aaggctgaaa	ttaagaaacc	960
aaaaaaaaaa	aaaagtccca	cagtagaaga	aataataact	tctgcgtttc	agtttagtgc	1020
ctttatgcaa	agcggccacc	gtgaaataca	catgtaacca	gtaacaaaca	ttttatggtc	1080
tactttaaaa	aggccctaaa	tatttaagtt	tgacttgaga	agatttttgc	ctcccttgga	1140
ggtagtccgg	gatagtgggg	ggcaagtggg	ttttgtagtt	gagcttaaat	ttggatacca	1200
acttagctcc	ttattagttc	tgtgacttcg	ggcaagtgat	tttacctctc	taaactttaa	1260
tattcttctt	gttaaaatgt	agctgtctct	taaatttagt	gagctaattg	gtattaaatt	1320
gcctagcata	gcgtctagct	taataaatgt	tagttttttc	ctttagtatc	agctaaattt	1380
ttttaacagt	ataatttaga	gcatataaaa	tgttctcaaa	taaataattg	gttaaatattg	1440
ttttgtacgt	gtaagaagcc	catagatata	cattagtatg	aaaatcaaat	atttttaaaa	1500
ctacaatttg	aagtgagaaa	tttttagcag	tgtttcttag	tgaagcagat	ttctggtttt	1560
tattcatatt	caatttgtga	ccatgttgta	gagacatatt	tgtctgtaca	aacctatctc	1620
atttcctgga	caacaaacca	tagagtgtca	taatcataga	gatgatcagg	gaggagaaaag	1680
acatgaagaa	aaaaatgagt	gttaattatt	taagcaaata	tttttagtga	gctgatggaa	1740
atacagacaa	ccatttagca	tgtttattta	catgtttatg	cctaaatact	ttcgtaatt	1800
tttcatttta	gttaagggga	ttctcattta	aagtaaggct	acccttttag	aaattatata	1860
taaaatataa	catgtataag	tatcaattct	gtcatgccct	ttaaaatggt	ttgagacagg	1920
gtgtcttaaa	aaagtgtgtt	gcttctgtaa	tgaagtgtct	aatctgtgcc	acctcattgc	1980
tatttctttt	ttattattga	cattttaaag	aatacataaa	agtagagaga	ataaaaataag	2040
ccccatatac	ccatcaccac	agtttaacca	gtatctaate	ttgcttcttt	tatcccttct	2100
tctctctcaa	gctggattat	tttaaaacaa	atatcaaaaa	tcagatcact	tcatacataa	2160
gtatttctgt	attacagtga	ctttcttata	ttaattattg	atataattta	attgttaata	2220
taattattgt	attaattttt	aattactctg	tatagtttat	aaactatttc	ttggaatatt	2280
ctagtttatt	tcagcaaagt	ctcttgtagc	tctcctgccc	tcttttcaag	acagagtgc	2340
gcaggaggct	accgcttcta	gtatatagta	tatcctataa	cctatgttga	aattgtaggc	2400
catgctgcta	gattggagtt	catattgttg	ctctgacatt	tactgtatg	gccttagcct	2460
aataatttaa	tcttttttgc	ttcaattttc	tcactcttaa	acgggagcaa	taatagtgc	2520
tcactcctagg	tttttgtgaa	gattatttga	ggcaaatctg	tgtaaagcat	ttaattttagc	2580

38

acatgctaag	tgctcagtaa	atgttagcta	aagttaaaaa	atttttaaaa	cctaaatgct	2640
agctgtagag	tgtttactgt	gtgccaagaa	ctttacaaa	aagaactcct	gaggtgggca	2700
tatcacttgc	ggtcaggagt	ttgagaccag	cctggcttac	atggtgaaac	cccactctcta	2760
ctaaaaaaa						2769

<210> 54  
 <211> 1389  
 <212> DNA  
 <213> Homo sapiens

<400> 54	
ctgcaggatt	60
cgccacgagg	
atctgttgac	
cacagatgta	
agaattttatt	
tctggagcct	
caatttcta	120
ccattgctct	
acatgtgcac	
tggtttcctg	
gggctgccac	
agcagaatac	
cacagacttg	180
ggggcttaaa	
caacaagtg	
attcctcaca	
gttctggagg	
ctccaagtct	
gagatcaagg	240
agtcggcatg	
tttggttcct	
cctgaggcct	
ctctccttgg	
cctgcagatg	
accacgttct	300
cactgtgttc	
tcagctggcc	
cttctctgtg	
catgtacatc	
cctgggtgtct	
cttctcctct	360
ttgttgacta	
taaggatacc	
agtcctgttg	
gaccagagcc	
ccactgtaaa	
ggcctcattt	420
taacttaatc	
cacttaaata	
acttatttcc	
acacacagta	
agtctgaggt	
tgagtctttt	480
gaattctggg	
ggagacactg	
cagcccatca	
cgtatgttga	
cgtgtgccac	
cgccacactg	540
cctggttact	
gcagagtgtt	
agtaagtttt	
gaaattagaa	
agtataagtg	
ctccagttct	600
gttatttttc	
aggattatta	
gggctattct	
gggtcccttc	
catctccata	
tgaatttgag	660
aatcagcttg	
tttattcctg	
ctaaaattta	
gttgggattt	
tttttttttt	
gagatggagc	720
ctggctgtgt	
caccaggctg	
gtgtgcagtg	
gtgcaatctc	
gcctcactgc	
aacctccacc	780
tcccgggttc	
aagtgattct	
cctgcctcag	
cctcctgagt	
agctgggact	
acatgcgcac	840
gccaccacac	
tcagctaatt	
tttgattttt	
tagtagagac	
gggggtttcac	
catgttgccc	900
aggatgggtct	
tgatctcttg	
acctcgtgat	
ccgcccacct	
tggcctccca	
aagtgtctgg	960
attacaggca	
tgagccaccg	
cgcttgccct	
cagttgggat	
tttcacaggg	
ctcgttttga	1020
gtctgcagat	
caatttggtt	
agtactgaca	
tcttaaaatt	
aagtctaccg	
atccacataa	1080
gtgggatgct	
agttaatatg	
gacaagtga	
tttattactc	
ctccaaaaaa	
ctattttcac	1140
aaactttata	
tcaacatgag	
aaatcatgtt	
gtatctttct	
acctggaggg	
agcaagaaag	1200
tcaggtgggtg	
tgctttttgt	
acatgggcaa	
cccaatttct	
tatctagact	
ttcttttgct	1260
cgctgggaaa	
aaaaaaaaaa	
tgaggtattg	
caaaaatgtg	
ataattgctt	
tccttagtat	1320
gaaaaactgt	
taaaatgagt	
aaagcctgtg	
ctcaactaga	
attctcacat	
ttccgttttg	1380
atgtatccgc	
atttcacctc	
gtgccgaatt	
cgatatcaag	
cttatcgata	
ccgtcgcacc	1389

<210> 55  
 <211> 748  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (15)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (17)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (29)  
 <223> n equals a,t,g, or c

<220>

39

<221> SITE  
 <222> (32)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (646)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (741)  
 <223> n equals a,t,g, or c

<400> 55  
 ggtgcggcgcg tttanantag ggatccccng gntgcaggaa ttcggcacga gcttattgka 60  
 tgtgctmcag tgacaggcat ctgggacttt tccaatctga agctgttagg aacaccacca 120  
 ccatgaatat tcttgatgt gttttctggc tctggggtgg agttgctggg, tcatggggca 180  
 ggcacatttt catttttact tcagtaaaaa atgtcyagkg ggccagccac tgtgcctggc 240  
 ccaratgact tcttcacaag gagacacata ctgtgtgtat cagtcaggat ccaaccagga 300  
 gacaaaccac acagtaattt aaacagcgat tgtttaatat acagaattgt taactatgat 360  
 aggggatttg agtaagagga actggttact aagaaataaa gagaatgcta acgaatgtag 420  
 aaatagactg ggcacaatgg ctcacacctg taatcccagc actttgggag gcaaggcagg 480  
 tggatcacga ggtcaagaga tcgagaccat cctggccaac atggtgaaac cccgtcccta 540  
 ctacaaaaag tagctgggcg tgggtggcgtg tgcctgtagt ccagctact caaggaggct 600  
 gaggcaggag aatagtttga acccaggagg cagaggttgc ggtganccga gatcttgcca 660  
 ctgcactcca gcctgggtga cagagagaga ctccatctca aaaaaaaaaa aaaaaaaaaa 720  
 actcgagggg gggcccggtg nccaattc 748

<210> 56  
 <211> 4202  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (57)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (58)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (61)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (4137)  
 <223> n equals a,t,g, or c

<400> 56  
 accagctcaa agattcatga gtttcacga gtcactgtga gtggagccca tgctgggntg 60  
 ntgcctctg tgtctgtgca tgcgctgtg tgtgtggcg tgtgtgcatt gctgggccag 120  
 cttgaaggga agggccgtca tgccctgca ctctgttttg caagatgccca aacccagtt 180

ctgatggggc	tccaacagcc	aggctgtggt	cctttgacgt	tcctcacctg	ttgccaacct	240
atcccgtagt	gaactgaaac	cccagtgaag	acagaactgt	gcctggggag	atgcaatgag	300
gtgagggctg	aactcaccct	tttatatttc	ttttcaagat	tggatcagag	ctcatctcca	360
tccagtcctg	tttctatgaa	ggcttcaatc	tgtttccatg	caaatttgct	aatcagagcc	420
cagagctgct	gggtccctca	tctccctcat	ctattataga	ttgacttaca	gcagggagag	480
aatctcttta	gtcattcctt	aatgggggtg	ggatcacaa	atggtctggt	ccaatctgca	540
tcttggtgtg	tccaagacc	ctatctcctc	cccaacattc	ttattgcctt	tggctcccag	600
taaggaacga	attggggggc	agggaggaga	acagggggga	tcaagaaggg	aaacccaatt	660
ccccctttga	aagtgggttc	tttgaactat	gtgtttgggg	gaagttcctc	tggatactaa	720
tttgaattta	tatacctcat	gttttggggg	tttgacgtat	atatatatat	atatatatat	780
atgcatatat	atttcataat	atttggaagg	tttttgatgc	tagaaaaatg	gaaacaagag	840
aaccttcaaa	aatggtactt	agatgggaac	tggaggccaa	tctttcataa	agccagcccc	900
atagctgctt	gctggttagc	ctccagccat	tttgacattg	gggtggatag	tcgattcacc	960
tgctgttcag	tcgattcacc	tgctgttcac	ccagttctgt	ggatgtgctg	gtgctgagcc	1020
tttgcctctc	ttccaaatgg	ttacagggat	gttgatcagc	tccaccagag	ggagctctga	1080
tgggaggaat	tgctctgcca	tccttgtccc	tgtgtctcct	gtcggcaggc	agccattgta	1140
tctcaccagc	agaccaggag	actggtccca	aggttactgc	accacagggc	aatttcctgc	1200
catagtttag	aaggaaacac	ctgaactaaa	tggaaagagc	atccctgcgg	tgtttaatat	1260
cacacccatg	ccctttgtca	ggttaccatg	tacagagatt	acttggagag	cctcatgccg	1320
tctctacctt	cgcacactgg	tcaagtatct	gctgagcttc	ttggccgcaa	ggatgcagaa	1380
ataggctgag	ggtccatggg	aagaaagaca	caatgaggca	gtaggagggtg	gggaagaaaa	1440
gaagacagac	tttcaaaatg	gaattaggca	ctggggagag	atcagtttcc	ccacatcagg	1500
gagaagaagg	tataggtggg	gaaggggggtg	gccaggagca	gaaggaagaa	gactcaagat	1560
ggaaagggag	ccgctgtgcc	tgtggcaata	ccagttggag	aggtcgactt	cataccttca	1620
agccttttcc	cctgggcttt	tgattgtgtc	tgtgccccct	ttcttgtcct	ctctgcagat	1680
gcccagtagg	ggctacctca	tcctcgtgct	gttcttgtgt	ggctttcttg	gcagtaggga	1740
tcttgaattt	cctttctaac	actgtgcccg	gcaaggcggg	gagcattcct	ctgccctttg	1800
tcttgtgcca	acctggaaag	gtgcagtcta	gatttcagtg	agaacctgc	cagctgagcc	1860
ctgtgcatct	actaccttga	cacagagtgt	tttccacta	gaagctctgc	tctgctctcc	1920
tggcccaagt	aggggattcc	atgccttccc	tttcatggtc	ttagcaccag	cagcctagtt	1980
tctcccttcc	agagtctcca	gggatgacaa	attggattgg	agacaaacct	cgtcagatgc	2040
tcattcccta	aaaggttaat	tgtgtatttg	tggctgcgtg	tgctttgtg	ttttcattct	2100
cttccctatt	ttgtacattt	tggtctcttc	tgtggtttta	tacttgggtca	aaagtactcg	2160
tcttgggtatt	gcactgttgt	gtgcatgaga	aaactggggg	aaggctcact	ggtacaagaa	2220
aggacccctg	acccctttcc	ttctctgtgg	ttcccggcat	tagattgggg	gttctgggag	2280
aggcaggtga	atgtcctaag	tgaattgttc	tgtttgaac	tggaaatgtt	ttgaagtctt	2340
tgggtgtgct	ccgtgaaagg	acatcgccac	ctggtgtctc	tgaggtgtct	ttgcagaaca	2400
ataaatggca	aatgaacaac	camaaaatg	ttacycttgt	tggccttctg	ctgttttag	2460
attagtgcac	ctatctgtga	gggatttggg	ttacctccct	gagtctgtaa	gcaaccacaa	2520
gccctgccac	tgggtggggg	aagtcctctc	ccaaccactt	aaaaacaaat	tttcccacat	2580
attaccccac	cccacacatt	tgacctggc	tagactttgt	ttgcctaaag	gaacagacca	2640
cattgctggg	aaaatgagta	agtgaacgtg	tgggagaaaa	acacttttag	aatcacgaat	2700
attcactttt	aaaggctctc	ttgcctggct	gcaatatagt	gtgtgtttta	attatttaca	2760
ggctgttgtt	tctcaataaa	atgtttaata	ttaatcattc	ccaaactgac	aagaacacaa	2820
aaataaaaatg	caaatacaga	gccagctttg	tcacccaaat	ctgtgtctat	ttctgatatg	2880
ccatggaatg	tggttttctt	ggaagccagg	gttgggtctc	ccacagaccc	caggctaagg	2940
tcaccagtta	ggaaccaggg	acttgggaag	cagagctgtg	agctcttcca	tcagggtatc	3000
gactccgcaa	aacgacttga	tgaatgcaat	tggcaaacct	ccatgttcgg	acttcatatg	3060
catgagccgt	tggacagagg	gtttcttagt	atatacttta	atgcatgttt	atgtgcaatc	3120
ttgttagtg	gtatacaagt	ttgtgaagaa	cttctcattt	caataggcag	ttaatgtaat	3180
gcattaaaag	cctgggaatt	tggggctata	tttttccctt	ctgactcaat	aatcttcaaa	3240
gaattcatag	gaaagtcagt	acttgcagac	aagtggtag	cttggctaaa	atgtacaaaa	3300
cacccagaac	ccacaaaaca	ctcagagggt	taggagaatg	ttttaatgct	taagaggcag	3360
gatcaagtga	agaggttaca	gaaatcagtg	tctctggctg	ggcagtcagg	agagcgggct	3420
caaattctgt	gactcacttc	tctgtgtctc	ggttggaaat	gaatgggtat	cctgggtccc	3480
accttcccac	acgctgtgat	acttcaaact	ccttgggtga	agggcctctt	ctcagcccaa	3540
gatcttgatt	gtgaacatta	acaaagagaa	cagtcacact	ccacagaaga	taactcatta	3600
atgacatttg	attcagtga	taaatatatc	atttaaaaaa	atattgtagg	gggatcatga	3660
aagtagtgga	ggtaattaca	atcaggagag	attgggtatta	aaattgagca	aagtcaccaac	3720

41

tctcaccaga	tgacaattat	gcatacctgct	agatgccccca	gggctgtcag	cctggaactg	3780
aaataaatgt	gttataagtg	gtgctggatg	ccttttttcag	ttcatttgaa	aacatggatt	3840
tgatcatgtc	agctcccttt	ctgctggaaa	aaaaagtagt	ttgcataatt	gggtttaact	3900
actctgtttt	gattctacag	agtaagtaat	actcaaagt	ggctcttactt	taacttcttg	3960
cctttgttac	ccccagaacc	atgcagacat	tgaatgtgg	tgtgcgtgtg	tgtgtgtgtg	4020
tgtgtgtgtg	tgtgtgttca	tataacagga	ggacagga	ggtaaggacc	cagaacaatg	4080
aagacttatt	gaaatgtggg	gggggtgtgt	gtgtgtgtgt	gttcatagaa	caggagnaca	4140
ggaaaggtaa	tgaccatcca	tggaagatga	aggggtagta	cttagccgga	cgcggtgggtc	4200
ga						4202

<210> 57  
 <211> 854  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (831)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (839)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (844)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (851)  
 <223> n equals a,t,g, or c

<400> 57	
cgaaattact	cctcactaaa gttgaacaaa agctggagct cgcgcgctg caggtcgaca 60
ctagtggatc	caaagaattc ggcacgagcg ggatccccgc aaaatgcaga ttccccaaaa 120
tttttgtaaa	tacagatgac acttatgaag agctccattt aatcgtttat aaggtaacaa 180
ctgttttctt	tccagcgtta atgattgtgc tgaagtggat ctttcttgca tgtgtacacg 240
agtgcattgt	caaacctctt aaatgtttct tggaaaagat attggaagtt ctgattatgg 300
taaaactcaa	aatgggtgtt cttccagcgt aataagttta ttttcagctc cttttaaaca 360
gttctgttat	tagtgaaaga ggaactgttt aagattgtga tttataaacg tgtgaagtct 420
cacgtgctct	caaaccacag gctgtcagag gttggtgctg cctgttctcg aaatggctct 480
ggatgggggc	cgtagccacg tgtctgtgca tatgctgctt ttgctctgat tttaaagctg 540
taggcttgct	aattccataa ggatcgtatt ttgkttctgt caggacatgg tcttgtaagg 600
atatgtactc	aggttgtgtt tctaattaaa ggcagttttt gattcaagaa agaagacgga 660
gcatgtgcac	gtgtttctcc tctttctgct ctgaggctgt ggagaagttt tcatttataa 720
aggctcagaa	atgatgccgt gggggamcag gaaggagcgg agaactagtc tcgagagtac 780
ttctagagcg	gccgcggggc catcgatttt ccacccgggt ggggtaccag ntaagtgtng 840
aagnattccc	ntta 854

<210> 58  
 <211> 1455  
 <212> DNA  
 <213> Homo sapiens



42

<220>  
<221> SITE  
<222> (5)  
<223> n equals a,t,g, or c

<220>  
<221> SITE  
<222> (39)  
<223> n equals a,t,g, or c

<220>  
<221> SITE  
<222> (40)  
<223> n equals a,t,g, or c

<400> 58  
ggcanagctg ggatgggggtc agcaccacaga agccagccnn ctctgacagc ttcctctttg 60  
gccaaagccct gcctctgtac agcctcgagt ggacagccag agggcacgag ggagcccaga 120  
gcccagatg gagccccagc tggggcctga ggctgccgcc ctccgccctg gctggctggc 180  
cctgtgtctg tgggtctcag ccctgagctg ttctttctcc ttgccagctt cttccctttc 240  
ttctctgggtg ccccaagtca gaaccagcta caattttgga aggactttcc tcgggtcttga 300  
taaagtcaat gcctgcatcg ggacatctat ttgcaagaag ttcttttaag aagaaataag 360  
atctgacaac tggctggctt cccaccttgg actgcctccc gattccttgc tttcttatcc 420  
tgcaaatatc tcagatgatt ccaaaatctg gcgccctgtg gagaçcttta gactggctcag 480  
caaatatcaa aacgagatct cagacaggaa aatctgtgcc tctgcatcag ccccaaagac 540  
ctgcagcatt gagcgtgtcc tgcggaaaac agagaggttc cagaaatggc tgcaggccaa 600  
gcgccctcacg ccggacctgg tgcaggactg tcaccagggc cagagagaac taaagtccct 660  
gtgtatgctg agataacacc agtgaaaaag cctggcatgg agcccagcac tgagaacttc 720  
cagaaagtgt tagccttctc ccaactgtgt tataccaacc acattttcaa atagtaatca 780  
ttaaagaggc ttctgcatca aaccttcaca tgcagctccc atgccacctc cagaattcac 840  
caacacacag gccaccagc aacagggtacc ttgtcacaat attttttgat gacaatccaa 900  
agccccggct ctttcccacc acactgtggt cccctagatg gggctgttgc tgagcccacc 960  
ccaatcccag atgtgatccc ccctgtgatc tacttccctg caagattcct ccagtcctgg 1020  
acagggtctt cctatgagat agaacctgat aaggagctag ggcaattctg acaacattac 1080  
caaagggcca cataacttct aaattttggt ctggtctgaa ggaaaacctg ttcttgccct 1140  
agtgtaggat gaactctctt atctctggct tctagaggga aaaaaaagca tacctctttt 1200  
actttttaag tacctccatc agagtcatga aatcacctgt caagactatc tatcttttat 1260  
gtttccattc tggttaagaac tctttaaatg aggacactgc tgattgctgg tgatgttttt 1320  
tgagcaaaca ctcggggggtg tggatgaaag ccaatcgcat gtcaaatgac tccttgggga 1380  
agctacttct cctctattca gatttcaact aaatcttcca agatgaaagc aaaaaaaaaa 1440  
aaaaaaaaaa aaaaaa 1455

<210> 59  
<211> 593  
<212> DNA  
<213> Homo sapiens

<400> 59  
ccacgcttcc ggagctctta cttctccagc aacgctcttc agtacataat aagcttaact 60  
gataaacaga atatttagaa aggtgagact tgggcttacc attgggttta aatcataggg 120  
acctagggcg agggttcagg gcttctctgg agcagatatt gtcaagttca tggccttagg 180  
tagcatgtat ctggtcttaa ctctgattgt agcaaaagt ctgagaggag ctgagccctg 240  
ttgtggccca ttaaagaaca gggctcctcag gccctgcccg cttcctgtcc actgccccct 300  
ccccatcccc agcccagccg agggaatccc gtgggttgc taccctaccta taagggtggt 360  
tataagctgc tgtcctggcc actgcattca aattccaatg tgtacttcat agtgtaaaaa 420  
tttatattat tgtgagggtt tttgtctttt tttttttttt tttttttttg gtatattgct 480  
gtatctactt taacttccag aaataaacgt tatataggaa ccgacaaaaa aaaaaaaaaa 540  
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 593

<210> 60  
 <211> 496  
 <212> DNA  
 <213> Homo sapiens

<400> 60  
 aattcggcctt tcgagcggcc gcccgggcag gtcctgcttt ttgcttataa ttgacaacat 60  
 gtgcaaaaat accaaatttg tgcctgtgc agtatgaaga attcagtga tttcatttaa 120  
 tgtattagct tgttttgctc tctgttcata tatggctcta ttcttagaaa tataatttga 180  
 atgtgatctt tcaatagtct gaatatttta caaattatag ctatgtcttg tgaaaataac 240  
 ctcaaaaaga aaaatacgac tctgttgctc tacttgatat ttcttgccct agtaatgtac 300  
 ttgacattta tgttcctaag cagtgttaagt accagtagaa ttctctgtc aaactcaatg 360  
 atcatttagt acttttgtct tctcccatgt gcttgaagga aaaataaagt gtcactaccg 420  
 tatttcttgt tttcatcaaa aaataaaaat aatttaaaaa aaaaaaaaaa aaaaaaaaaa 480  
 aarrggsgg ccccc 496

<210> 61  
 <211> 1292  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (71)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (697)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1280)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1287)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1291)  
 <223> n equals a,t,g, or c

<400> 61  
 aaacctcttc tataggtaaa gctgggtacgc ctgcagggtac cgggtccggaa ttcccggtc 60  
 gacccacgcg nccggaaaga ggaacatag aggtgccaaa ggaacaaaga cataatgatg 120  
 tcatccaagc caacaagcca tgctgaagta aatgaaacca taccacaacc ttaccacca 180  
 agcagcttta tggctcctgg atttcaacag cctctgggtt caatcaactt agaaaaccaa 240  
 gctcaggggtg ctcagcgtgc tcagccctac ggcacacat ctccgggaat ctttgctagc 300  
 agtcaaccgg gtcaaggaaa tatacaaatg ataaatccaa gtgtgggaac agcagtaatg 360  
 aactttaaag aagaagcaaa ggcactaggg gtgatccaga tcatgggttg attgatgcac 420  
 attgggtttg gaattgtttt gtgtttaata tccttctctt ttagagaagt attaggtttt 480  
 gcctctactg ctgktattgg tggataccca ttctggggtg gcctttcttt tattatctct 540

44

ggctctctct	ctgtgtcagc	atccaaggag	ctttcccgtt	gtctggtgaa	aggcagcctg	600
ggaatgaaca	ttggtaggtc	tatcttggcc	ttcattggag	tgattctgct	gctgggtggat	660
atgtgcatca	atggggtarc	tggccaagac	tactggnccg	tgctttctgg	aaaaggcatt	720
tcagccacgc	tgatgatctt	ctccytcttg	gagttcttcg	tagcttggtc	cacagcccat	780
tttgccaacc	aagcaaacac	cacaaccaat	atgtctgtcc	tggttattcc	aaatatgtat	840
gaaagcaacc	ctgkgacacc	agcgtcttct	tcagctcctc	ccagatgcaa	caactactca	900
gctaattgcc	ctaaaagaaa	aaggggtatc	agtctaattc	catggagaaa	aactacttgc	960
aaaaacttct	taagaagatg	tcttttattg	tctacaatga	tttctagtct	ttaaaaactg	1020
tgtttgagat	ttgttttttag	gttggtcgct	aatgatggcg	gtatctccct	tcactgtctc	1080
ttcctacatt	accactacta	catgctggca	'aagggtgaagg'	atcagaggac	tgaaaaatga	1140
ttctgcaact	ctcttaaagt	tagaaaatgtt	tctgttcata	ttactttttc	cttaataaaa	1200
tgtcattaga	aacaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaagggc	ggccgctcta	1260
gaggatccaq	gcttacgtan	gcgtgcntgc	na			1292

<210> 62  
 <211> 398  
 <212> DNA  
 <213> Homo sapiens

<400> 62	
ccccctggc	ccccccatta atgccatggt ttgggaggag cckgatcccc tgaacccpat 60
atttaacctc	tactgcccms ggaaatgcc tacattattt ttccctaatt ggaagtataa 120
ttagagtgat	gttggtaggg tagaaaaaga gggagtcact tgatgctttc aggttaatca 180
gagctatggg	tgctacaggc ttgtctttct aagtgcata ttcttatcta attctcagat 240
cagggttttg	aagmtwtggg ggtcttttta gattttaatc cctactttct ttatggtaca 300
aatatgtaca	aaagaaaaag gtcttatatt cttttacaca aatttataaa taaattttga 360
actccttctg	tttaaaaaaa aaaaaaaaaa aaaaaaaa 398

<210> 63  
 <211> 1202  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (282)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (596)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (607)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (1200)  
 <223> n equals a,t,g, or c

<400> 63	
gaattcggca	cgagattaag ttgtgcactt taattgggtg aattgtacat gtragttata 60
tatctractg	tagttgtwat taaaaaacia caggaggcca tgtgggctgc taggagtagc 120
aatgtctgty	cccagccagc aggtagagac cagggctgga cagagkagta tgggctgtgc 180

45

tgcagattat	ttgtggtacc	caactgttgc	ataaaacagg	gtgtgatctc	ttgcattgct	240
atgcatgagt	ggattcccag	taaattgtgc	caggctgcct	gnatgatgtg	tggcttgtgc	300
tttggatcgt	aatgcttacc	tatgctactt	aagttacata	ccctgtggcc	tttgtggcca	360
ggactgtggg	ctactacctg	kagtgattcg	ttaggggaaa	ggaccacag	cctgtgcagg	420
aggaaaaaag	catctctgag	tacagggtgg	atgagctgga	tgagctgccg	ggcaagagcc	480
acgcacaccc	aggtggtgag	tcttaaggat	aaggtggaat	ttgccccata	gctgtcctgg	540
acagaaactg	cccagagaag	aatgaatgga	ggacataggg	ctctgtggtc	ccaccntttt	600
ttggganacc	tgtgactggg	cctgtttacca	tgtcaactta	gccccaaacc	catctctgat	660
tgacttgggt	gcttattttg	gcacattctt	gctccacaca	gccacataca	tactggctgc	720
tccctsaagg	ccaggcagat	gcagcagctg	ttgggpcasc	aaagargaar	gtcctggaar	780
gttctggcct	gaacgctgca	tctgttgtgt	gacagccaca	actgctcagg	cttccttgtc	840
tgtgggtgca	ctgtggggag	gagtgttatg	ataagaacat	tggctctcag	tcttccctgg	900
ggagaagttt	ggcctcacgt	gggatttggg	cgttgccctt	aggaaggctc	tctgcatgtc	960
tagttccagt	ttgtactggg	aagaattaaa	aaagtctgpc	agcttcttta	gtttgtcctg	1020
tcttttggga	tgattctttc	tgagatcccc	tcctatcagc	tcaggagtgg	gattttctgg	1080
agaaggaaaag	tgtttttcc	gttcctcact	gctcaccttg	gggcattcag	gaacatgggc	1140
ctgatgaatt	tgcttgaagg	cagtctgtaa	tcccatcact	ttgggagcca	aagaggcggn	1200
ca						1202

&lt;210&gt; 64

&lt;211&gt; 1517

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 64

gattacgccca	actcgaattt	aaccctcact	aaagggaaaca	aaagctggag	ctccaccgcg	60
gtggcgcccg	ctctagaact	agtggatccc	cgggctgca	ggaattcggc	acgagggagc	120
ccagagccca	agatggagcc	ccagctgggg	cctgaggctg	ccgccctccg	ccctggctgg	180
ctggccctgc	tgctgtgggt	ctcagccctg	agctgttctt	tctccttgcc	agcttcttcc	240
ctttcttctc	tggtgccccca	agtcagaacc	agctacaatt	ttggaaggac	tttcctcggt	300
cttgataaat	gcaatgcctg	catcgggaca	tctattttgca	agaagttctt	taaagaaaga	360
aataagatct	gacaactggc	tggcttccca	ccttgggact	gcctcccgat	tcccttgctt	420
tcttatcctg	gcaaattact	caggatgatt	ccaaaatctg	gcgccctgtg	gagatcttta	480
gactgttcag	caaatatcaa	aacgagatct	cagacaggaa	aatctgtgcc	tctgcatcag	540
ccccaagac	ctgcagcatt	gagcgtgtcc	tgcggaatac	agagaggttc	cagaaatggc	600
tgcaggccaa	gcgcctcacg	ccggacctgg	tgcaggactg	tcaccagggc	cagagagaac	660
taaagttcct	gtgtatgctg	gagataacac	cagtgaataa	gccttggcat	ggagccccag	720
cactgagaac	ttccagaaag	tgttagcctt	ctcccaactg	tgttatacca	accacatttt	780
caaatagtaa	tcattaaaga	ggcttctgca	tcaaaccttc	acatgcagct	cccatgccac	840
cctccagaat	tcaccaacac	acaggcccac	cagcaacagg	cttacctttt	gcacaatatt	900
ttttgatgac	aatccaaagc	cccggctctt	tcccaccaca	ctgtggtccc	ctagatgggg	960
ctgttgctga	gcccacccca	atcccagatg	tgatccccct	gtgatctact	tctggggcaa	1020
gattctccag	tctggacagg	tcttccccta	tgagatagaa	cctgataagg	agctagggca	1080
attctgacaa	cattaccaaa	ggcccacata	acttctaaat	tttggctctg	tctgaaggaa	1140
aacctgttct	tgccctagtg	atggatgaac	tctcttatct	ctggcttcta	gagggaaaaa	1200
aaagcatacc	tcttttactt	tttaagtacc	tccatcagag	tcatgaaatc	acctgtcaag	1260
actatctatc	ttttatgttt	ccattctggt	aagaactctt	taaatgagga	cactgctgat	1320
tgctggtgat	gttttttgag	caaacactcg	ggggtatgga	tgaaagccaa	tcgcagggtca	1380
aatgactcct	tggggaagct	acttctcctc	tattcagatt	tcactaaaat	cttccaagat	1440
gaaagcaaaa	aaaaaaaaaa	aaaaaaaaaa	actcgagggg	gggcccgtac	ccaattcgcc	1500
ctatagttag	tcgtatt					1517

&lt;210&gt; 65

&lt;211&gt; 526

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

46

<220>  
 <221> SITE  
 <222> (66)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (106)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (484)  
 <223> n equals a,t,g, or c

<400> 65  
 ctctgacagc ttctcttttg gccaaagcct gcctctgtac agcctcgagt ggacagccag 60  
 aggtcnagac tggagcccag agcccaagat ggagcccag ctgggnccctg aggctgccgc 120  
 cctccgccct ggctggctgg cctgtctgct gtgggtctca gccctgagct gttctttctc 180  
 cttgccagct tcttcccttt cttctctggt gcccacagtc agaaccagct acaatttttg 240  
 aaggactttc ctgggtcttg ataaatgcaa tgcctgcac gccacatcta ttgcaagaa 300  
 gttcttttaa gaagaaataa gatctgacaa ctggctggct tcccacdttg ggactgcctc 360  
 ccgattccct ttggtttctt atccttgcaa attactccar atgattycca aaatctggsg 420  
 sccttggtga ratcttttaa ctggtcagca awtwtcaaac gaaatctcca aacaggaaat 480  
 cttntgcctc ctgcatccac ccccaaagaa cttgcacatt gacgtt 526

<210> 66  
 <211> 664  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (31)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (63)  
 <223> n equals a,t,g, or c

<400> 66  
 caggctctca atacggactc actcataggg naaagctggt acgcctgcag gtaccgggtcc 60  
 ggnaattccc gggctcgacc acgcgtcgcr gagctcttac ttctccagca acrtcttca 120  
 gtacataata agcttaactg ataaacagaa tatttagaaa ggtgagactt gggcttacca 180  
 ttgggtttta atcataggga cctagggcga ggggttcagg cttctctgga gcagatattg 240  
 tcaagttcat ggccttaggt agcatgtatc tgggtctaac tctgattgta gcaaaagttc 300  
 tgagaggagc tgagccctgt tgtggcccat taaagaacag ggtcctcagg ccctgccgcg 360  
 ttctgttcca ctgccccctc cccatcccca gccagccga ggaatcccg tgggttgctt 420  
 acctacctat aagggtggtt ataagctgct gtcctggcca ctgcattcaa attccaatgt 480  
 gtacttcata gtgtaaaaat ttatatattt gtgaggtttt ttgtcttttt tttttttttt 540  
 ttttttggtt tattgctgta tctactttta cttccagaaa taaacgttat atrggaaaaa 600  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 660  
 aaaa 664

<210> 67  
 <211> 156

47

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (156)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 67

Met Arg Leu Trp Lys Ala Val Val Val Thr Leu Ala Phe Met Ser Val  
 1 5 10 15

Asp Ile Cys Val Thr Thr Ala Ile Tyr Val Phe Ser His Leu Asp Arg  
 20 25 30

Ser Leu Leu Glu Asp Ile Arg His Phe Asn Ile Phe Asp Ser Val Leu  
 35 40 45

Asp Leu Trp Ala Ala Cys Leu Tyr Arg Ser Cys Leu Leu Leu Gly Ser  
 50 55 60

His His Trp Cys Gly Gln Glu Gln Cys Ala Gly Ala Pro Ala Ala Ala  
 65 70 75 80

Gly Leu Val Ala Gly His His Pro Arg Val Pro Leu Arg Gly His Leu  
 85 90 95

Cys His Gly Glu Ala Ala Ala Leu Leu Arg Gly Ala Gln Ala His Pro  
 100 105 110

Gly Pro Leu Val Leu Gly Pro Val Arg Val Asp Val His Phe Thr Arg  
 115 120 125

Arg Ile Leu Pro Ala Leu Val Ala Ala Val His Arg Ala Ala Arg His  
 130 135 140

Pro Gly Pro Gly Ala Arg Gly Gly His Arg Gly Xaa  
 145 150 155

&lt;210&gt; 68

&lt;211&gt; 70

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (70)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 68

Met Ala Ala Arg His Leu Pro Gly Phe His Thr Tyr Thr Asn Leu Leu  
 1 5 10 15

Phe Leu Leu Leu Pro Ser Leu Leu Met Gly Tyr Ser Glu Ser Pro Pro  
 20 25 30

Pro Ile Thr Asp Ser Trp Ala Pro Phe Ile Ser Leu Thr His His Val  
 35 40 45

48

Leu Ser Gln Ser Gln Ser Pro Leu Ser Ser Asn Cys Trp Ile Cys Leu  
 50 55 60

Ser Thr His Thr Gln Xaa  
 65 70

&lt;210&gt; 69

&lt;211&gt; 502

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (502)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 69

Met Trp Lys Leu Trp Arg Ala Glu Glu Gly Ala Ala Ala Leu Gly Gly  
 1 5 10 15

Ala Leu Phe Leu Leu Leu Phe Ala Leu Gly Val Arg Gln Leu Leu Lys  
 20 25 30

Gln Arg Arg Pro Met Gly Phe Pro Pro Gly Pro Pro Gly Leu Pro Phe  
 35 40 45

Ile Gly Asn Ile Tyr Ser Leu Ala Ala Ser Ser Glu Leu Pro His Val  
 50 55 60

Tyr Met Arg Lys Gln Ser Gln Val Tyr Gly Glu Ile Phe Ser Leu Asp  
 65 70 75 80

Leu Gly Gly Ile Ser Thr Val Val Leu Asn Gly Tyr Asp Val Val Lys  
 85 90 95

Glu Cys Leu Val His Gln Ser Glu Ile Phe Ala Asp Arg Pro Cys Leu  
 100 105 110

Pro Leu Phe Met Lys Met Thr Lys Met Gly Gly Leu Leu Asn Ser Arg  
 115 120 125

Tyr Gly Arg Gly Trp Val Asp His Arg Arg Leu Ala Val Asn Ser Phe  
 130 135 140

Arg Tyr Phe Gly Tyr Gly Gln Lys Ser Phe Glu Ser Lys Ile Leu Glu  
 145 150 155 160

Glu Thr Lys Phe Phe Asn Asp Ala Ile Glu Thr Tyr Lys Gly Arg Pro  
 165 170 175

Phe Asp Phe Lys Gln Leu Ile Thr Asn Ala Val Ser Asn Ile Thr Asn  
 180 185 190

Leu Ile Ile Phe Gly Glu Arg Phe Thr Tyr Glu Asp Thr Asp Phe Gln  
 195 200 205

His Met Ile Glu Leu Phe Ser Glu Asn Val Glu Leu Ala Ala Ser Ala

49

210                      215                      220

Ser Val Phe Leu Tyr Asn Ala Phe Pro Trp Ile Gly Ile Leu Pro Phe  
 225                      230                      235                      240

Gly Lys His Gln Gln Leu Phe Arg Asn Ala Ala Val Val Tyr Asp Phe  
                     245                      250                      255

Leu Ser Arg Leu Ile Glu Lys Ala Ser Val Asn Arg Lys Pro Gln Leu  
                     260                      265                      270

Pro Gln His Phe Val Asp Ala Tyr Leu Asp Glu Met Asp Gln Gly Lys  
                     275                      280                      285

Asn Asp Pro Ser Ser Thr Phe Ser Lys Glu Asn Leu Ile Phe Ser Val  
                     290                      295                      300

Gly Glu Leu Ile Ile Ala Gly Thr Glu Thr Thr Thr Asn Val Leu Arg  
 305                      310                      315                      320

Trp Ala Ile Leu Phe Met Ala Leu Tyr Pro Asn Ile Gln Gly Gln Val  
                     325                      330                      335

Gln Lys Glu Ile Asp Leu Ile Met Gly Pro Asn Gly Lys Pro Ser Trp  
                     340                      345                      350

Asp Asp Lys Cys Lys Met Pro Tyr Thr Glu Ala Val Leu His Glu Val  
                     355                      360                      365

Leu Arg Phe Cys Asn Ile Val Pro Leu Gly Ile Phe His Ala Thr Ser  
                     370                      375                      380

Glu Asp Ala Val Val Arg Gly Tyr Ser Ile Pro Lys Gly Thr Thr Val  
 385                      390                      395                      400

Ile Thr Asn Leu Tyr Ser Val His Phe Asp Glu Lys Tyr Trp Arg Asp  
                     405                      410                      415

Pro Glu Val Phe His Pro Glu Arg Phe Leu Asp Ser Ser Gly Tyr Phe  
                     420                      425                      430

Ala Lys Lys Glu Ala Leu Val Pro Phe Ser Leu Gly Arg Arg His Cys  
                     435                      440                      445

Leu Gly Glu His Leu Ala Arg Met Glu Met Phe Leu Phe Phe Thr Ala  
                     450                      455                      460

Leu Leu Gln Arg Phe His Leu His Phe Pro His Glu Leu Val Pro Asp  
 465                      470                      475                      480

Leu Lys Pro Arg Leu Gly Met Thr Leu Gln Pro Gln Pro Tyr Leu Ile  
                     485                      490                      495

Cys Ala Glu Arg Arg Xaa  
                     500

&lt;210&gt; 70

&lt;211&gt; 189



50

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (85)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (104)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (164)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (189)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 70

Met	Arg	Pro	Ala	Phe	Ala	Leu	Cys	Leu	Leu	Trp	Gln	Ala	Leu	Trp	Pro
1					5				10					15	

Gly	Pro	Gly	Gly	Gly	Glu	His	Pro	Thr	Ala	Asp	Arg	Ala	Gly	Cys	Ser
			20					25					30		

Ala	Ser	Gly	Ala	Cys	Tyr	Ser	Leu	His	His	Ala	Thr	Met	Lys	Arg	Gln
		35					40					45			

Ala	Ala	Glu	Glu	Ala	Cys	Ile	Leu	Arg	Gly	Gly	Ala	Leu	Ser	Thr	Val
	50					55					60				

Arg	Ala	Gly	Ala	Glu	Leu	Arg	Ala	Val	Leu	Ala	Leu	Leu	Arg	Ala	Gly
65					70					75					80

Pro	Gly	Pro	Gly	Xaa	Gly	Ser	Lys	Asp	Leu	Leu	Phe	Trp	Val	Ala	Leu
			85						90					95	

Glu	Arg	Arg	Arg	Ser	His	Cys	Xaa	Leu	Glu	Asn	Glu	Pro	Leu	Arg	Gly
			100					105					110		

Phe	Ser	Trp	Leu	Ser	Ser	Asp	Pro	Gly	Gly	Leu	Glu	Ser	Asp	Thr	Leu
		115					120					125			

Gln	Trp	Val	Glu	Glu	Pro	Gln	Arg	Ser	Cys	Thr	Ala	Arg	Arg	Trp	Val
	130					135					140				

Leu	Pro	Gly	His	Arg	Trp	Gly	Arg	Ala	Arg	Ser	Trp	Lys	Glu	Met	Arg
145					150					155				160	

Cys	His	Leu	Xaa	Ala	Asn	Ala	Thr	Cys	Ala	Ser	Thr	Ser	Leu	Arg	Ser
			165						170					175	

Cys	Val	Leu	Arg	Arg	Ala	Pro	Gly	Pro	Pro	Leu	Thr	Xaa
			180					185				

<210> 71  
 <211> 486  
 <212> PRT  
 <213> Homo sapiens

<400> 71

```

Met Gln Pro Ser Gly Leu Glu Gly Pro Gly Thr Phe Gly Arg Trp Pro
 1           5           10           15

Leu Leu Ser Leu Leu Leu Leu Leu Leu Leu Gln Pro Val Thr Cys
          20           25           30

Ala Tyr Thr Thr Pro Gly Pro Pro Arg Ala Leu Thr Thr Leu Gly Ala
          35           40           45

Pro Arg Ala His Thr Met Pro Gly Thr Tyr Ala Pro Ser Thr Thr Leu
          50           55           60

Ser Ser Pro Ser Thr Gln Gly Leu Gln Glu Gln Ala Arg Ala Leu Met
65           70           75           80

Arg Asp Phe Pro Leu Val Asp Gly His Asn Asp Leu Pro Leu Val Leu
          85           90           95

Arg Gln Val Tyr Gln Lys Gly Leu Gln Asp Val Asn Leu Arg Asn Phe
          100          105          110

Ser Tyr Gly Gln Thr Ser Leu Asp Arg Leu Arg Asp Gly Leu Val Gly
          115          120          125

Ala Gln Phe Trp Ser Ala Tyr Val Pro Cys Gln Thr Gln Asp Arg Asp
          130          135          140

Ala Leu Arg Leu Thr Leu Glu Gln Ile Asp Leu Ile Arg Arg Met Cys
145          150          155          160

Ala Ser Tyr Ser Glu Leu Glu Leu Val Thr Ser Ala Lys Ala Leu Asn
          165          170          175

Asp Thr Gln Lys Leu Ala Cys Leu Ile Gly Val Glu Gly Gly His Ser
          180          185          190

Leu Asp Asn Ser Leu Ser Ile Leu Arg Thr Phe Tyr Met Leu Gly Val
          195          200          205

Arg Tyr Leu Thr Leu Thr His Thr Cys Asn Thr Pro Trp Ala Glu Ser
          210          215          220

Ser Ala Lys Gly Val His Ser Phe Tyr Asn Asn Ile Ser Gly Leu Thr
225          230          235          240

Asp Phe Gly Glu Lys Val Val Ala Glu Met Asn Arg Leu Gly Met Met
          245          250          255

Val Asp Leu Ser His Val Ser Asp Ala Val Ala Arg Arg Ala Leu Glu
          260          265          270

```

52

Val Ser Gln Ala Pro Val Ile Phe Ser His Ser Ala Ala Arg Gly Val  
 275 280 285

Cys Asn Ser Ala Arg Asn Val Pro Asp Asp Ile Leu Gln Leu Leu Lys  
 290 295 300

Lys Asn Gly Gly Val Val Met Val Ser Leu Ser Met Gly Val Ile Gln  
 305 310 315 320

Cys Asn Pro Ser Ala Asn Val Ser Thr Val Ala Asp His Phe Asp His  
 325 330 335

Ile Lys Ala Val Ile Gly Ser Lys Phe Ile Gly Ile Gly Gly Asp Tyr  
 340 345 350

Asp Gly Ala Gly Lys Phe Pro Gln Gly Leu Glu Asp Val Ser Thr Tyr  
 355 360 365

Pro Val Leu Ile Glu Glu Leu Leu Ser Arg Gly Trp Ser Glu Glu Glu  
 370 375 380

Leu Gln Gly Val Leu Arg Gly Asn Leu Leu Arg Val Phe Arg Gln Val  
 385 390 395 400

Glu Lys Val Gln Glu Glu Asn Lys Trp Gln Ser Pro Leu Glu Asp Lys  
 405 410 415

Phe Pro Asp Glu Gln Leu Ser Ser Ser Cys His Ser Asp Leu Ser Arg  
 420 425 430

Leu Arg Gln Arg Gln Ser Leu Thr Ser Gly Gln Glu Leu Thr Glu Ile  
 435 440 445

Pro Ile His Trp Thr Ala Lys Leu Pro Ala Lys Trp Ser Val Ser Glu  
 450 455 460

Ser Ser Pro His Met Ala Pro Val Leu Ala Val Val Ala Thr Phe Pro  
 465 470 475 480

Val Leu Ile Leu Trp Leu  
 485

&lt;210&gt; 72

&lt;211&gt; 88

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (88)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 72

Met Val Ala Ser Gly Trp Leu Leu Leu Ala Gln Ala Ser Phe Leu Pro  
 1 5 10 15

Leu Ala Pro Pro Gly Ala Leu Gly Ala Gly Cys Trp Met Asp Gly Arg  
 20 25 30

53

Pro Leu Ala Pro Pro Gly Ala Leu Gly Ala Gly Cys Trp Met Gly Gly  
           35                          40                          45

Arg Pro Leu Ala Pro Pro Gly Ala Leu Gly Ala Gly Cys Trp Met Gly  
       50                          55                          60

Gly Arg His Gly Ala Pro Leu Leu Gly Cys Leu Cys Pro Ser Gly Leu  
       65                          70                          75                          80

Cys Ser Ser Tyr Val Cys Leu Xaa  
                           85

<210> 73  
 <211> 299  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (167)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 73  
 Met Met Ser Ser Lys Pro Thr Ser His Ala Glu Val Asn Glu Thr Ile  
       1                          5                          10                          15

Pro Asn Pro Tyr Pro Pro Ser Ser Phe Met Ala Pro Gly Phe Gln Gln  
                           20                          25                          30

Pro Leu Gly Ser Ile Asn Leu Glu Asn Gln Ala Gln Gly Ala Gln Arg  
           35                          40                          45

Ala Gln Pro Tyr Gly Ile Thr Ser Pro Gly Ile Phe Ala Ser Ser Gln  
       50                          55                          60

Pro Gly Gln Gly Asn Ile Gln Met Ile Asn Pro Ser Val Gly Thr Ala  
       65                          70                          75                          80

Val Met Asn Phe Lys Glu Glu Ala Lys Ala Leu Gly Val Ile Gln Ile  
                           85                          90                          95

Met Val Gly Leu Met His Ile Gly Phe Gly Ile Val Leu Cys Leu Ile  
           100                          105                          110

Ser Phe Ser Phe Arg Glu Val Leu Gly Phe Ala Ser Thr Ala Val Ile  
           115                          120                          125

Gly Gly Tyr Pro Phe Trp Gly Gly Leu Ser Phe Ile Ile Ser Gly Ser  
       130                          135                          140

Leu Ser Val Ser Ala Ser Lys Glu Leu Ser Arg Cys Leu Val Lys Gly  
       145                          150                          155                          160

Ser Leu Gly Met Asn Ile Xaa Ser Ser Ile Leu Ala Phe Ile Gly Val  
           165                          170                          175

Ile Leu Leu Leu Val Asp Met Cys Ile Asn Gly Val Ala Gly Gln Asp

54

180	185	190
Tyr Trp Ala Val Leu Ser Gly Lys Gly Ile Ser Ala Thr Leu Met Ile		
195	200	205
Phe Ser Leu Leu Glu Phe Phe Val Ala Cys Ala Thr Ala His Phe Ala		
210	215	220
Asn Gln Ala Asn Thr Thr Thr Asn Met Ser Val Leu Val Ile Pro Asn		
225	230	235
Met Tyr Glu Ser Asn Pro Val Thr Pro Ala Ser Ser Ser Ala Pro Pro		
245	250	255
Arg Cys Asn Asn Tyr Ser Ala Asn Ala Pro Lys Arg Lys Arg Gly Ile		
260	265	270
Ser Leu Ile Ser Trp Arg Lys Thr Thr Cys Lys Asn Phe Leu Arg Arg		
275	280	285
Cys Leu Leu Leu Ser Thr Met Ile Ser Ser Leu		
290	295	

<210> 74  
 <211> 48  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (48)  
 <223> Xaa equals stop translation

<400> 74

Met Ala Leu His Pro Gly Ser Ser His Leu Leu Val Ala Val Pro Val
1                      5                      10                      15
Ser Trp Phe Leu Phe Cys Ile Pro Gly Ile Ser Phe Ile Thr Leu Ser
20                      25                      30
Trp Ser Tyr Gln Glu Ser Pro Val Ser Phe Leu Ser Val Glu Gly Xaa
35                      40                      45

<210> 75  
 <211> 44  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (44)  
 <223> Xaa equals stop translation

<400> 75

Met Tyr Ser Leu Phe Leu Thr Cys Ile Phe Pro Phe Thr Leu Cys His  
 1 5 10 15

Lys Lys Ile Leu Met Val Ile His Asp Phe Thr Gly Pro Val His Val  
 20 25 30

Phe Pro Glu Lys Thr Val Leu Glu Trp Asn Tyr Xaa  
 35 40

<210> 76

<211> 140

<212> PRT

<213> Homo sapiens

<400> 76

Met Cys Ala Met Tyr Leu Met Ile Lys Ala Phe Leu Pro Lys Met Leu  
 1 5 10 15

Ala Gln Lys Ser Gly Asn Ile Ile Asn Met Ser Ser Val Ala Ser Ser  
 20 25 30

Val Lys Gly Val Val Asn Arg Cys Val Tyr Ser Thr Thr Lys Ala Ala  
 35 40 45

Val Ile Gly Leu Thr Lys Ser Val Ala Ala Asp Phe Ile Gln Gln Gly  
 50 55 60

Ile Arg Cys Asn Cys Val Cys Pro Gly Thr Val Asp Thr Pro Ser Leu  
 65 70 75 80

Gln Glu Arg Ile Gln Ala Arg Gly Asn Pro Glu Glu Ala Arg Asn Asp  
 85 90 95

Phe Leu Lys Arg Gln Lys Thr Gly Arg Phe Ala Thr Ala Glu Glu Ile  
 100 105 110

Ala Met Leu Cys Val Tyr Leu Ala Ser Asp Glu Ser Ala Tyr Val Thr  
 115 120 125

Gly Asn Pro Val Ile Ile Asp Gly Gly Trp Ser Leu  
 130 135 140

<210> 77

<211> 153

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (153)

<223> Xaa equals stop translation

<400> 77

Met Leu Val Val Cys Leu Leu Leu Ala Thr Gly Phe Cys Leu Phe Arg  
 1 5 10 15

Gly Leu Ile Ala Leu Asp Cys Pro Ser Glu Leu Cys Arg Leu Tyr Thr

```
<210> 78
<211> 180
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (180)  
<223> Xaa equals stop translation
```

```

<400> 78
Met Ala Ala Ala Ser Ala Gly Ala Thr Arg Leu Leu Leu Leu Leu
  1              5              10              15

Met Ala Val Ala Ala Pro Ser Arg Ala Arg Gly Ser Gly Cys Arg Ala
      20              25              30

Gly Thr Gly Ala Arg Gly Ala Gly Ala Glu Gly Arg Glu Gly Glu Xaa
      35              40              45

Pro Val Ser Ser Ala Ile Pro Arg Arg Val Cys Trp Ser Leu Leu Ser
      50              55              60

Pro Arg Pro Thr Arg Pro Gly Pro Ala Pro Cys Pro Leu Pro Ser
  65              70              75              80

Ala Gly Arg Gly Ala Ala Gly Leu Gly Pro Leu Ala Gln Gln Pro Val

```

85 57 95  
 90  
 Ser Pro Ala Pro Ala Ser Pro Met Ala Pro Cys Ser Pro Arg Gly Phe  
 100 105 110  
 Pro Pro Ala His Gly Val Glu Pro Glu Ile Leu Ala Thr Met Pro Val  
 115 120 125  
 Leu Thr Ser His Pro Pro Thr Pro Ser Pro Cys Ser Leu Gly Thr Cys  
 130 135 140  
 Arg Leu Leu Ser Ser Leu Cys Ala Phe Val Pro Gly Gly Leu Thr Leu  
 145 150 155 160  
 Leu Ser Leu Ala Gly Leu Gly Gly Pro Val Gln Ala Pro Ala Ala Pro  
 165 170 175  
 Pro Ser Leu Xaa  
 180

<210> 79  
 <211> 70  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (70)  
 <223> Xaa equals stop translation

<400> 79  
 Met Leu Met Gly Ser Ile Leu Tyr Val Leu Phe Cys Val Trp Leu Leu  
 1 5 10 15  
 Gln Cys Ile Phe Glu Ile Tyr Pro His Cys Cys Val Tyr Pro Lys Cys  
 20 25 30  
 Val Leu Phe His Cys Gln Ile Met Phe Cys Tyr Met Asn Ile Leu Gln  
 35 40 45  
 Asn Ile Cys Leu Phe Ile Tyr Trp Trp Ile Phe Ala Phe Val Pro Val  
 50 55 60  
 Trp Gly Tyr Tyr Glu Xaa  
 65 70

<210> 80  
 <211> 191  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (191)  
 <223> Xaa equals stop translation

<400> 80



58  
 Met Arg Ala Cys Pro Trp Ala Gln Val Pro Leu Tyr Leu Leu Leu Asp  
 1 5 10 15  
 Gly His Leu Ala Val Ser Gln Ala Gly Val Met Ala Gly Val Ser Gly  
 20 25 30  
 Gly Arg Gly Gly Arg Arg Leu Arg Gly Pro Ile Thr Ser Arg Val Ile  
 35 40 45  
 Thr Ser Cys Gln Gln Pro Gly Val Gly Val Trp Val Ser Leu Arg Pro  
 50 55 60  
 Glu Leu Leu Asn Leu Glu Ser Leu Gly Val Ala Ala Lys Gly Val Tyr  
 65 70 75 80  
 Asp Lys His Val Ser Leu Asp Ile Ser Gly Glu Arg Ser Gly Ala Leu  
 85 90 95  
 Val Thr Phe Ser Lys Gly Cys Trp Ala Ser Glu Gln Ser Pro Pro Met  
 100 105 110  
 Ser Gln Pro Leu Gln Gly Pro Ser Leu Ser Leu His Pro Arg Pro Ser  
 115 120 125  
 Ala Ala Leu Val Met Ser Arg Arg Lys Val Leu Gly Cys Ala Gln Ser  
 130 135 140  
 Gln Glu Ser Lys Ile Cys Gln Ala Lys Ala Pro Gly Lys Ser Arg Arg  
 145 150 155 160  
 Ser Leu Gly Trp Pro Pro Gly Cys Gly Ala Ala Arg Ala Lys Thr Val  
 165 170 175  
 Asn Thr Ala Leu Gln Leu Ser Glu Pro Gln Phe Ser Asn Leu Xaa  
 180 185 190

&lt;210&gt; 81

&lt;211&gt; 166

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (127)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (166)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 81

Met Cys Leu Ser Leu Leu Ala Ala Leu Ala Cys Ser Ala Gly Asp Thr  
 1 5 10 15

Trp Ala Ser Glu Val Gly Pro Val Leu Ser Lys Ser Ser Pro Arg Leu  
 20 25 30

Ile Thr Thr Trp Glu Lys Val Pro Val Gly Thr Asn Gly Gly Val Thr  
                   35                  40                  59                  45  
 Val Val Gly Leu Val Ser Ser Leu Leu Gly Gly Thr Phe Val Gly Ile  
           50                  55                  60  
 Ala Tyr Phe Leu Thr Gln Leu Ile Phe Val Asn Asp Leu Asp Ile Ser  
           65                  70                  75                  80  
 Ala Pro Gln Trp Pro Ile Ile Ala Phe Gly Gly Leu Ala Gly Leu Leu  
                   85                  90                  95  
 Gly Ser Ile Val Asp Ser Tyr Leu Gly Ala Thr Met Gln Tyr Thr Gly  
                   100                  105                  110  
 Leu Asp Glu Ser Thr Gly Met Val Val Asn Ser Pro Thr Asn Xaa Ala  
           115                  120                  125  
 Arg His Ile Ala Gly Lys Pro Ile Leu Asp Asn Asn Ala Val Asn Leu  
           130                  135                  140  
 Phe Ser Ser Val Leu Ile Ala Leu Leu Leu Pro Thr Ala Ala Trp Gly  
           145                  150                  155                  160  
 Phe Trp Pro Arg Gly Xaa  
                   165

<210> 82  
 <211> 42  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (42)  
 <223> Xaa equals stop translation

<400> 82  
 Met Cys Gly Leu Val Ile Leu Trp Pro Cys Ile Met Thr Leu Phe Ser  
           1                  5                  10                  15  
 Ser Leu Ser Thr Gly Asp Val Leu Leu Pro Cys Lys Ile Leu Val Gly  
                   20                  25                  30  
 Leu Arg Val Phe Ile Gly Ala Arg Val Xaa  
           35                  40

<210> 83  
 <211> 49  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (49)  
 <223> Xaa equals stop translation

60  
<400> 83  
Met Cys Phe Pro Ala Cys Leu Cys Ser Pro Leu Thr Cys Leu Leu Ser  
1 5 10 15  
Val Trp Lys Pro Gly Leu Ala His Ala Val Val His Cys Met Leu Glu  
20 25 30  
Pro Val Glu Phe Ala Arg Val Val Gln Tyr Glu Ala Gly His Val Leu  
35 40 45  
Xaa

<210> 84  
<211> 57,  
<212> PRT  
<213> Homo sapiens  
  
<220>  
<221> SITE  
<222> (57)  
<223> Xaa equals stop translation

<400> 84  
Met Leu Ile Ala Lys Leu Pro Val Leu Glu Ser Ile Cys Phe Phe Met  
1 5 10 15  
Leu Phe Leu Asn Pro Leu Val Ile Leu Leu Ser Leu Asn Asn Ala Leu  
20 25 30  
Pro Leu Val Phe His Pro His Ser Glu Phe Leu Glu Asp His Asn Arg  
35 40 45  
Gly Asp Thr Leu Pro Ser Ile Val Xaa  
50 55

<210> 85  
<211> 43  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (43)  
<223> Xaa equals stop translation

<400> 85  
Met Leu Val Ala Thr Ala Val Cys Cys Tyr Leu Phe Trp Leu Ile Ala  
1 5 10 15  
Ile Leu Ala Gln Leu Asn Pro Leu Phe Gly Pro Gln Leu Lys Asn Glu  
20 25 30  
Thr Ile Trp Tyr Val Arg Phe Leu Trp Glu Xaa  
35 40

61

<210> 86  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (41)  
 <223> Xaa equals stop translation

<400> 86  
 Met Leu Leu Leu Trp Ala Phe Ser Gly Val Cys Ala Val Pro Ala Arg  
           1                  5                  10                  15  
 Ala Thr Pro Val Pro Ser Ser Phe Cys Pro Gln Gly Pro Ser Leu Cys  
                   20                  25                  30  
 Pro Lys Gln Pro Ala Ser Leu Ala Xaa  
           35                  40

<210> 87  
 <211> 74  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (74)  
 <223> Xaa equals stop translation

<400> 87  
 Met His Ala Tyr Ala Cys Val Cys Ala Cys Met Leu Val Cys Val Cys  
           1                  5                  10                  15  
 Val Cys Val Cys Arg Ala Leu Val Ile Pro Thr Glu Gln Arg His Arg  
                   20                  25                  30  
 Arg Val Ala His Gly Arg Thr Ser Asp Ser Thr Leu Pro Cys Thr Val  
           35                  40                  45  
 Lys Ile Trp Pro Ser Glu Arg Gly Asp Gly Arg Gly Glu Arg Gly Glu  
           50                  55                  60  
 Arg Arg Arg Gly Thr Asp Trp Arg Gly Xaa  
           65                  70

<210> 88  
 <211> 47  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (34)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>

62.  
 <221> SITE  
 <222> (47)  
 <223> Xaa equals stop translation  
 <400> 88  
 Met His His Pro Asn Leu Cys Leu His Phe His Ala Ala Phe Ser Leu  
 1 5 10 15  
 Cys Val His Gly Cys Leu Cys Val Gln Phe Phe Pro Phe Tyr Lys Asp  
 20 25 30  
 Thr Xaa His Ile Gly Leu Glu Pro Thr Leu Met Thr Ser Ser Xaa  
 35 40 45

<210> 89  
 <211> 63  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (63)  
 <223> Xaa equals stop translation

<400> 89  
 Met Leu Phe Leu Asn Val Ile Leu Phe Ser Leu Thr Val Phe Thr Leu  
 1 5 10 15  
 Ile Ser Thr Ala His Thr Leu Asp Arg Ala Val Arg Ser Asp Trp Leu  
 20 25 30  
 Leu Leu Val Leu Ile Tyr Ala Cys Leu Glu Glu Leu Ile Pro Glu Leu  
 35 40 45  
 Ile Phe Asn Leu Tyr Cys Gln Gly Asn Ala Thr Leu Phe Phe Xaa  
 50 55 60

<210> 90  
 <211> 70  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (20)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (70)  
 <223> Xaa equals stop translation

<400> 90  
 Met Leu Leu Lys Leu His Thr Leu Trp Pro Leu Trp Pro Gly Leu Trp  
 1 5 10 15  
 Ala Thr Thr Xaa Ser Asp Ser Leu Gly Glu Arg Thr His Ser Leu Cys

20 25 63 30  
 Arg Arg Lys Lys Ala Ser Leu Ser Thr Gly Trp Met Ser Trp Met Ser  
 35 40 45  
 Cys Arg Ala Arg Ala Thr His Thr Gln Val Val Ser Leu Lys Asp Lys  
 50 55 60  
 Val Glu Phe Ala Pro Xaa  
 65 70

<210> 91  
 <211> 57  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (57)  
 <223> Xaa equals stop translation

<400> 91  
 Met Lys Glu Ser Arg Lys Met Leu Trp Val Phe Lys Met Leu Phe Phe  
 1 5 10 15  
 Lys Ile Val Leu Trp Val Asn Leu Leu Ser Ala Ala Leu Ser Cys Ile  
 20 25 30  
 Gln Lys Gln Met Leu Gly Ile Ala Pro Gln Lys Cys Val Pro Lys Leu  
 35 40 45  
 Cys Phe Gln Leu Tyr Ile Met Arg Xaa  
 50 55

<210> 92  
 <211> 68  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (68)  
 <223> Xaa equals stop translation

<400> 92  
 Met Tyr Phe Leu Leu Ser Val Thr Ser Glu Ser Val Trp Arg Ser Trp  
 1 5 10 15  
 Thr Leu Thr Phe His Ser Phe Ala Ile Leu Ser Leu Arg Cys Trp Thr  
 20 25 30  
 Ser Leu Leu Leu Leu Ile Pro Leu Thr Ser Cys Asn Phe Ser Ser Pro  
 35 40 45  
 Ser Trp Arg Met Thr Ala Ser Gln Val Pro Ser Lys Arg Lys Ala Ser  
 50 55 60

64

Met Thr Leu Xaa  
65

<210> 93  
<211> 45  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (45)  
<223> Xaa equals stop translation

<400> 93  
Met Lys Gly Trp Pro Val Phe Leu Leu Val Gln Ala Val Thr Phe Leu  
1 5 10 15

Ser Val Ala Gln Ser Gly Ala Met Ala Cys Ala Ala Ser Gly Val Val  
20 25 30

Tyr Ser Val Asp Val Pro Ala Cys Ser Ser Arg Ser Xaa  
35 40 45

<210> 94  
<211> 55  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (55)  
<223> Xaa equals stop translation

<400> 94  
Met Val Leu Ser Pro Trp Ala Cys Leu Phe Val Val Phe Phe Pro Tyr  
1 5 10 15

Ile Gln Ser Ser Leu Arg Ser Asp Lys His Leu Gln Leu Ser Asn Ile  
20 25 30

Leu Pro Thr Pro Ser His His Ile His Leu Pro Ala Ser Ile Cys Ile  
35 40 45

Gln Leu Arg Ala Gly Asn Xaa  
50 55

<210> 95  
<211> 41  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (41)  
<223> Xaa equals stop translation

65.  
 <400> 95  
 Met Cys Glu Tyr Val Leu Leu Leu Tyr Ile Val Leu Leu Cys Asn Arg  
 1 5 10 15  
 Ser Tyr Ala Val Phe Thr Gln Cys Val Leu Arg Ser Ser Pro Ile Asp  
 20 25 30  
 Ser Ser Arg Asn Ala Val Leu Leu Xaa  
 35 40

<210> 96  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> (41)  
 <223> Xaa equals stop translation

<400> 96  
 Met Thr Thr Pro Gly Leu Leu Ile Leu Phe Leu Ala His Val Cys Leu  
 1 5 10 15  
 Val Asn His Gln Gln Ala Ala Glu Pro Gly Trp Lys Gln His Cys Cys  
 20 25 30  
 Asn Trp Glu Gly His Arg Val Leu Xaa  
 35 40

<210> 97  
 <211> 50  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> (14)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (50)  
 <223> Xaa equals stop translation

<400> 97  
 Met Leu Cys His Val Tyr Leu Leu Leu Val Gly His Ala Xaa Phe Ser  
 1 5 10 15  
 Val Gly Leu Met Gly Gln Arg Lys Leu Arg Cys Ser Ile Asn Ser Ala  
 20 25 30  
 Leu Arg Ser Ala Val Ser Ser Ala Trp Asn Ser Ser Ile Cys Phe Asn  
 35 40 45  
 Ser Xaa  
 50



66

<210> 98  
<211> 58  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (58)  
<223> Xaa equals stop translation

<400> 98  
Met Ser Glu Trp Cys Gln Pro Asp Gln Ile Leu Leu Gln Phe Pro Val  
1 5 10 15  
Leu Ala Thr Met Ser Val Ala Phe Leu Ile Gln Arg Cys Phe Cys Phe  
20 25 30  
Trp Trp Phe Val Leu Asn Ala Phe Ser Ile Pro Ser Gly Thr Glu Lys  
35 40 45  
Lys Arg Ile Val Phe Lys Lys Trp Leu Xaa  
50 55

<210> 99  
<211> 52  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (52)  
<223> Xaa equals stop translation

<400> 99  
Met Lys Val Val Val Val Met Val Val Ile Leu Val Val Val Thr Leu  
1 5 10 15  
Val Val Val Val Met Val Val Ile Leu Val Met Val Val Met Val Val  
20 25 30  
Ala Leu Val Thr Leu Thr Trp Gly Pro Val Ala Val Thr Val Asp Ala  
35 40 45  
Gly Ser Trp Xaa  
50

<210> 100  
<211> 45  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (45)  
<223> Xaa equals stop translation

67

&lt;400&gt; 100

Met Pro His Phe Leu Arg Trp Leu Leu Thr Thr Phe Arg Ile Arg Ala  
 1 5 10 15

Ser Cys Gly Ser Thr Pro Cys Trp Ser Pro Ser His Leu Gly Cys Leu  
 20 25 30

Gln Pro Ala Leu Pro Arg Asp Leu Ser His Leu Glu Xaa  
 35 40 45

&lt;210&gt; 101

&lt;211&gt; 58

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (58)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 101

Met Ser Thr Lys Ile Leu Gln Phe Leu Phe Ser Ser Cys Cys Trp Val  
 1 5 10 15

Pro Pro Met Leu Phe Leu Phe Lys Asn Thr Lys Cys Arg Thr Ser Leu  
 20 25 30

Leu Tyr Cys Phe Tyr Phe Ile Leu Leu Thr Cys Ser Leu Ser Glu Tyr  
 35 40 45

Asp Ser Leu Leu Ser Ser Lys Val Phe Xaa  
 50 55

&lt;210&gt; 102

&lt;211&gt; 41

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (41)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 102

Met Phe Trp Phe Trp Phe Leu Leu Ser Leu Ser Phe Gln Gln Val Glu  
 1 5 10 15

Gln Gln Gln Val Phe Gln Cys Ile Cys Cys Thr Arg Thr Lys Tyr Lys  
 20 25 30

Ser Val Trp His Gln Lys Ser Lys Xaa  
 35 40

&lt;210&gt; 103

&lt;211&gt; 143

68

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (104)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (105)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (115)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (143)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 103

Met	Thr	Leu	Ile	Glu	Val	Leu	Val	Ser	Val	Leu	Ile	Leu	Ala	Val	Gly
1				5					10					15	

Leu	Leu	Arg	Ala	Ala	Val	Ile	Gln	Leu	Asn	Ala	Leu	Lys	Tyr	Thr	Asp
			20					25					30		

Ser	Ser	Arg	Met	Thr	Ser	Gln	Ala	Ser	Phe	Ile	Ala	Tyr	Asp	Met	Leu
		35					40					45			

Asp	Arg	Ile	Arg	Ala	Asn	Ser	Gly	Ala	Asp	Tyr	Ser	Trp	Gly	Gln	Gly
	50					55					60				

Glu	Arg	Ala	Pro	Ser	Thr	Ser	Val	Ala	Ser	Val	Arg	Asp	Leu	Asp	
65					70				75				80		

Leu	His	Asp	Phe	Glu	Ala	Asn	Ile	Val	Gly	Phe	Ala	Gly	Glu	Ser	Ala
			85						90					95	

Lys	Gly	Ser	Val	Ala	Val	Asn	Xaa	Xaa	Glu	Val	Thr	Ile	Ser	Ile	Ser
			100					105					110		

Trp	Asp	Xaa	Ser	Arg	Gly	Ala	Asn	Ala	Gln	Gly	Thr	Arg	Glu	Thr	Phe
		115					120					125			

Thr	Leu	Thr	Ser	Arg	Val	Ala	Val	Asp	Pro	Arg	Val	Leu	Pro	Xaa	
	130						135				140				

&lt;210&gt; 104

&lt;211&gt; 44

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

69

&lt;222&gt; (44)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 104

Met Ala Phe Phe Phe Ala Leu Phe Val Ile Phe Phe Val Ile Val Val  
 1 5 10 15

Gln Met Glu Ser His Ser Gly Leu Gly Lys Lys Ser Lys Ile Leu Ser  
 20 25 30

Gly Gly Gln Gly Glu Glu Val Tyr Phe Leu Asp Xaa  
 35 40

&lt;210&gt; 105

&lt;211&gt; 63

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (63)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 105

Met Tyr Phe Tyr Leu Ala Val Lys Pro Pro Leu Leu Trp Ala Arg Pro  
 1 5 10 15

Gln Val Ser Cys Arg Leu Ser Val Ser Leu Ala Trp Ser Tyr His Leu  
 20 25 30

His Leu Trp Ala Leu Phe Leu Phe Ser Ile Leu Leu Gln Cys Arg Ala  
 35 40 45

Arg Phe Leu Leu Leu Leu Val Leu Ser Gln Thr Gln Asp Leu Xaa  
 50 55 60

&lt;210&gt; 106

&lt;211&gt; 283

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (283)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 106

Met Gly Ser Pro Gly Met Val Leu Gly Leu Leu Val Gln Ile Trp Ala  
 1 5 10 15

Leu Gln Glu Ala Ser Ser Leu Ser Val Gln Gln Gly Pro Asn Leu Leu  
 20 25 30

Gln Val Arg Gln Gly Ser Gln Ala Thr Leu Val Cys Gln Val Asp Gln  
 35 40 45

Ala Thr Ala Trp Glu Arg Leu Arg Val Lys Trp Thr Lys Asp Gly Ala

```

              50              55              70              60
Ile Leu Cys Gln Pro Tyr Ile Thr Asn Gly Ser Leu Ser Leu Gly Val
65              70              75              80
Cys Gly Pro Gln Gly Arg Leu Ser Trp Gln Ala Pro Ser His Leu Thr
85              90              95
Leu Gln Leu Asp Pro Val Ser Leu Asn His Ser Gly Ala Tyr Val Cys
100              105              110
Trp Ala Ala Val Glu Ile Pro Glu Leu Glu Glu Ala Glu Gly Asn Ile
115              120              125
Thr Arg Leu Phe Val Asp Pro Asp Asp Pro Thr Gln Asn Arg Asn Arg
130              135              140
Ile Ala Ser Phe Pro Gly Phe Leu Phe Val Leu Leu Gly Val Gly Ser
145              150              155              160
Met Gly Val Ala Ala Ile Val Trp Gly Ala Trp Phe Trp Gly Arg Arg
165              170              175
Ser Cys Gln Gln Arg Asp Ser Gly Asn Ser Pro Gly Asn Ala Phe Tyr
180              185              190
Ser Asn Val Leu Tyr Arg Pro Arg Gly Ala Pro Lys Lys Ser Glu Asp
195              200              205
Cys Ser Gly Glu Gly Lys Asp Gln Arg Gly Gln Ser Ile Tyr Ser Thr
210              215              220
Ser Phe Pro Gln Pro Ala Pro Arg Gln Pro His Leu Ala Ser Arg Pro
225              230              235              240
Cys Pro Ser Pro Arg Pro Cys Pro Ser Pro Arg Pro Gly His Pro Val
245              250              255
Ser Met Val Arg Val Ser Pro Arg Pro Ser Pro Thr Gln Gln Pro Arg
260              265              270
Pro Lys Gly Phe Pro Lys Val Gly Glu Glu Xaa
275              280

```

&lt;210&gt; 107

&lt;211&gt; 98

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (98)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 107

```

Met Cys Lys Leu Cys Phe Tyr Leu Tyr Leu Cys Thr Trp Phe Pro Phe
1              5              10              15

```

Gly Ala Ser Gly Leu Phe Trp Asp Lys Trp Cys Leu Pro Arg His Leu  
 20 25 30  
 Pro Val Val Ser Gly Gln Glu Gln Leu Ser Ser Ser Leu Pro Ala Ala  
 35 40 45  
 Leu Leu Phe Leu Gly Arg Arg Trp Arg Pro Pro Leu Arg Val Ser Pro  
 50 55 60  
 Gly Leu Ser Phe Arg Gly Gly Arg Ala Gly Glu Pro Gln Gly Trp Gly  
 65 70 75 80  
 Asp Ser Trp Glu Met Glu Val Ala Pro Ala Pro Leu Asp Gln Tyr Trp  
 85 90 95  
 Leu Xaa

<210> 108  
 <211> 62  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> (62)  
 <223> Xaa equals stop translation

<400> 108  
 Met Cys Leu Leu Leu Trp Leu Thr Thr Phe Gln Arg Thr Ser Gly  
 1 5 10 15  
 Ala Leu Arg Arg Gly Gly Leu Ser Ser Pro Ala Trp Ala Met Arg Ser  
 20 25 30  
 Pro Ser Val Tyr Ser Thr Gln Thr Pro Ser Pro Met Met Ser Thr Gly  
 35 40 45  
 Thr Leu Arg Gly Leu Ser Gly Ala Met Cys Asn Leu Ser Xaa  
 50 55 60

<210> 109  
 <211> 47  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> (47)  
 <223> Xaa equals stop translation

<400> 109  
 Met Lys Leu Cys Lys Leu Thr Gln Cys Ser Phe Leu Leu Lys Ser Leu  
 1 5 10 15  
 Ile Leu Leu Leu Glu Gln Leu Asn Val Ser Met Gly Phe Val Ala Ala  
 20 25 30

72

Phe Asp Val Leu Val Gly Cys Ser Ile Cys Phe Glu Lys His Xaa  
 35 40 45

&lt;210&gt; 110

&lt;211&gt; 47

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (47)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 110

Met Thr Thr Phe Ser Leu Cys Ser Gln Leu Ala Leu Leu Cys Ala Cys  
 1 5 10 15

Thr Ser Leu Val Ser Leu Pro Pro Phe Val Asp Tyr Lys Asp Thr Ser  
 20 25 30

Pro Val Gly Pro Glu Pro His Cys Lys Gly Leu Ile Leu Thr Xaa  
 35 40 45

&lt;210&gt; 111

&lt;211&gt; 42

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (32)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (33)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (41)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (42)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 111

Met Asn Ile Leu Val Cys Val Phe Trp Leu Trp Gly Gly Val Ala Gly  
 1 5 10 15

Ser Trp Gly Arg His Ile Phe Ile Phe Thr Ser Val Lys Asn Val Xaa  
 20 25 30

Xaa Ala Ser His Cys Ala Trp Pro Xaa Xaa

35 40 73

<210> 112  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens

<400> 112  
 Met Gly Gly Ile Ala Leu Pro Ser Leu Ser Leu Cys Leu Leu Ser Ala  
 1 5 10 15  
 Gly Ser His Cys Ile Ser Pro Ala Asp Gln Glu Thr Gly Pro Lys Val  
 20 25 30  
 Thr Ala Pro Gln Gly Asn Phe Leu Pro  
 35 40

<210> 113  
 <211> 44  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (44)  
 <223> Xaa equals stop translation

<400> 113  
 Met Ile Val Leu Lys Trp Ile Phe Leu Ala Cys Val His Glu Cys Met  
 1 5 10 15  
 Cys Lys Pro Leu Lys Cys Phe Leu Glu Lys Ile Leu Glu Val Leu Ile  
 20 25 30  
 Met Val Lys Leu Lys Met Gly Val Leu Pro Ala Xaa  
 35 40

<210> 114  
 <211> 182  
 <212> PRT  
 <213> Homo sapiens

<400> 114  
 Met Glu Pro Gln Leu Gly Pro Glu Ala Ala Ala Leu Arg Pro Gly Trp  
 1 5 10 15  
 Leu Ala Leu Leu Leu Trp Val Ser Ala Leu Ser Cys Ser Phe Ser Leu  
 20 25 30  
 Pro Ala Ser Ser Leu Ser Ser Leu Val Pro Gln Val Arg Thr Ser Tyr  
 35 40 45  
 Asn Phe Gly Arg Thr Phe Leu Gly Leu Asp Lys Cys Asn Ala Cys Ile  
 50 55 60  
 Gly Thr Ser Ile Cys Lys Lys Phe Phe Lys Glu Glu Ile Arg Ser Asp



```

<400> 115
Met Ala Leu Gly Ser Met Tyr Leu Val Leu Thr Leu Ile Val Ala Lys
  1             5             10             15

Val Leu Arg Gly Ala Glu Pro Cys Cys Gly Pro Leu Lys Asn Arg Val
          20             25             30

Leu Arg Pro Cys Pro Leu Pro Val His Cys Pro Leu Pro Ile Pro Ser
      35             40             45

Pro Ala Glu Gly Ile Pro Trp Val Ala Tyr Leu Pro Ile Arg Trp Phe
  50             55             60

Ile Ser Cys Cys Pro Gly His Cys Ile Gln Ile Pro Met Cys Thr Ser
  65             70             75             80

Xaa

```

```
<210> 116
<211> 49
<212> PRT
<213> Homo sapiens
```

75.

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (49)

&lt;223&gt; Xaa, equals stop translation

&lt;400&gt; 116

Met Ser Cys Glu Asn Asn Leu Lys Lys Lys Asn Thr Thr Leu Leu Ser  
 1 5 10 15

Tyr Leu Ile Phe Leu Ala Leu Val Met Tyr Leu Thr Phe Met Phe Leu  
 20 25 30

Ser Ser Val Ser Thr Ser Arg Ile Ser Leu Ser Asn Ser Met Ile Ile  
 35 40 45

Xaa

&lt;210&gt; 117

&lt;211&gt; 204

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (31)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (93)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (99)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (115)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (151)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (204)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 117

Met Val Gly Leu Met His Ile Gly Phe Gly Ile Val Leu Cys Leu Ile  
 1 5 10 15

Ser Phe Ser Phe Arg Glu Val Leu Gly Phe Ala Ser Thr Ala Xaa Ile

```
<210> 119
<211> 13
<212> PRT
<213> Homo sapiens
```

77

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (13)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 119

Leu Val Leu Trp Ile Val Met Leu Thr Tyr Ala Thr Xaa  
 1 5 10

&lt;210&gt; 120

&lt;211&gt; 80

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (80)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 120

Met Glu Pro Gln Leu Gly Pro Glu Ala Ala Ala Leu Arg Pro Gly Trp  
 1 5 10 15

Leu Ala Leu Leu Leu Trp Val Ser Ala Leu Ser Cys Ser Phe Ser Leu  
 20 25 30

Pro Ala Ser Ser Leu Ser Ser Leu Val Pro Gln Val Arg Thr Ser Tyr  
 35 40 45

Asn Phe Gly Arg Thr Phe Leu Gly Leu Asp Lys Cys Asn Ala Cys Ile  
 50 55 60

Gly Thr Ser Ile Cys Lys Lys Phe Phe Lys Glu Arg Asn Lys Ile Xaa  
 65 70 75 80

&lt;210&gt; 121

&lt;211&gt; 146

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (96)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (107)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (111)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

78

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (115)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (122)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (132)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 121

Met	Glu	Pro	Gln	Leu	Gly	Pro	Glu	Ala	Ala	Ala	Leu	Arg	Pro	Gly	Trp
1				5						10				15	

Leu	Ala	Leu	Leu	Leu	Trp	Val	Ser	Ala	Leu	Ser	Cys	Ser	Phe	Ser	Leu
		20						25					30		

Pro	Ala	Ser	Ser	Leu	Ser	Ser	Leu	Val	Pro	Gln	Val	Arg	Thr	Ser	Tyr
		35					40					45			

Asn	Phe	Gly	Arg	Thr	Phe	Leu	Gly	Leu	Asp	Lys	Cys	Asn	Ala	Cys	Ile
	50					55					60				

Gly	Thr	Ser	Ile	Cys	Lys	Lys	Phe	Phe	Lys	Glu	Glu	Ile	Arg	Ser	Asp
65					70					75					80

Asn	Trp	Leu	Ala	Ser	His	Leu	Gly	Thr	Ala	Ser	Arg	Phe	Pro	Leu	Xaa
				85					90					95	

Ser	Tyr	Pro	Cys	Lys	Leu	Leu	Gln	Met	Ile	Xaa	Lys	Ile	Trp	Xaa	Pro
		100						105					110		

Cys	Gly	Xaa	Leu	Leu	Thr	Gly	Gln	Gln	Xaa	Ser	Asn	Glu	Ile	Ser	Lys
		115					120					125			

Gln	Glu	Ile	Xaa	Cys	Leu	Leu	His	Pro	Pro	Pro	Lys	Asn	Leu	His	Ile
	130						135					140			

Asp Val

145

&lt;210&gt; 122

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (81)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 122

79  
 Met Ala Leu Gly Ser Met Tyr Leu Val Leu Thr Leu Ile Val Ala Lys  
 1 5 10 15  
 Val Leu Arg Gly Ala Glu Pro Cys Cys Gly Pro Leu Lys Asn Arg Val  
 20 25 30  
 Leu Arg Pro Cys Pro Leu Pro Val His Cys Pro Leu Pro Ile Pro Ser  
 35 40 45  
 Pro Ala Glu Gly Ile Pro Trp Val Ala Tyr Leu Pro Ile Arg Trp Phe  
 50 55 60  
 Ile Ser Cys Cys Pro Gly His Cys Ile Gln Ile Pro Met Cys Thr Ser  
 65 70 75 80

Xaa

<210> 123  
 <211> 337  
 <212> PRT  
 <213> Homo sapiens

<400> 123  
 Glu Pro His Arg Gly Pro His Leu Pro Pro Asp Leu Gly His His His  
 1 5 10 15  
 Gly Gln Arg Pro Gly Leu Gln Asn Ile Asn Val Phe Leu Arg Asn Thr  
 20 25 30  
 Val Lys Val Thr Gly Val Val Val Phe Met Phe Ser Leu Ser Trp Gln  
 35 40 45  
 Leu Ser Leu Val Thr Phe Met Gly Phe Pro Ile Ile Met Met Val Ser  
 50 55 60  
 Asn Ile Tyr Gly Lys Tyr Tyr Lys Arg Leu Ser Lys Glu Val Gln Asn  
 65 70 75 80  
 Ala Leu Ala Arg Ala Ser Asn Thr Ala Glu Glu Thr Ile Ser Ala Met  
 85 90 95  
 Lys Thr Val Arg Ser Phe Ala Asn Glu Glu Glu Glu Ala Glu Val Tyr  
 100 105 110  
 Leu Arg Lys Leu Gln Gln Val Tyr Lys Leu Asn Arg Lys Glu Ala Ala  
 115 120 125  
 Ala Tyr Met Tyr Tyr Val Trp Gly Ser Gly Leu Thr Leu Leu Val Val  
 130 135 140  
 Gln Val Ser Ile Leu Tyr Tyr Gly Gly His Leu Val Ile Ser Gly Gln  
 145 150 155 160  
 Met Thr Ser Gly Asn Leu Ile Ala Phe Ile Ile Tyr Glu Phe Val Leu  
 165 170 175  
 Gly Asp Cys Met Glu Asn Val Ser Phe Ser Leu Ser Pro Gly Lys Val

180 80 190  
 Thr Ala Leu Val Gly Pro Ser Gly Ser Gly Lys Ser Ser Cys Val Asn  
 195 200 205  
 Ile Leu Glu Asn Phe Tyr Pro Leu Glu Gly Gly Arg Val Leu Leu Asp  
 210 215 220  
 Gly Lys Pro Ile Ser Ala Tyr Asp His Lys Tyr Leu His Arg Val Ile  
 225 230 235 240  
 Ser Leu Val Ser Gln Glu Pro Val Leu Phe Ala Arg Ser Ile Thr Asp  
 245 250 255  
 Asn Ile Ser Tyr Gly Leu Pro Thr Val Pro Phe Glu Met Val Val Glu  
 260 265 270  
 Ala Ala Gln Lys Ala Asn Ala His Gly Phe Ile Met Glu Leu Gln Asp  
 275 280 285  
 Gly Tyr Ser Thr Glu Thr Gly Glu Lys Gly Ala Gln Leu Ser Gly Gly  
 290 295 300  
 Gln Lys Gln Arg Val Ala Trp Pro Gly Leu Trp Cys Gly Thr Pro Gln  
 305 310 315 320  
 Ser Ser Ser Trp Met Lys Pro Pro Ala Leu Trp Met Pro Arg Ala Ser  
 325 330 335  
 Ile

<210> 124  
 <211> 315  
 <212> PRT  
 <213> Homo sapiens

<400> 124  
 Met Ser Ser Ala Thr Trp Thr Ala Ala Ser Trp Arg Thr Ser Ala Thr  
 1 5 10 15  
 Ser Thr Ser Leu Thr Arg Cys Trp Ile Ser Gly Gln Pro Ala Cys Thr  
 20 25 30  
 Ala Ala Ala Cys Cys Trp Gly Ala Thr Ile Gly Val Ala Lys Asn Ser  
 35 40 45  
 Ala Leu Gly Pro Arg Arg Leu Arg Ala Ser Trp Leu Val Ile Thr Leu  
 50 55 60  
 Val Cys Leu Phe Val Gly Ile Tyr Ala Met Val Lys Leu Leu Leu Phe  
 65 70 75 80  
 Ser Glu Val Arg Arg Pro Ile Arg Asp Pro Trp Phe Trp Ala Leu Phe  
 85 90 95  
 Val Trp Thr Tyr Ile Ser Leu Gly Ala Ser Phe Leu Leu Trp Trp Leu  
 100 105 110

81

Leu Ser Thr Val Arg Pro Gly Thr Gln Ala Leu Glu Pro Gly Ala Ala  
 115 120 125  
 Thr Glu Ala Glu Gly Phe Pro Gly Ser Gly Arg Pro Pro Pro Glu Gln  
 130 135 140  
 Ala Ser Gly Ala Thr Leu Gln Lys Leu Leu Ser Tyr Thr Lys Pro Asp  
 145 150 155 160  
 Val Ala Phe Leu Val Ala Ala Ser Phe Phe Leu Ile Val Ala Ala Leu  
 165 170 175  
 Gly Glu Thr Phe Leu Pro Tyr Tyr Thr Gly Arg Ala Ile Asp Gly Ile  
 180 185 190  
 Val Ile Gln Lys Ser Met Asp Gln Phe Ser Thr Ala Val Val Ile Val  
 195 200 205  
 Cys Leu Leu Ala Ile Gly Ser Ser Phe Ala Ala Gly Ile Arg Gly Gly  
 210 215 220  
 Ile Phe Thr Leu Ile Phe Ala Arg Leu Asn Ile Arg Leu Arg Asn Cys  
 225 230 235 240  
 Leu Phe Arg Ser Leu Val Ser Gln Glu Thr Ser Phe Phe Asp Glu Asn  
 245 250 255  
 Arg Thr Gly Asp Leu Ile Ser Arg Leu Thr Ser Asp Thr Thr Met Val  
 260 265 270  
 Ser Asp Leu Val Ser Arg Thr Ser Met Ser Ser Cys Gly Thr Gln Ser  
 275 280 285  
 Arg Ser Arg Ala Trp Trp Ser Ser Cys Ser Ala Ser His Gly Ser Ser  
 290 295 300  
 Pro Trp Ser Pro Ser Trp Ala Ser Pro Ser Ser  
 305 310 315

&lt;210&gt; 125

&lt;211&gt; 167

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 125

His Leu Leu Arg Pro Ala His Cys Ala Phe Arg Asp Gly Gly Gly Gly  
 1 5 10 15  
 Arg Thr Glu Gly Gln Cys Pro Arg Leu His His Gly Thr Pro Gly Arg  
 20 25 30  
 Leu Gln His Arg Asp Arg Gly Glu Gly Arg Pro Ala Val Arg Trp Pro  
 35 40 45  
 Glu Ala Ala Gly Gly Met Ala Arg Ala Leu Val Arg Asn Pro Pro Val  
 50 55 60



Leu Ile Leu Asp Glu Ala Thr Ser Ala Leu Asp Ala Glu Ser Glu Tyr  
 65 70 75 80  
 Leu Ile Gln Gln Ala Ile His Gly Asn Leu Gln Lys His Thr Val Leu  
 85 90 95  
 Ile Ile Ala His Arg Leu Ser Thr Val Glu His Ala His Leu Ile Val  
 100 105 110  
 Val Leu Asp Lys Gly Arg Val Val Gln Gln Gly Thr His Gln Gln Leu  
 115 120 125  
 Leu Ala Gln Gly Gly Leu Tyr Ala Lys Leu Val Gln Arg Gln Met Leu  
 130 135 140  
 Gly Leu Gln Pro Ala Ala Asp Phe Thr Ala Gly His Asn Glu Pro Val  
 145 150 155 160  
 Ala Asn Gly Ser His Lys Ala  
 165

&lt;210&gt; 126

&lt;211&gt; 227

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (71)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 126

Arg Leu Thr Lys Thr Ile Ser Phe Ser Leu Gln Asn Gln Thr Ala Phe  
 1 5 10 15  
 Ile Asn Ser Leu Ala Lys Thr Pro Tyr Gln Ala Leu Thr Gly Ala Ala  
 20 25 30  
 Leu Ala Gly Ser Tyr Pro Ile Trp Glu Asn Glu Asn Thr Leu Ser Trp  
 35 40 45  
 Tyr Leu Pro Ser Pro Thr Thr Leu Leu Ser Pro Pro Val Leu Phe Cys  
 50 55 60  
 Val Ile Gln Leu Ile Phe Xaa Leu Pro Ala Asn Trp Ser Gly Thr Cys  
 65 70 75 80  
 Thr Leu Val Phe Gln Ala Pro Thr Ile Asn Ile Leu Pro Pro Asn Gln  
 85 90 95  
 Thr Ile Leu Ile Ser Val Glu Ala Ser Ile Ser Ser Ser Pro Ile Arg  
 100 105 110  
 Asn Lys Trp Ala Leu His Leu Ile Thr Leu Leu Thr Gly Leu Gly Ile  
 115 120 125  
 Thr Ala Ala Leu Gly Thr Gly Ile Ala Gly Ile Thr Thr Ser Ile Thr  
 130 135 140

83

Ser Tyr Gln Thr Leu Phe Thr Thr Leu Ser Asn Thr Val Glu Asp Met  
 145 150 155 160

His Thr Ser Ile Thr Ser Leu Gln Arg Gln Leu Asp Phe Leu Val Gly  
 165 170 175

Val Ile Leu Gln Asn Trp Arg Val Leu Asp Leu Leu Thr Thr Glu Lys  
 180 185 190

Gly Gly Thr Cys Ile Tyr Leu Gln Glu Glu Cys Cys Phe Cys Val Asn  
 195 200 205

Glu Ser Gly Ile Val His Ile Ala Val Arg Arg Leu His Asp Arg Ala  
 210 215 220

Ala Glu Leu  
 225

<210> 127  
 <211> 29  
 <212> PRT  
 <213> Homo sapiens

<400> 127  
 Tyr Pro Ile Trp Glu Asn Glu Asn Thr Leu Ser Trp Tyr Leu Pro Ser  
 1 5 10 15

Pro Thr Thr Leu Leu Ser Pro Pro Val Leu Phe Cys Val  
 20 25

<210> 128  
 <211> 27  
 <212> PRT  
 <213> Homo sapiens

<400> 128  
 Arg Val Leu Asp Leu Leu Thr Thr Glu Lys Gly Gly Thr Cys Ile Tyr  
 1 5 10 15

Leu Gln Glu Glu Cys Cys Phe Cys Val Asn Glu  
 20 25

<210> 129  
 <211> 10  
 <212> PRT  
 <213> Homo sapiens

<400> 129  
 Phe Ser Leu Gly Arg Arg His Cys Leu Gly  
 1 5 10

<210> 130  
 <211> 123  
 <212> PRT

84

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (64)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (83)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 130

Glu	His	Pro	Thr	Ala	Asp	Arg	Ala	Gly	Cys	Ser	Ala	Ser	Gly	Ala	Cys
1				5				10					15		

Tyr	Ser	Leu	His	His	Ala	Thr	Met	Lys	Arg	Gln	Ala	Ala	Glu	Glu	Ala
			20					25					30		

Cys	Ile	Leu	Arg	Gly	Gly	Ala	Leu	Ser	Thr	Val	Arg	Ala	Gly	Ala	Glu
		35					40					45			

Leu	Arg	Ala	Val	Leu	Ala	Leu	Leu	Arg	Ala	Gly	Pro	Gly	Pro	Gly	Xaa
	50					55					60				

Gly	Ser	Lys	Asp	Leu	Leu	Phe	Trp	Val	Ala	Leu	Glu	Arg	Arg	Arg	Ser
65					70					75					80

His	Cys	Xaa	Leu	Glu	Asn	Glu	Pro	Leu	Arg	Gly	Phe	Ser	Trp	Leu	Ser
			85						90					95	

Ser	Asp	Pro	Gly	Gly	Leu	Glu	Ser	Asp	Thr	Leu	Gln	Trp	Val	Glu	Glu
		100						105					110		

Pro	Gln	Arg	Ser	Cys	Thr	Ala	Arg	Arg	Trp	Val
	115						120			

&lt;210&gt; 131

&lt;211&gt; 344

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (19)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 131

Ser	Arg	Pro	Pro	Val	Gly	Ser	Ser	Pro	Gln	Leu	Glu	Gly	Asp	Ala	Met
1				5				10					15		

Pro	Pro	Xaa	Arg	Gln	Arg	Tyr	Leu	Cys	Lys	Tyr	Gln	Phe	Glu	Val	Leu
			20					25					30		

Cys	Pro	Ala	Pro	Arg	Pro	Gly	Ala	Ala	Ser	Asn	Leu	Ser	Tyr	Arg	Ala
		35				40						45			

Pro	Phe	Gln	Leu	His	Ser	Ala	Ala	Leu	Asp	Phe	Ser	Pro	Pro	Gly	Thr
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

50                      55                      85                      60  
 Glu Val Ser Ala Leu Cys Arg Gly Gln Leu Pro Ile Ser Val Thr Cys  
 65                      70                      75                      80  
 Ile Ala Asp Glu Ile Gly Ala Arg Trp Asp Lys Leu Ser Gly Asp Val  
 85                      90                      95  
 Leu Cys Pro Cys Pro Gly Arg Tyr Leu Arg Ala Gly Lys Cys Ala Glu  
 100                      105                      110  
 Leu Pro Asn Cys Leu Asp Asp Leu Gly Gly Phe Ala Cys Glu Cys Ala  
 115                      120                      125  
 Thr Gly Phe Glu Leu Gly Lys Asp Gly Arg Ser Cys Val Thr Ser Gly  
 130                      135                      140  
 Glu Gly Gln Pro Thr Leu Gly Gly Thr Gly Val Pro Thr Arg Arg Pro  
 145                      150                      155                      160  
 Pro Ala Thr Ala Thr Ser Pro Val Pro Gln Arg Thr Trp Pro Ile Arg  
 165                      170                      175  
 Val Asp Glu Lys Leu Gly Glu Thr Pro Leu Val Pro Glu Gln Asp Asn  
 180                      185                      190  
 Ser Val Thr Ser Ile Pro Glu Ile Pro Arg Trp Gly Ser Gln Ser Thr  
 195                      200                      205  
 Met Ser Thr Leu Gln Met Ser Leu Gln Ala Glu Ser Lys Ala Thr Ile  
 210                      215                      220  
 Thr Pro Ser Gly Ser Val Ile Ser Lys Phe Asn Ser Thr Thr Ser Ser  
 225                      230                      235                      240  
 Ala Thr Pro Gln Ala Phe Asp Ser Ser Ser Ala Val Val Phe Ile Phe  
 245                      250                      255  
 Val Ser Thr Ala Val Val Val Leu Val Ile Leu Thr Met Thr Val Leu  
 260                      265                      270  
 Gly Leu Val Lys Leu Cys Phe His Glu Ser Pro Ser Ser Gln Pro Arg  
 275                      280                      285  
 Lys Glu Ser Met Gly Pro Pro Gly Trp Arg Val Ile Leu Lys Pro Ala  
 290                      295                      300  
 Ala Leu Gly Ser Ser Ser Ala His Cys Thr Asn Asn Gly Val Lys Val  
 305                      310                      315                      320  
 Gly Asp Cys Asp Leu Arg Asp Arg Ala Glu Gly Ala Leu Leu Ala Glu  
 325                      330                      335  
 Ser Pro Leu Gly Ser Ser Asp Ala  
 340

&lt;210&gt; 132

&lt;211&gt; 6

<212> PRT  
<213> Homo sapiens

<400> 132  
Val Glu Gly Gly His Ser  
1 5

<210> 133  
<211> 7  
<212> PRT  
<213> Homo sapiens

<400> 133  
Arg Tyr Leu Thr Leu Thr His  
1 5

<210> 134  
<211> 6  
<212> PRT  
<213> Homo sapiens

<400> 134  
Cys Asn Thr Pro Trp Ala  
1 5

<210> 135  
<211> 8  
<212> PRT  
<213> Homo sapiens

<400> 135  
Ala Pro Val Ile Phe Ser His Ser  
1 5

<210> 136  
<211> 6  
<212> PRT  
<213> Homo sapiens

<400> 136  
Arg Asn Val Pro Asp Asp  
1 5

<210> 137  
<211> 6  
<212> PRT  
<213> Homo sapiens

<400> 137  
Gly Leu Glu Asp Val Ser  
1 5

<210> 138

87.

&lt;211&gt; 23

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 138

Val Glu Gly Gly His Ser Leu Asp Asn Ser Leu Ser Ile Leu Arg Thr  
1 5 10 15

Phe Tyr Met Leu Gly Val Arg  
20

&lt;210&gt; 139

&lt;211&gt; 190

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 139

Thr Trp Leu Arg Leu Gly Ser Ser Gln Ile Trp Leu Gly Thr Ala Pro  
1 5 10 15

Arg Gly Pro Arg Ile His Pro Glu Gln Ala Gly Leu Ala Gly Ala Pro  
20 25 30

Val Lys Ser Thr Ser Ser Glu Glu Ser Gln Pro Gly Gly Gln Cys Gln  
35 40 45

Ser Ser Gly Gly Ala Gln Thr Leu Pro Ser Leu Arg Ala Ala Pro Val  
50 55 60

Ala Ala Leu Gly Ser Leu Ser Ser Tyr Pro Asp Ser Cys Pro Arg Ala  
65 70 75 80

Thr Thr Pro Glu Leu Cys Pro Gly Ala Pro Thr Leu His Leu Ala Asp  
85 90 95

Ser Ile Ser Gly Pro Val Ser Pro Pro Gly Ser Ser Leu Gly Pro Asp  
100 105 110

Ala Trp Thr Leu Cys Ala Lys His His Gln Ala Lys Gly Met Thr Leu  
115 120 125

Gly Thr Pro Lys Val Leu Arg Leu Gln Pro Val Ser Pro Cys Trp Gly  
130 135 140

Pro Lys Ser Trp Arg Val Pro Gly Pro Phe Gln Pro Gly Arg Arg Arg  
145 150 155 160

Gly Glu Ser Arg Gln Gln Gly Arg Gly Lys Arg Arg Ser Ala Arg Ser  
165 170 175

Ala Gln Ser Pro Thr Gly Pro Glu Ser Ala Ala Trp Pro Cys  
180 185 190

&lt;210&gt; 140

&lt;211&gt; 129

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

88

&lt;400&gt; 140

Thr Val Ala Thr Ala Cys Val Trp Ala Ala Cys Thr Gly Cys Trp Ala  
 1 5 10 15

Arg Pro Pro Val Pro Thr Trp Ala Gly Cys Ala Ala Arg Cys Ala Ala  
 20 25 30

Glu Asp Ala Arg Ala Gly Val Gly Asp Leu Pro Ala Thr Gly Gly Ala  
 35 40 45

Ala Thr Gly Arg Arg Ala Leu Thr Pro Ala Pro Pro Arg Gly Pro Cys  
 50 55 60

Ile Leu Ser Pro Gln Pro Trp Ala Leu Gly Leu Pro Gly Ala Pro Leu  
 65 70 75 80

Pro Ala Ala Leu Pro Gly Arg Ala Arg Gly Arg Pro Gly Leu Pro Ala  
 85 90 95

Leu Pro Ala Leu Ser Thr Leu Pro Gly Cys Pro Ala Leu Asp Pro Ala  
 100 105 110

Gly Ala Gly Thr Leu Cys Pro Pro Pro Gly Ala Ala Glu Pro Ala Gly  
 115 120 125

Pro

&lt;210&gt; 141

&lt;211&gt; 90

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 141

Arg Ser Gly Gln Pro Gly Glu Gly Ser Met Leu Arg Lys Phe Ser Leu  
 1 5 10 15

Gln Arg Leu Leu Ser Pro Leu Asp Gln Ala Gln Thr Arg Trp Gly Leu  
 20 25 30

Ala Leu Ala Cys Val Ala Gly Asp Lys Gly Pro Pro Arg Pro Trp Asn  
 35 40 45

Ile Ser Ser Ala Pro Ala His Pro His Val Thr Thr Pro Gly Met Glu  
 50 55 60

Thr Ser Gly Gly Pro Ala Arg Asp Gly Gly Leu Ile Leu Glu Arg Glu  
 65 70 75 80

Ala Ala Phe Asn Lys Pro Ala Pro Gly Glu  
 85 90

&lt;210&gt; 142

&lt;211&gt; 307

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

<220>  
 <221> SITE  
 <222> (135)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (197)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (203)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (219)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (255)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 142  
 Arg Cys Gln Arg Asn Lys Asp Ile Met Met Ser Ser Lys Pro Thr Ser  
           1                  5                  10                  15  
 His Ala Glu Val Asn Glu Thr Ile Pro Asn Pro Tyr Pro Pro Ser Ser  
                   20                  25                  30  
 Phe Met Ala Pro Gly Phe Gln Gln Pro Leu Gly Ser Ile Asn Leu Glu  
           35                  40                  45  
 Asn Gln Ala Gln Gly Ala Gln Arg Ala Gln Pro Tyr Gly Ile Thr Ser  
           50                  55                  60  
 Pro Gly Ile Phe Ala Ser Ser Gln Pro Gly Gln Gly Asn Ile Gln Met  
           65                  70                  75                  80  
 Ile Asn Pro Ser Val Gly Thr Ala Val Met Asn Phe Lys Glu Glu Ala  
                   85                  90                  95  
 Lys Ala Leu Gly Val Ile Gln Ile Met Val Gly Leu Met His Ile Gly  
           100                  105                  110  
 Phe Gly Ile Val Leu Cys Leu Ile Ser Phe Ser Phe Arg Glu Val Leu  
           115                  120                  125  
 Gly Phe Ala Ser Thr Ala Xaa Ile Gly Gly Tyr Pro Phe Trp Gly Gly  
           130                  135                  140  
 Leu Ser Phe Ile Ile Ser Gly Ser Leu Ser Val Ser Ala Ser Lys Glu  
           145                  150                  155                  160  
 Leu Ser Arg Cys Leu Val Lys Gly Ser Leu Gly Met Asn Ile Gly Arg  
           165                  170                  175



90

Ser Ile Leu Ala Phe Ile Gly Val Ile Leu Leu Leu Val Asp Met Cys  
 180 185 190

Ile Asn Gly Val Xaa Gly Gln Asp Tyr Trp Xaa Val Leu Ser Gly Lys  
 195 200 205

Gly Ile Ser Ala Thr Leu Met Ile Phe Ser Xaa Leu Glu Phe Phe Val  
 210 215 220

Ala Cys Ala Thr Ala His Phe Ala Asn Gln Ala Asn Thr Thr Thr Asn  
 225 230 235 240

Met Ser Val Leu Val Ile Pro Asn Met Tyr Glu Ser Asn Pro Xaa Thr  
 245 250 255

Pro Ala Ser Ser Ser Ala Pro Pro Arg Cys Asn Asn Tyr Ser Ala Asn  
 260 265 270

Ala Pro Lys Arg Lys Arg Gly Ile Ser Leu Ile Ser Trp Arg Lys Thr  
 275 280 285

Thr Cys Lys Asn Phe Leu Arg Arg Cys Leu Leu Leu Ser Thr Met Ile  
 290 295 300

Ser Ser Leu  
 305

<210> 143  
 <211> 246  
 <212> PRT  
 <213> Homo sapiens

<400> 143  
 Met Gly Arg Leu Asp Gly Lys Val Ile Ile Leu Thr Ala Ala Ala Gln  
 1 5 10 15

Gly Ile Gly Gln Ala Ala Ala Leu Ala Phe Ala Arg Glu Gly Ala Lys  
 20 25 30

Val Ile Ala Thr Asp Ile Asn Glu Ser Lys Leu Gln Glu Leu Glu Lys  
 35 40 45

Tyr Pro Gly Ile Gln Thr Arg Val Leu Asp Val Thr Lys Lys Lys Gln  
 50 55 60

Ile Asp Gln Phe Ala Asn Glu Val Glu Arg Leu Asp Val Leu Phe Asn  
 65 70 75 80

Val Ala Gly Phe Val His His Gly Thr Val Leu Asp Cys Glu Glu Lys  
 85 90 95

Asp Trp Asp Phe Ser Met Asn Leu Asn Val Arg Asn Val Met Tyr Leu  
 100 105 110

Met Ile Lys Ala Phe Leu Pro Lys Met Leu Ala Gln Lys Ser Gly Asn  
 115 120 125

91

Ile Ile Asn Met Ser Ser Val Ala Ser Ser Val Lys Gly Val Val Asn  
 130 135 140

Arg Cys Val Tyr Ser Thr Thr Lys Ala Ala Val Ile Gly Leu Thr Lys  
 145 150 155 160

Ser Val Ala Ala Asp Phe Ile Gln Gln Gly Ile Arg Cys Asn Cys Val  
 165 170 175

Cys Pro Gly Thr Val Asp Thr Pro Ser Leu Gln Glu Arg Ile Gln Ala  
 180 185 190

Arg Gly Asn Pro Glu Glu Ala Arg Asn Asp Phe Leu Lys Arg Gln Lys  
 195 200 205

Thr Gly Arg Phe Ala Thr Ala Glu Glu Ile Ala Met Leu Cys Val Tyr  
 210 215 220

Leu Ala Ser Asp Glu Ser Ala Tyr Val Thr Gly Asn Pro Val Ile Ile  
 225 230 235 240

Asp Gly Gly Trp Ser Leu  
 245

<210> 144  
 <211> 234  
 <212> PRT  
 <213> Homo sapiens

<400> 144

Gly Thr Ile Gly Leu Tyr Trp Val Gly Ser Ile Ile Met Ser Val Val  
 1 5 10 15

Val Phe Val Pro Gly Asn Ile Val Gly Lys Tyr Gly Thr Arg Ile Cys  
 20 25 30

Pro Ala Phe Phe Leu Ser Ile Pro Tyr Thr Cys Leu Pro Val Trp Ala  
 35 40 45

Gly Phe Arg Ile Tyr Asn Gln Pro Ser Glu Asn Tyr Asn Tyr Pro Ser  
 50 55 60

Lys Val Ile Gln Glu Ala Gln Ala Lys Asp Leu Leu Arg Arg Pro Phe  
 65 70 75 80

Asp Leu Met Leu Val Val Cys Leu Leu Leu Ala Thr Gly Phe Cys Leu  
 85 90 95

Phe Arg Gly Leu Ile Ala Leu Asp Cys Pro Ser Glu Leu Cys Arg Leu  
 100 105 110

Tyr Thr Gln Phe Gln Glu Pro Tyr Leu Lys Asp Pro Ala Ala Tyr Pro  
 115 120 125

Lys Ile Gln Met Leu Ala Tyr Met Phe Tyr Ser Val Pro Tyr Phe Val  
 130 135 140

Thr Ala Leu Tyr Gly Leu Val Val Pro Gly Cys Ser Trp Met Pro Asp

```

<210> 145
<211> 238
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (184)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 145
Met Ser Asn His Asp Pro Arg Gly Cys Thr Arg Arg Arg Ala Gln Lys
  1              5              10              15

Pro Leu Ala Ile Gln Pro Arg Leu Phe His Ala Ser Ala Pro Asp Glu
      20              25              30

Gly Thr Gln Gly Thr Leu Lys Gly Thr Gln Lys Gly Gly Cys Ile Leu
      35              40              45

Val Gln Cys Gln Ser Glu Gly Gly Ala Ala Gly Ala Trp Thr Gly Pro
      50              55              60

Pro Ser Pro Ala Arg Asp Arg Arg Val Arg Pro Pro Gly Thr Lys Ala
      65              70              75              80

Gln Arg Leu Glu Arg Arg Arg His Val Pro Arg Leu His Gly Leu Gly
      85              90              95

Val Gly Gly Cys Glu Val Arg Thr Gly Ile Val Ala Arg Ile Ser Gly
      100             105             110

Ser Thr Pro Trp Ala Gly Gly Lys Pro Leu Gly Leu His Gly Ala Met
      115             120             125

Gly Glu Ala Gly Ala Gly Asp Thr Gly Cys Cys Ala Lys Gly Pro Ser
      130             135             140

Pro Ala Ala Pro Leu Pro Ala Glu Gly Arg Gly Gln Gly Ala Gly Pro
145             150             155             160

```

```

<400> 146
Ser Thr Xaa Thr Xaa Thr Ile Gly Xaa Ala Gly Thr Pro Ala Gly Thr
 1      5      10      15

Gly Pro Glu Phe Pro Gly Arg Pro Thr Arg Pro Gly Glu Xaa Pro Val
      20      25      30

Asp Phe Ser Lys Gln Tyr Ser Ala Ser Trp Met Cys Leu Ser Leu Leu
      35      40      45

Ala Ala Leu Ala Cys Ser Ala Gly Asp Thr Trp Ala Ser Glu Val Gly
 50      55      60

```

94  
 Pro Val Leu Ser Lys Ser Ser Pro Arg Leu Ile Thr Thr Trp Glu Lys  
 65 70 75 80  
 Val Pro Val Gly Thr Asn Gly Gly Val Thr Val Val Gly Leu Val Ser  
 85 90 95  
 Ser Leu Leu Gly Gly Thr Phe Val Gly Ile Ala Tyr Phe Leu Thr Gln  
 100 105 110  
 Leu Ile Phe Val Asn Asp Leu Asp Ile Ser Ala Pro Gln Trp Pro Ile  
 115 120 125  
 Ile Ala Phe Gly Gly Leu Ala Gly Leu Leu Gly Ser Ile Val Asp Ser  
 130 135 140  
 Tyr Leu Gly Ala Thr Met Gln Tyr Thr Gly Leu Asp Glu Ser Thr Gly  
 145 150 155 160  
 Met Val Val Asn Ser Pro Thr Asn Xaa Ala Arg His Ile Ala Gly Lys  
 165 170 175  
 Pro Ile Leu Asp Asn Asn Ala Val Asn Leu Phe Ser Ser Val Leu Ile  
 180 185 190  
 Ala Leu Leu Leu Pro Thr Ala Ala Trp Gly Phe Trp Pro Arg Gly  
 195 200 205

&lt;210&gt; 147

&lt;211&gt; 116

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 147

Met Ser Gln Arg Ala Gly Arg Arg Pro Gly Gly Trp Asn Pro Ser Leu  
 1 5 10 15  
 Ser Val Val Glu Val Cys Arg Gly Cys Arg Gly Thr Gly Pro Leu Pro  
 20 25 30  
 Trp Gly Ala Ser Leu Phe Pro Cys Ser Ala Ser Pro Leu Phe Pro Leu  
 35 40 45  
 Pro Leu Asn Arg Arg Gly Asp Val His Gly Thr Leu Gly Gly Arg Met  
 50 55 60  
 Leu Asn Arg Val Glu Cys Arg Asp Gly Val Ala Ala Trp Leu Cys  
 65 70 75 80  
 Leu His Asp Ala Ala Ala Ile Arg Gly Ala Val Gly Arg Cys Pro Met  
 85 90 95  
 Trp Thr Gln Pro Thr His Trp Val Leu Leu Leu Cys Trp Ala Leu His  
 100 105 110  
 Phe Tyr Cys Arg  
 115

95

<210> 148  
 <211> 81  
 <212> PRT  
 <213> Homo sapiens

<400> 148  
 Met Thr Ala His Ser Phe Ala Leu Pro Val Ile Ile Phe Thr Thr Phe  
           1                  5                  10                  15  
 Trp Gly Leu Val Gly Ile Ala Gly Pro Trp Phe Val Pro Lys Gly Pro  
                   20                  25                  30  
 Asn Arg Gly Val Ile Ile Thr Met Leu Val Ala Thr Ala Val Cys Cys  
           35                  40                  45  
 Tyr Leu Phe Trp Leu Ile Ala Ile Leu Ala Gln Leu Asn Pro Leu Phe  
           50                  55                  60  
 Gly Pro Gln Leu Lys Asn Glu Thr Ile Trp Tyr Val Arg Phe Leu Trp  
           65                  70                  75                  80  
 Glu

<210> 149  
 <211> 110  
 <212> PRT  
 <213> Homo sapiens

<400> 149  
 Ala Gln Arg Ala Ala Arg Leu Gly Thr Arg Ala Pro Ala Ala Pro Ala  
           1                  5                  10                  15  
 Ala Arg Pro Cys Ile Leu Pro Gly His Pro Ala Pro Gly His Asp Gly  
                   20                  25                  30  
 Ala Leu Ile Arg Pro Pro Gly His His Leu His His Val Leu Gly Pro  
           35                  40                  45  
 Arg Arg His Arg Gly Pro Trp Phe Val Pro Lys Gly Pro Asn Arg Gly  
           50                  55                  60  
 Val Ile Ile Thr Met Leu Val Ala Thr Ala Val Cys Cys Tyr Leu Phe  
           65                  70                  75                  80  
 Trp Leu Ile Ala Ile Leu Ala Gln Leu Asn Pro Leu Phe Gly Pro Gln  
                   85                  90                  95  
 Leu Lys Asn Glu Thr Ile Trp Tyr Val Arg Phe Leu Trp Glu  
                   100                  105                  110

<210> 150  
 <211> 135  
 <212> PRT  
 <213> Homo sapiens

<400> 150

Met Thr Leu Glu Glu His Arg Asp Arg Pro Arg Leu Gly Met Cys Met  
 1 5 10 15  
 Cys Val Cys Ala Cys Val Tyr Ala Cys Met Leu Met His Val Cys Val  
 20 25 30  
 His Ala Cys Leu Cys Val Cys Val Cys Val Cys Val Glu Pro Trp Ser  
 35 40 45  
 Ser Arg Gln Ser Lys Asp Thr Gly Gly Trp His Met Glu Glu Gln Val  
 50 55 60  
 Thr Pro Pro Ser Leu Ala Gln Leu Lys Ser Gly Gln Val Arg Gly Glu  
 65 70 75 80  
 Met Gly Glu Gly Arg Gly Glu Lys Gly Glu Glu Ala Leu Thr Gly Gly  
 85 90 95  
 Ala Glu Ala Leu Ser Leu Leu Gly Arg Arg Ser Pro Ser Thr Pro Leu  
 100 105 110  
 Phe Leu Asp Arg Glu Asp Lys Gln Ala Lys Asp Ala Arg Asn Leu Ser  
 115 120 125  
 Ser Thr Val Ala Pro Asp Phe  
 130 135

<210> 151  
 <211> 82  
 <212> PRT  
 <213> Homo sapiens

<400> 151  
 His Glu Lys Ile Leu Thr Pro Ile Trp Pro Ser Ser Thr Asp Leu Glu  
 1 5 10 15  
 Lys Pro His Glu Met Leu Phe Leu Asn Val Ile Leu Phe Ser Leu Thr  
 20 25 30  
 Val Phe Thr Leu Ile Ser Thr Ala His Thr Leu Asp Arg Ala Val Arg  
 35 40 45  
 Ser Asp Trp Leu Leu Leu Val Leu Ile Tyr Ala Cys Leu Glu Glu Leu  
 50 55 60  
 Ile Pro Glu Leu Ile Phe Asn Leu Tyr Cys Gln Gly Asn Ala Thr Leu  
 65 70 75 80  
 Phe Phe

<210> 152  
 <211> 71  
 <212> PRT  
 <213> Homo sapiens

<400> 152

97

Pro Ala Asn Lys Ala Gly Ala Ala Ile Glu Ala Gly Ile Gly Ile Ser  
 1 5 10 15

Leu Met Val Leu Ser Pro Trp Ala Cys Leu Phe Val Val Phe Phe Pro  
 20 25 30

Tyr Ile Gln Ser Ser Leu Arg Ser Asp Lys His Leu Gln Leu Ser Asn  
 35 40 45

Ile Leu Pro Thr Pro Ser His His Ile His Leu Pro Ala Ser Ile Cys  
 50 55 60

Ile Gln Leu Arg Ala Gly Asn  
 65 70

&lt;210&gt; 153

&lt;211&gt; 75

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 153

Ala Gly Ser Pro Ala Gly Thr Gly Pro Glu Phe Pro Gly Arg Pro Thr  
 1 5 10 15

Arg Pro Ile Ser Thr His Val Phe Glu Tyr Glu Cys Ile Cys Lys Ile  
 20 25 30

Pro Arg Phe Met Cys Glu Tyr Val Leu Leu Leu Tyr Ile Val Leu Leu  
 35 40 45

Cys Asn Arg Ser Tyr Ala Val Phe Thr Gln Cys Val Leu Arg Ser Ser  
 50 55 60

Pro Ile Asp Ser Ser Arg Asn Ala Val Leu Leu  
 65 70 75

&lt;210&gt; 154

&lt;211&gt; 483

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (194)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (205)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 154

Met Pro Ser Gly Met Ser Ala Ala Val Pro Ile Ser Gly Leu Leu Asp  
 1 5 10 15

Leu Ser His Asn Ser Ile Ser Gln Glu Ser Ala Leu Tyr Leu Leu Glu  
 20 25 30



98

Thr Leu Pro Ser Cys Pro Arg Val Arg Glu Ala Ser Val Asn Leu Gly  
 35 40 45  
 Ser Glu Gln Ser Phe Arg Ile His Phe Ser Arg Glu Asp Gln Ala Gly  
 50 55 60  
 Lys Thr Leu Arg Leu Ser Glu Cys Ser Phe Arg Pro Glu His Val Ser  
 65 70 75 80  
 Arg Leu Ala Thr Gly Leu Ser Lys Ser Leu Gln Leu Thr Glu Leu Thr  
 85 90 95  
 Leu Thr Gln Cys Cys Leu Gly Gln Lys Gln Leu Ala Ile Leu Leu Ser  
 100 105 110  
 Leu Val Gly Arg Pro Ala Gly Leu Phe Ser Leu Arg Val Gln Glu Pro  
 115 120 125  
 Trp Ala Asp Arg Ala Arg Val Leu Ser Leu Leu Glu Val Cys Ala Gln  
 130 135 140  
 Ala Ser Gly Ser Val Thr Glu Ile Ser Ile Ser Glu Thr Gln Gln Gln  
 145 150 155 160  
 Leu Cys Val Gln Leu Glu Phe Pro Arg Gln Glu Glu Asn Pro Glu Ala  
 165 170 175  
 Val Ala Leu Arg Leu Ala His Cys Asp Leu Gly Ala His His Ser Leu  
 180 185 190  
 Leu Xaa Gly Gln Leu Met Glu Thr Cys Ala Arg Leu Xaa Gln Leu Ser  
 195 200 205  
 Leu Ser Gln Val Asn Leu Cys Glu Asp Asp Ala Ser Ser Leu Leu  
 210 215 220  
 Leu Gln Ser Leu Leu Leu Ser Leu Ser Glu Leu Lys Thr Phe Arg Leu  
 225 230 235 240  
 Thr Ser Ser Cys Val Ser Thr Glu Gly Leu Ala His Leu Ala Ser Gly  
 245 250 255  
 Leu Gly His Cys His His Leu Glu Glu Leu Asp Leu Ser Asn Asn Gln  
 260 265 270  
 Phe Asp Glu Glu Gly Thr Lys Ala Leu Met Arg Ala Leu Glu Gly Lys  
 275 280 285  
 Trp Met Leu Lys Arg Leu Asp Leu Ser His Leu Leu Leu Asn Ser Ser  
 290 295 300  
 Thr Leu Ala Leu Leu Thr His Arg Leu Ser Gln Met Thr Cys Leu Gln  
 305 310 315 320  
 Ser Leu Arg Leu Asn Arg Asn Ser Ile Gly Asp Val Gly Cys Cys His  
 325 330 335  
 Leu Ser Glu Ala Leu Arg Ala Ala Thr Ser Leu Glu Glu Leu Asp Leu

340 " " " 99 345 350  
 Ser His Asn Gln Ile Gly Asp Ala Gly Val Gln His Leu Ala Thr Ile  
 355 360 365  
 Leu Pro Gly Leu Pro Glu Leu Arg Lys Ile Asp Leu Ser Gly Asn Ser  
 370 375 380  
 Ile Ser Ser Ala Gly Gly Val Gln Leu Ala Glu Ser Leu Val Leu Cys  
 385 390 395 400  
 Arg Arg Leu Glu Glu Leu Met Leu Gly Cys Asn Ala Leu Gly Asp Pro  
 405 410 415  
 Thr Ala Leu Gly Leu Ala Gln Glu Leu Pro Gln His Leu Arg Val Leu  
 420 425 430  
 His Leu Pro Phe Ser His Leu Gly Pro Gly Gly Ala Leu Ser Leu Ala  
 435 440 445  
 Arg Pro Trp Met Asp Pro Pro Ile Trp Lys Arg Ser Ala Trp Arg Lys  
 450 455 460  
 Thr Thr Trp Leu Glu Gly Ser Cys Val Ser Val Trp Ser Ser Arg Cys  
 465 470 475 480  
 Ser Asp Arg

<210> 155  
 <211> 221  
 <212> PRT  
 <213> Homo sapiens

<400> 155  
 His Gln Leu Ser Arg Gly Ser Ala Val Gly Arg Val Ser Arg Ser Leu  
 1 5 10 15  
 Gln Ala Pro Gly Gly Val Asp Ala Trp Leu Gln Cys Pro Gly Gly Ser  
 20 25 30  
 His Ser Pro Gly Ala Gly Ser Gly Ala Ala Pro Ala Pro Glu Gly Pro  
 35 40 45  
 Thr Pro Thr Ile Gln Pro Ser Gly Pro Arg Trp Gly Pro Glu Pro Gly  
 50 55 60  
 Gln Ala Leu Asp Gly Ser Pro His Leu Glu Glu Ile Ser Leu Ala Glu  
 65 70 75 80  
 Asn Asn Leu Ala Gly Gly Val Leu Arg Phe Cys Met Glu Leu Pro Leu  
 85 90 95  
 Leu Arg Gln Ile Asp Leu Val Ser Cys Lys Ile Asp Asn Gln Thr Ala  
 100 105 110  
 Lys Leu Leu Thr Ser Ser Phe Thr Ser Cys Pro Ala Leu Glu Val Ile  
 115 120 125

100

Leu Leu Ser Trp Asn Leu Leu Gly Asp Glu Ala Ala Ala Glu Leu Ala  
 130 135 140

Gln Val Leu Pro Gln Met Gly Arg Leu Lys Arg Val Asp Leu Glu Lys  
 145 150 155 160

Asn Gln Ile Thr Ala Leu Gly Ala Trp Leu Leu Ala Glu Gly Leu Ala  
 165 170 175

Gln Gly Ser Ser Ile Gln Val Ile Arg Leu Trp Asn Asn Pro Ile Pro  
 180 185 190

Cys Asp Met Ala Gln His Leu Lys Ser Gln Glu Pro Arg Leu Asp Phe  
 195 200 205

Ala Phe Phe Asp Asn Gln Pro Gln Ala Pro Trp Gly Thr  
 210 215 220

&lt;210&gt; 156

&lt;211&gt; 89

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 156

Glu Lys Leu Phe Cys Phe Glu Met Leu Leu Ile Cys Lys Phe Ser Pro  
 1 5 10 15

Asn Ser Val Pro Pro Glu Thr Cys Ala Ile Leu Asn Gln Gly Leu Met  
 20 25 30

Asp Leu Gly Leu Cys Arg Met Cys Leu Gly Asn Asn Met Phe Ala Gly  
 35 40 45

Ser Met Leu Gly Lys Ser His Arg His Ser Pro Phe Ser Ile Asn Gln  
 50 55 60

Arg His Asn Ala Leu Arg Lys Ala Ala Gly Thr Pro Ala Gln Lys Ser  
 65 70 75 80

Leu Gly Ile Val Gln Val Ser Pro Asn  
 85

&lt;210&gt; 157

&lt;211&gt; 58

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 157

Gly Cys Ala Gly Cys Ala Leu Val Thr Ile Cys Leu Gln Ala Val Cys  
 1 5 10 15

Leu Val Lys Ala Ile Ala Ile Leu His Ser Arg Leu Thr Arg Asp Thr  
 20 25 30

Met His Cys Gly Arg Pro Gln Gly Pro Leu Pro Arg Lys Ala Trp Val  
 35 40 45

101

Leu Ser Arg Phe Pro Pro Thr Glu Thr Ala  
50 55

<210> 158  
<211> 48  
<212> PRT  
<213> Homo sapiens

<400> 158  
Pro Glu Thr Gln Cys Thr Ala Glu Gly Arg Arg Asp Pro Cys Pro Glu  
1 5 10 15

Lys Pro Gly Tyr Cys Pro Gly Phe Pro Gln Leu Arg Gln Pro Glu Ile  
20 25 30

Trp Pro Arg Gly Lys Gly Lys Thr Leu His Pro Pro Ala Arg His Met  
35 40 45

<210> 159  
<211> 8  
<212> PRT  
<213> Homo sapiens

<400> 159  
Ser Glu Ile Gly Glu Asn Arg Pro  
1 5

<210> 160  
<211> 8  
<212> PRT  
<213> Homo sapiens

<400> 160  
His Asp Thr Asp Ser Phe Ala His  
1 5

<210> 161  
<211> 7  
<212> PRT  
<213> Homo sapiens

<400> 161  
Ala Leu Arg Lys Ala Ala Gly  
1 5

<210> 162  
<211> 148  
<212> PRT  
<213> Homo sapiens

102

&lt;400&gt; 162

Met Arg Gly Pro Val Cys Gly Phe Ser Leu Val Glu Met Leu Leu Ala  
 1 5 10 15

Leu Ala Leu Gly Leu Met Leu Ile Leu Gly Val Thr Gln Ile Ala Leu  
 20 25 30

Ser Ser Arg Thr Thr Tyr Ala Ser Gln Ser Ala Ala Ser Leu Leu Gln  
 35 40 45

Asp Asp Ala Arg Phe Ala Leu Gly Lys Leu Ile Gln Glu Ile Arg Gln  
 50 55 60

Ala Gly Met Phe Gly Cys Leu Ser Ala Ala Ser Ile Ser Asn Ala Pro  
 65 70 75 80

Ala Gly Phe Asp Arg Pro Ile Gly Trp Ser Thr Thr Gly Ser Ser Arg  
 85 90 95

Ser Leu Thr Leu Val Thr Ala Asp Val Gly Glu Gly Gly Ser Lys Pro  
 100 105 110

Asp Trp Thr Val Leu Ser Asp Cys Thr Gly Ser Ala His Ala Tyr Val  
 115 120 125

Gly Ser Pro Pro Ala Ala Asn Ala Arg Ala Asn Pro Leu Pro Thr Cys  
 130 135 140

Ala Lys Leu Thr  
 145

&lt;210&gt; 163

&lt;211&gt; 137

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 163

Met Gly Tyr Tyr Leu Ser Arg Ser Arg Gln Ala Gly Met Val Leu Leu  
 1 5 10 15

Ile Ser Leu Val Phe Leu Leu Leu Leu Ala Leu Leu Gly Val Ser Ser  
 20 25 30

Met Gln Gly Ala Ile Ser Gln Glu Lys Ile Thr Gly Ser Leu Arg Gln  
 35 40 45

Arg Asn Gln Ser Phe Gln Gln Ala Glu Ser Gly Leu Arg Leu Gly Glu  
 50 55 60

Ser Leu Val Gln Ala Ser Gly Phe Ala Leu Arg Pro Cys His Ser Thr  
 65 70 75 80

Ala Ala Cys Ala Pro Pro Ala Glu Ser Val Ser Val Val Gly Pro Gly  
 85 90 95

Thr Asn Pro Val Ser Thr Val Thr Trp Ile Gly Met Lys Asp Gly Val  
 100 105 110

103

Tyr Gly Ile Gln Asn Leu Gly Pro Gly Thr Gly Leu Val Asn Ser Arg  
 115 120 125

Gln Arg Pro Arg Pro Arg Ser Ile Ala  
 130 135

<210> 164  
 <211> 209  
 <212> PRT  
 <213> Homo sapiens

<400> 164  
 Glu Asn Glu Ser Thr Lys Glu Pro Ser Leu Leu Gln Tyr Leu Cys Val  
 1 5 10 15

Gln Ser Pro Ala Gly Leu Asn Gly Phe Asn Val Leu Leu Ser Gly Ser  
 20 25 30

Gln Thr Pro Pro Thr Val Gly Pro Ser Ser Gly Gln Leu Pro Ser Phe  
 35 40 45

Ser Val Pro Cys Met Val Leu Pro Ser Pro Pro Leu Gly Pro Phe Pro  
 50 55 60

Val Leu Tyr Ser Pro Ala Met Pro Gly Pro Val Ser Ser Thr Leu Gly  
 65 70 75 80

Ala Leu Pro Asn Thr Gly Pro Val Asn Phe Ser Leu Pro Gly Leu Gly  
 85 90 95

Ser Ile Ala Gln Leu Leu Val Gly Pro Thr Ala Val Val Asn Pro Lys  
 100 105 110

Ser Ser Thr Leu Pro Ser Ala Asp Pro Gln Leu Gln Ser Gln Pro Ser  
 115 120 125

Leu Asn Leu Ser Pro Val Met Ser Arg Ser His Ser Val Val Gln Gln  
 130 135 140

Pro Glu Ser Pro Val Tyr Val Gly His Pro Val Ser Val Val Lys Leu  
 145 150 155 160

His Gln Ser Pro Val Pro Val Thr Pro Lys Ser Ile Gln Arg Thr His  
 165 170 175

Arg Glu Thr Phe Phe Lys Thr Pro Gly Ser Leu Gly Asp Pro Val Leu  
 180 185 190

Lys Arg Arg Glu Arg Asn Asn His Glu Thr Pro Ala Arg Pro Arg Gly  
 195 200 205

Asp

<210> 165  
 <211> 454  
 <212> PRT

104

&lt;213&gt; Homo sapiens

&lt;400&gt; 165

```

Arg His Glu Arg His Glu Tyr Arg Arg Ala Leu Asp His Glu Glu Glu
 1           5           10           15

Ala Leu Ser Ser Gly Ser Val Gln Glu Ala Glu Ala Met Leu Asp Glu
          20           25           30

Pro Gln Glu Gln Ala Glu Gly Ser Leu Thr Val Tyr Val Ile Ser Glu
 35           40           45

His Ser Ser Leu Leu Pro Gln Asp Met Met Ser Tyr Ile Gly Pro Lys
 50           55           60

Arg Thr Ala Val Val Arg Gly Ile Met His Arg Glu Ala Phe Asn Ile
 65           70           75           80

Ile Gly Arg Arg Ile Val Gln Val Ala Gln Ala Met Ser Leu Thr Glu
          85           90           95

Asp Val Leu Ala Ala Ala Leu Ala Asp His Leu Pro Glu Asp Lys Trp
          100          105          110

Ser Ala Glu Lys Arg Arg Pro Leu Lys Ser Ser Leu Gly Tyr Glu Ile
          115          120          125

Thr Phe Ser Leu Leu Asn Pro Asp Pro Lys Ser His Asp Val Tyr Trp
          130          135          140

Asp Ile Glu Gly Ala Val Arg Arg Tyr Val Gln Pro Phe Leu Asn Ala
          145          150          155          160

Leu Gly Ala Ala Gly Asn Phe Ser Val Asp Ser Gln Ile Leu Tyr Tyr
          165          170          175

Ala Met Leu Gly Val Asn Pro Arg Phe Asp Ser Ala Ser Ser Ser Tyr
          180          185          190

Tyr Leu Asp Met His Ser Leu Pro His Val Ile Asn Pro Val Glu Ser
          195          200          205

Arg Leu Gly Ser Ser Ala Ala Ser Leu Tyr Pro Val Leu Asn Phe Leu
          210          215          220

Leu Tyr Val Pro Glu Leu Ala His Ser Pro Leu Tyr Ile Gln Asp Lys
          225          230          235          240

Asp Gly Ala Pro Val Ala Thr Asn Ala Phe His Ser Pro Arg Trp Gly
          245          250          255

Gly Ile Met Val Tyr Asn Val Asp Ser Lys Thr Tyr Asn Ala Ser Val
          260          265          270

Leu Pro Val Arg Val Glu Val Asp Met Val Arg Val Met Glu Val Phe
          275          280          285

Leu Ala Gln Leu Arg Leu Leu Phe Gly Ile Ala Gln Pro Gln Leu Pro
          290          295          300

```

105

Pro Lys Cys Leu Leu Ser Gly Pro Thr Ser Glu Gly Leu Met Thr Trp  
 305 310 315 320

Glu Leu Asp Arg Leu Leu Trp Ala Arg Ser Val Glu Asn Leu Ala Thr  
 325 330 335

Ala Thr Thr Thr Leu Thr Ser Leu Ala Gln Leu Leu Gly Lys Ile Ser  
 340 345 350

Asn Ile Val Ile Lys Asp Asp Val Ala Ser Glu Val Tyr Lys Ala Val  
 355 360 365

Ala Ala Val Gln Lys Ser Ala Glu Glu Leu Ala Ser Gly His Leu Ala  
 370 375 380

Ser Ala Phe Val Ala Ser Gln Glu Ala Val Thr Ser Ser Glu Leu Ala  
 385 390 395 400

Phe Phe Asp Pro Ser Leu Leu His Leu Leu Tyr Phe Pro Asp Asp Gln  
 405 410 415

Lys Phe Ala Ile Tyr Ile Pro Leu Phe Leu Pro Met Ala Val Pro Ile  
 420 425 430

Leu Leu Ser Leu Val Lys Ile Phe Leu Glu Thr Arg Lys Ser Trp Arg  
 435 440 445

Lys Pro Glu Lys Thr Asp  
 450

&lt;210&gt; 166

&lt;211&gt; 66

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 166

Lys Leu Leu Leu Thr Lys Val Glu Gln Lys Leu Glu Leu Ala Arg Leu  
 1 5 10 15

Gln Val Asp Thr Ser Gly Ser Lys Glu Phe Gly Thr Ser Gly Ile Pro  
 20 25 30

Ala Lys Cys Arg Phe Pro Lys Ile Phe Val Asn Thr Asp Asp Thr Tyr  
 35 40 45

Glu Glu Leu His Leu Ile Val Tyr Lys Val Thr Thr Val Phe Leu Pro  
 50 55 60

Ala Leu  
 65

&lt;210&gt; 167

&lt;211&gt; 79

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens



106.

&lt;400&gt; 167

Met Glu Pro Gln Leu Gly Pro Glu Ala Ala Ala Leu Arg Pro Gly Trp  
 1 5 10 15

Leu Ala Leu Leu Leu Trp Val Ser Ala Leu Ser Cys Ser Phe Ser Leu  
 20 25 30

Pro Ala Ser Ser Leu Ser Ser Leu Val Pro Gln Val Arg Thr Ser Tyr  
 35 40 45

Asn Phe Gly Arg Thr Phe Leu Gly Leu Asp Lys Cys Asn Ala Cys Ile  
 50 55 60

Gly Thr Ser Ile Cys Lys Lys Phe Phe Lys Glu Arg Asn Lys Ile  
 65 70 75

&lt;210&gt; 168

&lt;211&gt; 209

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 168

Gln Leu Pro Leu Trp Pro Ser Pro Ala Ser Val Gln Pro Arg Val Asp  
 1 5 10 15

Ser Gln Arg Ala Arg Gly Ser Pro Glu Pro Lys Met Glu Pro Gln Leu  
 20 25 30

Gly Pro Glu Ala Ala Ala Leu Arg Pro Gly Trp Leu Ala Leu Leu Leu  
 35 40 45

Trp Val Ser Ala Leu Ser Cys Ser Phe Ser Leu Pro Ala Ser Ser Leu  
 50 55 60

Ser Ser Leu Val Pro Gln Val Arg Thr Ser Tyr Asn Phe Gly Arg Thr  
 65 70 75 80

Phe Leu Gly Leu Asp Lys Cys Asn Ala Cys Ile Gly Thr Ser Ile Cys  
 85 90 95

Lys Lys Phe Phe Lys Glu Glu Ile Arg Ser Asp Asn Trp Leu Ala Ser  
 100 105 110

His Leu Gly Leu Pro Pro Asp Ser Leu Leu Ser Tyr Pro Ala Asn Tyr  
 115 120 125

Ser Asp Asp Ser Lys Ile Trp Arg Pro Val Glu Ile Phe Arg Leu Val  
 130 135 140

Ser Lys Tyr Gln Asn Glu Ile Ser Asp Arg Lys Ile Cys Ala Ser Ala  
 145 150 155 160

Ser Ala Pro Lys Thr Cys Ser Ile Glu Arg Val Leu Arg Lys Thr Glu  
 165 170 175

Arg Phe Gln Lys Trp Leu Gln Ala Lys Arg Leu Thr Pro Asp Leu Val  
 180 185 190

107

Gln Asp Cys His Gln Gly Gln Arg Glu Leu Lys Phe Leu Cys Met Leu  
 195 200 205

Arg

<210> 169  
 <211> 146  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (96)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (107)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (111)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (115)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (122)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (132)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 169  
 Met Glu Pro Gln Leu Gly Pro Glu Ala Ala Ala Leu Arg Pro Gly Trp  
 1 5 10 15

Leu Ala Leu Leu Leu Trp Val Ser Ala Leu Ser Cys Ser Phe Ser Leu  
 20 25 30

Pro Ala Ser Ser Leu Ser Ser Leu Val Pro Gln Val Arg Thr Ser Tyr  
 35 40 45

Asn Phe Gly Arg Thr Phe Leu Gly Leu Asp Lys Cys Asn Ala Cys Ile  
 50 55 60

Gly Thr Ser Ile Cys Lys Lys Phe Phe Lys Glu Glu Ile Arg Ser Asp  
 65 70 75 80

Asn Trp Leu Ala Ser His Leu Gly Thr Ala Ser Arg Phe Pro Leu Xaa  
 85 90 95

108

Ser Tyr Pro Cys Lys Leu Leu Gln Met Ile Xaa Lys Ile Trp Xaa Pro  
100 105 110

Cys Gly Xaa Leu Leu Thr Gly Gln Gln Xaa Ser Asn Glu Ile Ser Lys  
115 120 125

Gln Glu Ile Xaa Cys Leu Leu His Pro Pro Pro Lys Asn Leu His Ile  
130 135 140

Asp Val  
145

<210> 170

<211> 97

<212> PRT

<213> Homo sapiens

<400> 170

Gly Pro Arg Ala Arg Val Gln Gly Phe Ser Gly Ala Asp Ile Val Lys  
1 5 10 15

Phe Met Ala Leu Gly Ser Met Tyr Leu Val Leu Thr Leu Ile Val Ala  
20 25 30

Lys Val Leu Arg Gly Ala Glu Pro Cys Cys Gly Pro Leu Lys Asn Arg  
35 40 45

Val Leu Arg Pro Cys Pro Leu Pro Val His Cys Pro Leu Pro Ile Pro  
50 55 60


Ser Pro Ala Glu Gly Ile Pro Trp Val Ala Tyr Leu Pro Ile Arg Trp  
65 70 75 80

Phe Ile Ser Cys Cys Pro Gly His Cys Ile Gln Ile Pro Met Cys Thr  
85 90 95

Ser

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/19330

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) : CO7H 21/04; C07K 1/00; C12N 1/20; C12P 21/06 US CL : 530/350; 536/23.5; 435/252.3; 435/69.1 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 530/350; 536/23.5; 435/252.3; 435/69.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Sequence databases: emb158, genbank111, n-issued, n-geneseq35, emb1-e3st58, agenseq35, a-issued, pir60		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database Genebank, accession U66676, Homo sapiens cDNA similar to ATP-binding cassette transporter. 12 September 1996, Vol. 5, No. 10, pages 1649-1655, see attached sequence alignment.	1-10, 21
X	Database Genbank, accession N46602, Homo sapiens cDNA clone IMAGE: 276822, April 1993, see attached sequence alignment.	1-10, 21
Y	US 5,352,596 A (CHEUNG et al) 04 October 1994, col 35, line 1, and see attached sequence alignment.	11,12, 14-16
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family	
Date of the actual completion of the international search 01 DECEMBER 1999		Date of mailing of the international search report 7 fevrier 2000    07/02/2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer  MICHAEL T. BRANNOCK Telephone No. (703) 308-0196